

**Notes on aquatic and semiaquatic bugs
(Hemiptera: Heteroptera: Nepomorpha, Gerromorpha)
from Malesia with description of three new species**

Ping-ping CHEN¹⁾, Nico NIESER²⁾ & Ivor LANSBURY³⁾

¹⁾ Section of Entomology, Plant Protection Service, P.O.Box 9102, NL-6700 HC, Wageningen & c/o Dept. of Entomology, National Museum of Natural History – Naturalis, P.O.Box 9517, NL-2300 RA, Leiden, The Netherlands; e mail: p.chen@minlnv.nl

²⁾ c/o Dept. of Collections, National Museum of Natural History – Naturalis, P.O.Box 9517, NL-2300 RA, Leiden, The Netherlands; e-mail: nico.nieser@hotmail.com

³⁾ 10 Chilswell Road, Oxford OX1 4PJ, England, U.K.

Abstract. *Micronecta lumutensis* sp. nov. (Micronectidae) from Kalimantan (Borneo, Indonesia), *Enithares rinjani* sp. nov. (Notonectidae) from Lombok (Indonesia) and *Neusterensifer stysi* sp. nov. (Veliidae) from New Guinea are newly described. In addition, distributional records are given for three species of the genus *Enithares* Spinola, 1937, *Neusterensifer sepik* (Polhemus & Polhemus, 1994), and *Tenagogonus kampaspe* (Kirkaldy, 1900). Morphological variability in *T. kampaspe* is discussed.

Key words. Heteroptera, Gerridae, *Tenagogonus*, Micronectidae, *Micronecta*, Notonectidae, *Enithares*, Veliidae, *Neusterensifer*, new species, new records, Indonesia, Papua New Guinea

Introduction

While sorting out material in the Leiden and Oxford Museums we came across some specimens belonging to new or rarely reported species of aquatic and semiaquatic bugs which are described or reported in this paper.

In the Old World, the family Micronectidae is, apart from *Synaptonecta* Lundblad, 1933 (two species) and *Papuanecta* Tinerella, 2008 (four species, see TINERELLA (2008)), dominantly represented by the genus *Micronecta* Kirkaldy, 1897, which has numerous species in the Old World tropics, especially on the Asian continent, and the richness of species is gradually declining eastward except for a secondary centre of diversification in Australia (WRÓBLEWSKI 1968, 1972; ANDERSEN & WEIR 2004). CHEN et al. (2005) report 32 species of *Micronecta* from Malesia. However, the micronectine fauna of Borneo is poorly known, with only three reported species. As a comparison, eight species have been found in Sulawesi. In this paper, the fourth species of *Micronecta* is added to Kalimantan.