

## Two new species of *Kokeshia* from India and Thailand (Hemiptera: Heteroptera: Schizopteridae)

Dávid RÉDEI

Hungarian Natural History Museum, Department of Zoology, H-1088 Budapest, Baross u. 13, Hungary;  
e-mail: redei@zoo.zoo.nhmus.hu

**Abstract.** *Kokeshia stysi* sp. nov. (India) and *K. oroszi* sp. nov. (Thailand) (Heteroptera: Schizopteridae) are described. A key for the identification of the known species of *Kokeshia* is presented.

**Key words.** Heteroptera, Dipsocoromorpha, Schizopteridae, *Kokeshia*, new species, India, Thailand

### Introduction

The schizopterid genus *Kokeshia* was established by MIYAMOTO (1960) to accommodate his new species *K. esakii* Miyamoto, 1960, from Japan. Shortly after that, the morphology of the alimentary tract of *K. esakii* was studied by the same author (MIYAMOTO 1961). ŠTYS (1985) described two further new species, *K. martensi* Štys, 1985, and *K. similis* Štys, 1985, both from Nepal, provided an extensive morphological treatment of *K. martensi*, and discussed the diagnostic characters as well as the evolutionary significance of the genus. An additional species, *K. hsiao* Ren & Zheng, 1992, was described by REN & ZHENG (1992) from Zhejiang Province, eastern China. *Kokeshia* currently contains the above four species, and no other authors have ever provided original data on the genus.

Among unidentified material deposited in the Hungarian Natural History Museum (HNHM), Budapest, specimens representing two undescribed species of *Kokeshia* were found. These are described as new in the present paper.

### Materials and methods

Structures were examined using an optical microscope (XSZ-N107). Drawings were made using a camera lucida. Measurements were taken using a micrometer eyepiece at magnifications up to 400x. Male genitalia were examined after a short clearing in 10% KOH solution. The terminology follows ŠTYS (1985) except parts of the phallus.