

## Chrysoidea: Chrysididae (zlatěnkovití)

Pavel TYRNER

Tylova 2073, CZ-436 01 Litvínov, Czech Republic; e-mail: ptyrner@seznam.cz

**Abstract.** This paper presents a checklist of Chrysididae of the Czech Republic and Slovakia. So far 98 species are known from the Czech Republic (81 from Bohemia, 91 from Moravia), and 124 species from Slovakia. *Chrysis valida* Mocsáry, 1912 (Bohemia and Moravia), *C. leptomandibularis* Niehuis, 2000 (Bohemia and Moravia), *C. phryne* Abeille de Perrin, 1878 (Moravia), and *C. subsinuata* Marquet, 1879 (Moravia) are new for the Czech Republic; *C. grohmanni* Dahlbom, 1854 and *C. chryso stigma* Mocsáry, 1889 are new for Bohemia; *Hedychridium krajniki* Balthasar, 1946, *Chrysis angustula* Schenck, 1856, and *C. pseudobrevitarsis* Linsenmaier, 1951 are new for Moravia. *Cleptes splendens* (Fabricius, 1793), *Pseudomalus triangulifer* Abeille de Perrin, 1877, *Hedychridium mediocre* Linsenmaier, 1987, *Chrysis cerastes* Abeille de Perrin, 1877, *C. equestris* Dahlbom, 1845, *C. leptomandibularis*, *C. pseudobrevitarsis* Linsenmaier, 1951, *C. pyrrhina* Dahlbom, 1845, *C. separata* Trautmann, 1926, *C. valida*, and *Chrysura filiformis* Mocsáry, 1889 are new for Slovakia. Several problems in synonymy are explained. *Trichrysis pumilionis* Linsenmaier, 1987 is accepted as a new valid name for *Chrysidea pumila* (Klug, 1845).

**Keywords:** Hymenoptera, Chrysoidea, Chrysididae, checklist, new records, Czech Republic, Bohemia, Moravia, Slovakia

### Introduction

The family Chrysididae is distributed all over the World and contains more than 3,000 species; moreover, about 1,000 species are expected to be found and described (KIMSEY & BOHART 1990). The family is divided into four subfamilies, two of them (Cleptinae and Chrysidinae) occur in central Europe. The species rich subfamily Chrysidinae contains tribes Allosceliini, Elampini, Chrysidini, and

Čeďed' zlatěnkovití (Chrysididae) je rozšířena po celém světě a čítá více než 3 000 druhů, přičemž okolo 1 000 druhů bude pravděpodobně ještě objeveno a popsáno (KIMSEY & BOHART 1990). Děľíme ji do čtyř podčeledí, z nichž dvě (Cleptinae a Chrysidinae) žijí i ve střední Evropě. Druhově početná podčeleď Chrysidinae zahrnuje triby Allosceliini, Elampini, Chrysidini a Parnopini a kromě