

Vespoidea: Pompilidae (hrabalkovití)

Jakub STRAKA

Charles University in Prague, Faculty of Science, Department of Zoology, Viničná 7, CZ-128 44 Praha 2, Czech Republic; e-mail: straka-jakub@mbox.vol.cz

Abstract. Checklist of Pompilidae of the Czech Republic and Slovakia is presented: 100 species are known from the Czech Republic (85 from Bohemia, 92 from Moravia), and 99 from Slovakia. *Anoplius alpinobalticus* Wolf, 1965 and *Priocnemis fastigiata* Haupt, 1934 are new species for Bohemia, *Arachnospila conjungens* (Kohl 1898), *Arachnospila rufa* (Haupt, 1927) and *Dipogon monticolus* Wahis, 1972 are new species for Slovakia. Occurrence of the following rare species was confirmed: *Priocnemis pelliplus* Wahis, 1998, *Poecilagenia rubicans* (Lepelletier, 1845), *Arachnospila westerlundii* (Morawitz, 1893), *Tachyagetes filicornis* (Tournier, 1889), and *Anoplius tenuicornis* (Tournier, 1889) for the Czech Republic, *Dipogon vechti* Day, 1979 and *Auplopus rectus* (Haupt, 1926) for Bohemia, *Aporus pollux* (Kohl, 1888), *Dicyrtomellus tingitanus* (Wolf, 1966), and *Priocnemis pelliplus* for Slovakia. The following species have been conditionally removed from the Slovak fauna until the material is revised: *Priocnemis baltica* Blüthgen, 1944, *P. bellieri* Sichel, 1860, *P. diversa* Junco, 1947, *P. fallax* Verhoeff, 1922, *P. massaliensis* Soyer, 1945, *Arachnospila consobrina* (Dahlbom, 1843), *Evagetes sahlbergi* (Morawitz, 1893), *Anoplius tenuicornis* (Tournier, 1889), and *Episyron funerarium* (Tournier, 1889). Majority of these species are highly unlikely to occur in Slovakia.

Key words. Hymenoptera, Vespoidea, Pompilidae, checklist, new records, Czech Republic, Bohemia, Moravia, Slovakia

Introduction

The family Pompilidae (spider wasps) comprises about 120 described genera and about 5000 described species (WASBAUER 1995). It is divided into three cosmopolitan subfamilies (Pepsinae, Pompilinae, and Ceropalinae), whose members can be found almost

Čeď hrabalkovití (Pompilidae) je ve světě zastoupena přibližně 120 rody se zhruba 5000 dosud popsanými druhy (WASBAUER 1995). Dělíme ji na tři podčeďedi (Pepsinae, Pompilinae a Ceropalinae), které jsou rozšířeny kosmopolitně a s jejich zástupci