


**ENUMERATIO INSECTORUM  
BOHEMOSLOVAKIAE  
CHECK LIST**  
**of Czechoslovak Insects III (Hymenoptera)**

of Czechoslovak Insects III (Hymenoptera)

Prepared cooperatively by specialists  
on the various groups of Hymenoptera  
under the direction of

JOSEF ŠEDIVÝ



ACTA FAUNISTICA  
ENTOMOLOGICA  
MUSEI NATIONALIS PRAGAE

19 — 1989

Národní muzeum — Přírodovědecké muzeum, Praha

Sectio entomologica Musei rerum naturalium, Praha-Kunratice, Československo

Acta faun. ent. Mus. Nat. Pragae

VYDÁNO -- EDITUM

30. XI. 1989

NÁRODNÍ MUZEUM

PRAHA

Vedoucí redaktor: RNDr. Jiří Dlabač, CSc.

Redakční rada:

RNDr. Adolf Čejchan, CSc., RNDr. Ludvík Hoberlandt, CSc.,  
RNDr. Josef Jelínek, CSc., RNDr. Jan Ježek, CSc.,  
RNDr. František Krampl, CSc.

Recenzent: RNDr. Pavel Štys, CSc.

Návrh obálky: redakce AF, podle RNDr. Josefa Šedivého, DrSc.

Jazyková úprava: dr. ing. Radoslav Obrtel, CSc.

## CONTENTS

	Page
Introduction by J. Šedivý . . . . .	6
Explanations by J. Šedivý . . . . .	7
Hymenoptera by J. Šedivý . . . . .	9
Symphyta by K. Beneš . . . . .	14
Xyeloidea . . . . .	14
Xyelidae . . . . .	14
Pamphilioidea . . . . .	14
Pamphiliidae . . . . .	14
Megalodontidae . . . . .	15
Tenthredinoidea . . . . .	15
Argidae . . . . .	15
Blasticotomidae . . . . .	15
Cimbicidae . . . . .	15
Diprionidae . . . . .	16
Tenthredinidae . . . . .	16
Siricoidea . . . . .	24
Siricidae . . . . .	24
Xiphidiidae . . . . .	24
Orussoidea . . . . .	25
Orussidae . . . . .	25
Cephidoidea . . . . .	25
Cephidae . . . . .	25
Apocrita . . . . .	27
Parasitica . . . . .	27
Ichneumonoidea . . . . .	27
Braconidae by M. Čapek and J. Lukáš . . . . .	27
Aphidiidae by B. Starý . . . . .	45
Ichneumonidae by J. Šedivý . . . . .	49
Stephanoidea by J. Šedivý . . . . .	95
Stephanidae . . . . .	95
Chalcidoidea by V. Kalina . . . . .	97
Leucospidae . . . . .	98
Chalcididae . . . . .	98
Eurytomidae . . . . .	99
Torymidae . . . . .	100
Ormymidae . . . . .	102
Eucharitidae . . . . .	102

Perilampidae . . . . .	102
Pteromalidae . . . . .	102
Eupelmidae . . . . .	111
Encyrtidae . . . . .	112
Signiphoridae . . . . .	118
Aphelinidae . . . . .	118
Elasmidae . . . . .	119
Tetracampidae . . . . .	119
Eulophidae . . . . .	119
Trichogrammatidae . . . . .	126
Mymaridae . . . . .	127
Mymarommatidae . . . . .	127
Cynipoidea by P. Mikula . . . . .	129
Ibaliidae . . . . .	130
Anacharitidae . . . . .	130
Aspiceridae . . . . .	130
Figitidae . . . . .	130
Eucolidae . . . . .	130
Alloxystidae . . . . .	130
Cynipidae . . . . .	131
Evanoidea by J. Šedivý . . . . .	133
Evanidae . . . . .	133
Aulacidae . . . . .	133
Gasteruptiidae . . . . .	134
Proctotrupoidea by J. Macek . . . . .	136
Heloridae . . . . .	136
Proctotrupidae . . . . .	136
Diapriidae . . . . .	136
Scelionidae . . . . .	139
Platygasteridae . . . . .	140
Ceraphronoidea . . . . .	141
Ceraphronidae . . . . .	141
Megaspilidae . . . . .	141
Trigonaloidea by J. Šedivý . . . . .	142
Trigonalidae . . . . .	142
Aculeata . . . . .	143
Chrysidoidea by J. Strejček . . . . .	143
Dryinidae . . . . .	144
Embolemidae . . . . .	145
Bethylidae . . . . .	145
Cleptidae . . . . .	146
Chrysididae by Z. Pádr . . . . .	147
Scolioidea by Z. Pádr . . . . .	149
Tiphidae . . . . .	150
Methochidae . . . . .	150
Mutillidae . . . . .	150
Scoliidae . . . . .	151
Sapygidae . . . . .	151

<b>Formicoidea by P. Werner</b>	.	.	.	.	.	.	.	.	153
<b>Formicidae</b>	.	.	.	.	.	.	.	.	154
<b>Vespoidea by Z. Pádr</b>	.	.	.	.	.	.	.	.	157
<b>Masaridae</b>	.	.	.	.	.	.	.	.	157
<b>Eumenidae</b>	.	.	.	.	.	.	.	.	157
<b>Vespidae</b>	.	.	.	.	.	.	.	.	
<b>Pompiloidea by J. Šedivý</b>	.	.	.	.	.	.	.	.	161
<b>Pompilidae</b>	.	.	.	.	.	.	.	.	161
<b>Sphecoidea by Z. Pádr</b>	.	.	.	.	.	.	.	.	165
<b>Sphecidae</b>	.	.	.	.	.	.	.	.	166
<b>Ampulicidae</b>	.	.	.	.	.	.	.	.	166
<b>Philantidae</b>	.	.	.	.	.	.	.	.	166
<b>Nyssonidae</b>	.	.	.	.	.	.	.	.	166
<b>Larridae</b>	.	.	.	.	.	.	.	.	167
<b>Astatidae</b>	.	.	.	.	.	.	.	.	168
<b>Pemphredonidae</b>	.	.	.	.	.	.	.	.	168
<b>Trypoxylonidae</b>	.	.	.	.	.	.	.	.	169
<b>Crabronidae</b>	.	.	.	.	.	.	.	.	169
<b>Apoidea by M. Kocourek</b>	.	.	.	.	.	.	.	.	173
<b>Colletidae</b>	.	.	.	.	.	.	.	.	174
<b>Andrenidae</b>	.	.	.	.	.	.	.	.	174
<b>Halictidae</b>	.	.	.	.	.	.	.	.	176
<b>Melittidae</b>	.	.	.	.	.	.	.	.	178
<b>Megachilidae</b>	.	.	.	.	.	.	.	.	179
<b>Anthophoridae</b>	.	.	.	.	.	.	.	.	181
<b>Apidae</b>	.	.	.	.	.	.	.	.	184
<b>Index by J. Šedivý</b>	.	.	.	.	.	.	.	.	185

## INTRODUCTION

The present check-list of insects of the order Hymenoptera is the first comprehensive evaluation of their occurrence in the territory of Czechoslovakia. It is the result of long-lasting faunal research carried out by professional entomologists and members of the Czechoslovak Entomological Society, sponsored by the Czechoslovak Academy of Sciences.

The list is based on materials deposited in state and private collections and on data published in the literature. The different superfamilies and families have been worked out at different levels. Those groups which are being studied actively in Czechoslovakia have been worked out to a greater depth, in contrast to those in which faunal studies are not under way at present. In those cases the list is based on findings obtained and determined one or two decades ago. This is why even the classification of the different superfamilies is different. In some of them, subfamilies, tribes, subgenera and subspecies are not included. Many taxa ranked as subspecies in the present list and recorded as such in the literature are actually not biological subspecies (ecological or geographical ones) at all; their names concern infrasubspecific taxa. In the present list it has not been possible to consider such differences, and all infraspecific taxa have been listed as subspecies.

The check-list contains only synonyms published in recent years or in such cases where a particular species was reported from Czechoslovakia under an invalid name.

The introductory chapters to superfamilies and families comprise data on the present status of taxonomy and faunal research of the particular groups. A brief outline of the history of faunal research in Czechoslovakia is included, stating places where collections of the entomologists mentioned are deposited. Biological data are included in a general way, not encompassing specific data on individual taxa. The introductory chapters are concluded by a statement on the level at which the occurrence in different parts of Czechoslovakia is known. Concerning taxa higher than subfamily, essentially the classification used in the Catalog of Hymenoptera of America North of Mexico, I-III, (Krombein et al. 1979, Washington, 2735 pp.) has been adopted. The basic works

according to which lower taxonomical units have been arranged are given in the introductory chapters to superfamilies or families.

The main objective of compiling this check-list has been to provide an idea of the richness of hymenopterous fauna of Czechoslovakia. Some of the species listed are in danger of extinction due to large-scale alterations of their habitats. In compiling the list, one of the major endeavours has been to eliminate erroneous data on the occurrence of certain species in Czechoslovakia. The inclusion of the most important synonyms should facilitate orientation in the earlier literature.

The list corresponds with the present knowledge of the distribution of the Hymenoptera in Czechoslovakia. It can be assumed to be incomplete and to contain inaccuracies and/or errors, which to remove requires obtaining further new data on the individual species.

The names of entomologists responsible for the individual parts of the check-list are given in the headings of the superfamilies or families. Their addresses are given at the end of the paper.

The paper was submitted to press in March, 1986. After that date, only minor corrections were made in galley-proof reading.

I thank Dr. Ing. R. Obrtel, CSc., for emendations and/or translation of the texts into English.

Prague, March 1986

Josef Šedivý

#### EXPLANATIONS of the layout, abbreviations and symbols

##### Layout

All names of taxa from suborders down to tribi are listed according to the system and printed in CAPITAL LETTERS on the top left-hand side of each page. Within each genus or subgenus, the names of species and subspecies are listed alphabetically and are printed in low-case letters on the extreme left margin of the text. Synonyms of all taxa are indented.

##### Abbreviations and symbols used in the text:

B = Bohemia; M = Moravia incl. Silesia; S = Slovakia

\* = finding hitherto not published for a particular region

? = data in the literature (mostly foreign) pertaining to Czechoslovakia

auct. = auctorum

p.p. = pro parte

s. str. = sensu stricto

( ) = Author's name and year given in parentheses indicate that the species has been included in a genus different from that in which it was originally described

misident. = misidentifications

introd. = introduction



## HYMENOPTERA

JOSEF ŠEDIVÝ

The Hymenoptera is the most numerous insect order (in terms of species) in the territory of Czechoslovakia. Most numerous data on the occurrence and distribution are available on the groups Symphyta and Aculeata. In faunal respect, most of the families and some of the genera in these groups have been elaborated in the "Prodromus Hymenopterorum Čechoslovakiae/Bohemoslovakiae" [Pars 1—10, 1937—1971] and in the respective monographs of the series "Fauna ČSR" [Chrysididae, Sphecidae], as well as in faunal papers the most important of which are cited in the introductory chapters to superfamilies and families. A review of the genera and selected species recorded in Czechoslovakia has been given in the "Klíč zvířeny ČSR, II" (Key to the Fauna of Czechoslovakia, II, 1957, Bouček et al., in Kratochvíl, Prague: 35—406).

Fewer data are available on the distribution of Parasitica, the most numerous group of Hymenoptera, a comprehensive elaboration of whose difficult families, subfamilies and genera is still lacking. Further investigations, carried out in our territory, and further taxonomical studies may provide data on the distribution of additional species or of those new for science.

The order Hymenoptera comprises many economically important phytophagous pests and useful predators as well as parasitoids of arthropods, and pollinators of cultivated plants.

The larval stages of Symphyta, being the oldest hymenopterous groups evolutionarily, are phytophagous, xylophagous or seed-eating rarely parasitic. Most species of Hymenoptera are comprised in the group of Parasitica. Their larvae are predators of arthropods or their internal or external parasitoids. The Parasitica also comprises phytophagous species whose larvae produce plant galls (Cynipidae), bore tunnels in plant stems (*Harmolita*) or consume seed endosperm (*Torymus*, *Systole*). Many species are hyperparasitic Mesochorinae, Ceraphronidae etc.).

Members of the group of Aculeata are often important pollinators of entomophilous plants. Their larvae feed on nectar and pollen. The Aculeata also comprise species cleptoparasitic on pollen-collecting solitary bees. Larvae of other aculeate species develop on arthropods paralysed by the female's sting.

Data on the total number of genera and species recorded in Czechoslovakia are given in tab. 1, not including subgenera, misidentified species, subspecies and varieties. In tab. 2 the superfamilies are ranged by

the numbers of species and their percentages of the total number of species of Hymenoptera so far recorded in Czechoslovakia.

Table 1: Numbers of Hymenoptera taxa in Czechoslovakia

	Gene- ra	Spe- cies		Gene- ra	Spe- cies
HYMENOPTERA	1599	7483	Cynipidae	20	162
Symphyta	120	617	Evanoidea	5	24
Xyeloidea	2	6	Evaniidae	2	2
Xyelidae	2	6	Aulacidae	2	5
Pamphilioidea	7	46	Gasteruptiidae	1	17
Pamphiliidae	5	40	Proctotrupoidea	97	300
Megalodontidae	2	6	Heloridae	1	4
Tenthredinoidea	100	537	Proctotrupidae	8	28
Argidae	3	24	Diapriidae	40	176
Blasticotomidae	1	1	Scelionidae	25	52
Cimbicidae	7	24	Platygastridae	23	40
Diprionidae	6	16	Ceraphronoidea	9	20
Tenthredinidae	83	472	Ceraphronidae	5	13
Siricoidea	5	11	Megaspilidae	4	7
Siricidae	4	8	Trigonaloidea	1	1
Xiphydriidae	1	3	Trigonalidae	1	1
Orussoidea	1	1	Aculeata	255	1546
Orussidae	1	1	Chrysoidea	52	214
Cephidoidea	5	16	Dryinidae	17	48
Cephidae	5	16	Embolemidae	1	1
Apocrita	1479	6865	Bethylidae	20	45
Parasitica	1224	5320	Cleptidae	1	9
Ichneumonoidea	612	3088	Chrysidae	13	112
Braconidae	168	1005	Scolioidea	18	40
Aphidiidae	27	125	Tiphidae	2	9
Ichneumonidae	420	1957	Methochidae	1	1
Stephanoidea	1	1	Mutillidae	10	20
Stephanidae	1	1	Scoliidae	2	5
Chalcidoidea	461	1674	Sapygidae	3	5
Leucospidae	1	4	Formicoidea	32	108
Chalcididae	15	40	Formicidae	32	108
Eurytomidae	9	60	Vespoidea	28	89
Torymidae	16	107	Masaridae	1	1
Ormyridae	1	8	Eumenidae	23	72
Eucharitidae	2	4	Vespidae	4	16
Perilampidae	1	16	Pompiloidea	25	109
Pteromalidae	175	474	Pompilidae	25	109
Eupelmidae	7	40	Sphecoidea	45	287
Encyrtidae	115	347	Sphecidae	3	16
Signiphoridae	3	3	Ampulicidae	2	2
Aphelinidae	13	54	Philantidae	2	25
Elasmidae	2	8	Nyssonidae	9	59
Tetracampidae	5	8	Larridae	9	28
Eulophidae	69	449	Astatidae	1	11
Trichogrammatidae	15	25	Pemphredonidae	8	47
Mymaridae	10	26	Trypoxylonidae	2	9
Mymarommataidae	1	1	Crabronidae	9	90
Cynipoidea	38	212	Apoidae	55	699
Ibaliidae	1	2	Colletidae	2	44
Anacharitidae	1	6	Andrenidae	3	162
Asiceridae	1	1	Halictidae	8	133
Figitidae	4	13	Melittidae	3	13
Eucoilidae	9	18	Megachilidae	19	132
Alloxystidae	2	10	Anthophoridae	17	179
			Apidae	3	36

**Table 2: Superfamilies ranked by percent of total species of Hymenoptera**

	Number of species	Percent of total species
Ichneumonoidea	3088	41,27
Chalcidoidea	1674	22,37
Apoidea	699	9,34
Tenthredinoidea	537	7,17
Proctotrupoidea	300	4,00
Sphecoidea	287	3,83
Chrysidoidea	214	2,84
Cynipoidea	212	2,83
Pompiloidea	109	1,45
Formicoidea	108	1,44
Vespoidea	89	1,18
Pamphilioidea	46	0,61
Scolioidea	40	0,53
Evanoidea	24	0,32
Ceraphronoidea	20	0,26
Cephoidea	16	0,21
Siricoidea	11	0,14
Xyeloidea	6	0,08
Orussoidea	1	0,01
Stephanoidea	1	0,01
Trigonaloidea	1	0,01
	7483	99,90



## S Y M P H Y T A

KAREL BENEŠ

The very beginnings of the research into sawflies in our countries coincide with the research history of other groups of Hymenoptera. Among the first collectors of Symphyta in our territory who contributed to our present knowledge since the middle of the last century, the most important were L. Kirchner, followed later by H. Gradiš, R. v. Stein, G. Strobl and particularly J. Slavíček of Milkov (coll. Moravian Museum, Brno). At the beginning of the century, the most important collectors were A. Kubes, L. Duda, and cecidologists E. Bayer and E. Baudyš; the latter published, for more than fifty years, since 1914 records of sawfly galls in a number of papers.

Between World Wars I and II, F. Gregor sen. founded a collection which is the basis of the collection of Symphyta in the National Museum, Praha. To the same time belong collections and/or faunal papers by E. Hachler, L. Bařa, O. Šuster, A. Hoffer, V. Balthazar and J. Obenberger. On the basis of these collections a list of Symphyta of the Czech countries was published [Gregor and Bařa 1940—42]. After the World War II, the Symphyta were collected by R. Obrtel, M. Kocourek and particularly by Z. Pádr.

The first faunal records from Slovakia, based on the collections of the Hungarian National Museum in Budapest, are found in Mocsáry (1897). In this century, F. Gregor sen. collected there and in the southern parts M. Kocourek and Z. Pádr; those collections of the latter two are now deposited in the Slovakian National Museum, Bratislava. In the High Tatra Mts., L. H. Woollatt collected in 1970 about fifty species which are now deposited in the Hope Museum, Oxford. In recent years, J. Jendeková contributed to the knowledge of the fauna of southern Slovakia.

Our knowledge of the Czechoslovak sawflies was summarized by Bouček and Pádr (1957). A number of papers on forest pests belonging to the Symphyta was published by S. Kolubajiv, A. Kalandra, M. Kudela and recently namely by V. Martinek and J. Kríštek. The present author occasionally dealt, within the last 25 years, with the taxonomy and life history and recently also with the faunistics of Symphyta in the Krkonoše Mts.

The suborder Symphyta includes generalized phytophagous Hymenoptera. Their larvae (except for Orussidae, parasitoids of wood-boring larvae of Buprestidae and Sircidae) feed mostly externally on the foliage of different plants but several groups develop in galls, leaf-mines

and fruits or bore in wood or stems. Many species are serious forest or agricultural pests.

The classification of the higher taxa in this list is based mainly on Benson (1951—58) and Smith (1979), except for the superfamily Xyeloidae which is considered as a separate group (sensu Rasnitsyn 1969).

References: Benson R. B., 1951—58: Hymenoptera Symphyta. Handbooks for the Identification of British Insects, 6 (2a—c), 252 pp. R. Ent. Soc. London. — Bouček Z. and Pádr Z. (in Kratochvíl J. ed.), 1957: Šíropasí — Symphyta. Klíč zvěřeny ČSR, 2. pp. 46—87. Nakl. ČSAV, Praha. — Gregor F. and Bata I., 1940—42: Podřád Symphyta. Prodromus Hymenopterorum patriae nostrae, 4—5. Sbor. ent. odd. Zem. mus., Praha, 18: 203—240; 19: 191—215; 20: 259—344. — Mocsáry A., 1897: Hymenoptera. Fauna Regni Hungariae, pp. 520—532. Budapest. — Rasnitsyn A. P., 1969: Proischožděníje i evoljucijsa nižšich perepončatokrylych. 196 pp. Nauka, Moskva. Smith D. R. (in Krombein K. et al.), 1979: Symphyta. Catalog of Hymenoptera in America North of Mexico, 1. pp. 3—137. Smith. Inst. Press, Washington, D. C.

SYMPHYTA		<i>nemoralis</i> (Thomson, 1871)	
XYELOIDEA		<i>pumilionis</i> (Giraud, 1861)	S
XYELIDAE		<b>Cephalcia</b> Panzer, 1805	
XYELINAE		<i>abietis</i> (Linnaeus, 1758)	B M S
XYELINI		<i>alashanica europaea</i>	
<b>Xyela</b> Dalman, 1819		<i>Beneš</i> , 1976	B M S
<i>Xyelatana</i> Benson, 1938	B M	<i>arvensis</i> Panzer, 1805	B M S
<i>curva</i> Benson, 1938	B M S	<i>erythrogaster</i> (Hartig, 1837)	B M S
<i>julii</i> (Brébison, 1818)	B M	<i>fallenii</i> (Dalman, 1823)	B M S
<i>longula</i> Dalman, 1819	B M	<i>hartigi</i> (Bremi, 1849)	B M
<i>obscura</i> (Strobl, 1895)	S	<i>lariciphila</i> (Wachtl, 1898)	B M
<i>julii v. tetrica</i> Gregor, 1940		<i>alpina</i> auct. nec Klug, 1808	
<b>Pleroneura</b> Konow, 1897		PAMPHILIINAE	
<i>coniferarum</i> (Hartig, 1837)	B M	<b>Pamphilius</b> Latreille, 1802	
<i>dahli</i> (Hartig, 1837)	M S	<i>albopictus</i> (Thomson, 1871)	B
PAMPHILIOIDEA		<i>alternans</i> (Costa, 1859)	B M
PAMPHILIIDAE		<i>aurantiacus</i> (Giraud, 1857)	B M
CEPHALCIINAE		<i>neglectus</i> Zaddach, 1865 nec auct.	
CAENOLYDINI		<i>balteatus</i> (Fallén, 1808)	B M
<b>Caenolyda</b> Konow, 1897		<i>betulae</i> (Linnaeus, 1758)	B M
<i>reticulata</i> (Linnaeus, 1758)	B M	<i>brevicornis</i> Hellén, 1948	M
CEPHALCIINI		<i>fumipennis</i> (Curtis, 1831)	M
<b>Acantholyda</b> Costa, 1894		<i>gyllenhali</i> (Dahlbom, 1835)	B
<i>erythrocephala</i> (Linnaeus, 1758)	B M S	<i>histrio</i> Latreille, 1811	B M
<i>flaviceps</i> (Retzius, 1783)	B S	<i>hortorum</i> (Klug, 1808)	B M S
<i>hieroglyphica</i> (Christ, 1791)	B M	<i>inanitus</i> (Villers, 1789)	B M S
<i>laricis</i> (Giraud, 1861)	B M	<i>latifrons</i> (Fallén, 1808)	B
<i>posticalis</i> (Matsumura, 1912)	B M S	<i>lethierryi</i> (Konow, 1887)	B

<i>marginatus</i> (Lepeletier, 1823)	B M	<i>fuscipennis</i> (Herrich- -Schaeffer 1833)	B M S
<i>nemorum</i> (Gmelin, 1788)	B M S	<i>gracilicornis</i> (Klug, 1812)	B M S
<i>pallipes</i> (Zetterstedt, 1838)	B M	<i>melanochroa</i> (Gmelin in Linnaeus, 1790)	B M S
<i>stramineipes</i> (Hartig, 1837)	M	<i>metallica</i> (Klug, 1834)	M S
<i>sylvarum</i> (Stephens, 1835)	B M	<i>nigripes nigripes</i> (Retzius, 1783)	B M S
<i>sylvaticus</i> (Linnaeus, 1758)	B M S	<i>enodis</i> auct. nec Linnaeus, 1758	
<i>thorwaldi</i> Kontuniemi, 1946	M	<i>ochropus</i> (Gmelin in Linnaeus, 1790)	B M S
<i>vafer</i> (Linnaeus, 1767)	B M S	<i>rosae</i> auct. nec Linnaeus, 1758	
<i>varius</i> (Lepeletier, 1823)	B M	<i>pagana</i> (Panzer, 1798)	B M S
NEUROTOMINI		<i>pleuritica</i> (Klug, 1834)	S
<b>Neurotoma</b> Konow, 1897		<i>pullata</i> (Zaddach, 1859)	B
<i>fausta</i> (Klug, 1808)	B S	<i>rustica</i> (Linnaeus, 1758)	B M S
<i>mandibularis</i> (Zaddach, 1865)	B M	<i>ustulata</i> (Linnaeus, 1758)	B M S
<i>nemoralis</i> (Linnaeus, 1758)	B M	STERICTIPHORINAE	
<i>saltuum</i> (Linnaeus, 1758)	B M	STERICTIPHORINI	
<i>flaviventris</i> Ratzeburg, 1783		<b>Sterictiphora</b> Billberg, 1820	
MEGALODONTIDAE		<i>furcata</i> (Villers, 1789)	M S
MEGALODONTINAE		<i>geminata</i> (Gmelin, in Linnaeus, 1790)	B M S
<b>Megalodontes</b> Latreille, 1802		<b>Aprosthemata</b> Konow, 1899	
<i>cephalotes</i> (Fabricius, 1781)	M S	<i>axillaris</i> (Zaddach, 1863)	B
<i>flavicornis</i> (Klug, 1824)	S	<i>brevicornis</i> (Fallén, 1808)	B
<i>klugi</i> (Leach, 1817)	M S	<i>maculata</i> (Jurine, 1807)	M S
<i>laticeps</i> Konow, 1897	S	<i>melanura melanura</i> (Klug, 1812)	B M
<i>plagiocephalus</i> (Fabricius, 1804)	B M S	<i>melanura friesei</i> (Konow, 1892)	B
<b>Melanopus</b> Konow, 1897		<i>melanura humerata</i> (Konow, 1892)	M
<i>fabricii</i> (Leach, 1817)	M S	<i>tarda</i> (Klug, 1812)	M
TENTHREDINOIDEA		BLASTICOTOMIDAE	
ARGIDAE		BLASTICOTOMINAE	
ARGINAE		<b>Blasticotoma</b> Klug, 1834	
<b>Arge</b> Schrank, 1802		<i>filiceti filicetti</i> Klug, 1834	B M
<i>berberidis</i> Schrank, 1802	B M S	CIMBICIDAE	
<i>ciliaris</i> (Linnaeus, 1767)	B M	ZARAEINAE	
<i>clavicornis</i> (Fabricius, 1793)	B M	<b>Zaraea</b> Leach, 1817	
<i>cyanocrocea</i> (Foerster, 1771)	B M S	<i>aenea</i> (Klug, 1839)	B
<i>dimidiata</i> (Fallén, 1808)	B M S		
<i>enodis</i> (Linnaeus, 1767)	B M S		
<i>coeruleipennis</i> (Retzius, 1783)			

<i>bigens</i> (Kangas, 1946)	B		<i>juniperi</i> (Linnaeus, 1758)	M S
<i>fasciata</i> (Linnaeus, 1758)	B M S		<i>obscuratus</i> (Hartig, 1837)	S
<i>lonicerae</i> (Linnaeus, 1758)	B M S		DIPRIONINAE	
<i>hugarica</i> (Mossáry, 1883)	M		<b>Microdiprion</b> Enslin, 1917	
<i>mutica</i> (Thomson, 1871)	B		<i>pallipes</i> (Fallén, 1808)	B M S
<b>Abia</b> Leach, 1817			<b>Neodiprion</b> Rohwer, 1918	
<i>candens</i> Konow, 1887	B S		<i>sertifer</i> (Geoffroy in Fourcroy, 1785)	B M S
<i>fulgens</i> Zaddach, 1863	B M S		<b>Diprion</b> Schrank, 1802	
<i>nitens</i> (Linnaeus, 1758)	B M		<i>pinii</i> (Linnaeus, 1758)	B M S
<i>sericea</i> (Linnaeus, 1767)	B M S		<i>similis</i> (Hartig, 1834)	B M
CIMBICINAE			<b>Gilpinia</b> Benson, 1939	
CIMBICINI			<i>abieticola</i> (Dalla-Torre, 1894)	B M
<b>Palaeocimbex</b> Semenov-Tianshanskij, 1935			<i>frutetorum</i> (Fabricius, 1793)	B M S
<i>quadrimaculata</i> (Müller, 1766)	M		<i>hercyniae</i> (Hartig, 1837)	B M
<b>Cimbex</b> Olivier, 1790			<i>laricis</i> (Jurine, 1807)	B M
<i>connata</i> (Schrank, 1776)	B S		<i>pallida</i> (Klug, 1812)	B M
<i>fagi</i> Zaddach, 1863	B		<i>polytoma</i> (Hartig, 1834)	B M S
<i>femorata</i> (Linnaeus, 1758)	B M S		<i>socia</i> (Klug, 1812)	B
<i>lutea</i> (Linnaeus, 1761)	B S		<i>variegata</i> (Hartig, 1834)	B M S
TRIGHIOSOMINI			<i>virens</i> (Klug, 1812)	B M
<b>Trichiosoma</b> Leach, 1817			<b>Macrodiprion</b> Enslin, 1917	
<i>latreillei</i> Leach, 1817	B S		<i>nemoralis</i> Enslin, 1917	M S
<i>lucorum</i> (Linnaeus, 1758)	B M S		TENTHREDINIDAE	
<i>sorbi</i> Hartig, 1840	B S		SELANDRIJINAE	
<i>sylvaticum</i> Leach, 1817	B		STRONGYLOGASTRINI	
<i>tibiale</i> Stephens, 1835	B M S		<b>Strongylogaster</b> Dahlbom, 1835	
<i>vitellinae</i> (Linnaeus, 1761)	B M S		<i>Thrinax</i> Konow, 1885	
<b>Pseudoclavellaria</b> Schulz, 1906			<i>contigua</i> Konow, 1885	B
<i>amerinae</i> (Linnaeus, 1758)	B M S		<i>lineata lineata</i> (Christ, 1791)	B M
CORYNINAE			<i>macula</i> (Klug, 1814)	M
<b>Corynis</b> Thunberg, 1789			<i>mixta</i> (Klug, 1814)	B M
<i>crassicornis</i> (Rossi, 1790)	B M S		<i>xanthocera</i> (Stephens, 1835)	B
<i>obscura</i> (Fabricius, 1775)	B M S		<b>Strombocerina</b> Malaise, 1942	
DIPRIONIDAE			<i>delicatula</i> (Fallén, 1808)	B M S
MONOCTENINAE			ANEUGMENINI	
<b>Monocetus</b> Dahlbom, 1835			<b>Aneugmenus</b> Hartig, 1837	
			<i>coronatus</i> (Klug, 1814)	B M
			<i>padi</i> (Linnaeus, 1761)	B M

<i>temporalis</i> (Thomson, 1871)	B M S	<i>gonager</i> (Fabricius, 1781) B M S
<b>Birka</b> Malaise, 1944		<i>haematodes</i> (Schrank, 1781) B M S
<i>Melisandra</i> Benson, 1939 p.p.		<i>harwoodi</i> Benson, 1947 B
<i>annulitarsis</i> (Thomson, 1870)	B M	<i>liogaster</i> Thomson, 1871 B M
<i>cinereipes</i> (Klug, 1814)	B M S	<i>rugosulus</i> Dalla-Torre, 1894
<b>Nesoselandria</b> Rohwer, 1910		<i>madiodus</i> (Klug., 1814) B M S
<i>Melisandra</i> Benson, 1939 p.p.		<i>megapterus</i> Cameron, 1881 B M S
<i>morio</i> (Fabricius, 1781)	B M S	<i>niger</i> (Linnaeus, 1767) B M S
SELANDRIINI		<i>nigratus</i> (Müller, 1776) B M S
<b>Selandria</b> Leach, 1817		<i>nitens</i> Zaddach, 1859 B M S
<i>excisa</i> (Konow, 1885)	B M	<i>picipes</i> (Klug, 1814) B M S
<i>sixii</i> Vollenhoven, 1858)	B M	<i>pratensis</i> (Linnaeus, 1758) B M S
<i>serva</i> (Fabricius, 1793)	B M S	<i>puncticollis</i> Thomson, 1871 B M S
<b>Brachythops</b> Haliday, 1839		<i>quadrinotatus</i> Biro, 1884 S
<i>flavens</i> (Klug, 1814)	B M	<i>sanguinicollis</i> (Klug, 1814) B M S
HEPTAMELINI		<i>taeniatus</i> Zaddach, 1859 B M
<b>Heptamelus</b> Haliday, 1855		<i>thoracicus</i> Fallén, 1808 B M S
<i>ochroleucus</i> (Stephens, 1835)	M	<i>triplicatus</i> (Klug, 1814) B M S
DOLERINAE		<i>uliginosus</i> (Klug, 1814) B M
<b>Dolerus</b> Panzer, 1801		<i>yukonensis</i> <i>yukonensis</i> (Norton, 1872) B M
<i>aeneus</i> (Hartig, 1837)	B M S	<i>saxatilis</i> Hartig, 1837
<i>aericeps</i> Thomson, 1871	B M	<b>Loderus</b> Konow, 1890
<i>anticus</i> (Klug, 1814)	B M S	<i>eversmanni</i> (Kirby, 1882) B M S
<i>anthracinus</i> (Klug, 1814)	B M S	<i>genucinctus</i> Zaddach, 1859 M
<i>asper</i> Zaddach, 1859	B M	<i>pratorum</i> (Fallén, 1808) B M S
<i>bimaculatus</i> (Geoffroy in Fourcroy, 1785)	B M S	<i>vestigialis</i> (Klug, 1814) B M S
<i>brevitarsis</i> Hartig, 1837	B M S	NEMATINAE
<i>coracinus</i> (Klug, 1814)	B M	CLADIINI
<i>cothurnatus</i> Lepeletier, 1823	B M S	<b>Cladius</b> Illiger, 1807
<i>palustris</i> Klug, 1814		<i>diformis</i> (Panzer, 1799) B
<i>ferrugatus</i> Lepeletier, 1823	B M	<i>pectinicornis</i> (Geoffroy in Fourcroy, 1785) B M S
<i>germanicus germanicus</i> (Fabricius, 1775)	B M S	<b>Priophorus</b> Dahlbom, 1835
<i>germanicus etruscus</i> (Klug, 1814)	M S	<i>morio</i> (Lepeletier, 1823) B M S
<i>gessneri gessneri</i> André, 1880	B M	<i>pallipes</i> (Lepeletier, 1823) B M S
<i>gibbosus</i> Hartig, 1837	B M	<b>Trichiocampus</b> Hartig, 1837
		<i>pilicornis</i> (Curtis, 1833) B M
		<i>rufipes</i> (Lepeletier, 1823) B M
		<i>ulmi</i> auct. nec Linnaeus, 1758
		<i>ulmi</i> (Linnaeus, 1758) M
		<i>laevifrons</i> Benson, 1936
		<i>viminalis</i> (Fallén, 1808) B M S

<b>NEMATINI</b>			
<b>Hoplocampus</b> Hartig, 1837			
<i>alpina</i> (Zetterstedt, 1838)	B	S	
<i>ariae</i> Benson, 1933	B		
<i>brevis</i> (Klug, 1814)	M		
<i>chrysorrhoea</i> (Klug, 1814)	B		
<i>crataegi</i> (Klug, 1814)	B	M	S
<i>flava</i> (Linnaeus, 1761)	B	M	
<i>minuta</i> (Christ, 1791)	B	M	S
<i>pectoralis</i> Thomson, 1871	B		
<i>plagiata</i> (Klug, 1814)	B		
<i>rutilicornis</i> (Klug, 1814)	B	M	
<i>testudinea</i> (Klug, 1814)	B		
<b>Hemicreba</b> Stephens, 1835			
<i>australis</i> (Lepeletier, 1823)	B	M	S
<i>alni</i> Linnaeus, 1758			
<i>crocea</i> (Geoffroy in Fourcroy, 1785)	B	M	
<b>Platycampus</b> Schiödte, 1839			
<i>luridiventris</i> (Fallén, 1808)	B		
<b>Anoplonyx</b> Marlatt, 1896			
<i>duplex</i> (Lepeletier, 1823)	B	M	
<i>ovatus</i> (Zaddach, 1883)	M	S	
<i>pectoralis</i> (Lepeletier, 1823)	M		
<b>Dineura</b> Dahlbom, 1835			
<i>stilata</i> (Klug, 1814)	B		
<i>testaceipes</i> (Klug, 1814)	B	M	
<i>virididorsata</i> (Retzius, 1783)	B		
<b>Mesoneura</b> Hartig, 1837			
<i>opaca</i> (Fabricius, 1775)	B	M	
<b>Nematinus</b> Rohwer, 1911			
<i>abdominalis</i> (Panzer, 1799)	B	M	S
<i>acuminatus</i> (Thomson, 1871)	B	M	
<i>luteus</i> (Panzer, 1805)	B	M	S
<i>willigkiae</i> (Stein, 1926)	S		
<b>Pachynematus</b> Konow, 1890			
<i>albibennnis</i> (Hartig, 1837)	B	M	
<i>apicalis</i> (Hartig, 1837)	M	S	
<i>calcicola</i> Benson, 1948	B		
<i>clitellatus</i> (Lepeletier, 1823)	B	M	S
<i>declinatus</i> (Foerster, 1854)	M		
<i>kirbyi</i> (Dahlbom, 1835)	B	M	S
<i>kubesi</i> Konow, 1905	B	M	
<i>laevigatus</i> (Zaddach, 1875)	M		
<i>lichtwardti</i> Konow, 1903	B	M	
<i>moerens</i> (Foerster, 1854)	M		
<i>nigerrimus</i> Konow, 1903	M		
<i>obductus</i> (Hartig, 1837)	B	M	
<i>parvilabris</i> (Thomson, 1862)	B		
<i>pullus</i> Konow, 1890	B	S	
<i>rumicis</i> (Linnaeus, 1758)	B	M	S
<i>subaequalis</i> (Foerster, 1854)	B	M	
<i>vagus</i> (Fabricius, 1781)	B	M	S
<i>xanthocarpus</i> (Hartig, 1837)	B	M	
<b>Pikonema</b> Ross, 1937			
<i>imperfecta</i> (Zaddach, 1875)	B	M	
<i>insignis</i> (Hartig, 1840)	B	M	
<i>montana</i> (Zaddach, 1882)	B	M	S
<i>pallescents</i> (Hartig, 1837)	B	M	S
<i>scutellata</i> (Hartig, 1837)	B	M	S
<i>styx</i> (Benson, 1954)	M		
<b>Stauronematus</b> Benson, 1948			
<i>compresicornis</i> (Fabricius, 1804),	B	M	
<b>Pristiphora</b> Latreille, 1810			
<i>abbreviata</i> (Hartig, 1837)	B		
<i>abietina</i> (Christ, 1791)	B	M	S
<i>albilabris</i> (Thomson, 1862)	B		
<i>anderschi</i> (Zaddach, 1875)	B	M	
<i>aquilegiae</i> (Vollenhoven, 1866)	B	M	
<i>bifida</i> (Hellén, 1948)	B		
<i>biscalis</i> (Foerster, 1854)	B	M	
<i>borea</i> (Konow, 1904)	B	S	
<i>carinata</i> (Hartig, 1837)	B	M	
<i>cincta</i> Newman, 1837	B	M	S
<i>quercus</i> Hartig, 1837			
<i>compressa</i> (Hartig, 1837)	B	M	

<i>conjugata</i> (Dahlbom, 1835)	B M S	<i>septentrionalis</i> (Linnaeus, 1758)	B M S
<i>crassicornis</i> (Hartig, 1837)	B M	<i>varus</i> (Villaret, 1832)	B M
<i>decipiens</i> (Enslin, 1916)	B M	<b>Nematus</b> Panzer, 1801	
<i>denudata</i> (Konow, 1904)	M	<i>bergmanni</i> Dahlbom, 1835	B M S
<i>erichsonii</i> (Hartig, 1837)	B	<i>bipartitus</i> (Lepeletier, 1823)	B M
<i>friesei</i> (Konow, 1904)	M	<i>capito</i> (Konow, 1903)	B M
<i>fulvipes</i> (Fallén, 1808)	B M S	<i>coerulocarpus</i> Hartig, 1837	B M S
<i>geniculata</i> (Hartig, 1840)	B M S	<i>crassus</i> (Fallén, 1808)	B M
<i>gerula</i> (Konow, 1904)	B M	<i>dispar</i> Brischke, 1875	B M
<i>laricis</i> (Hartig, 1837)	B M S	<i>ferrugineus</i> Foerster, 1854	B M S
<i>leucopodia</i> (Hartig, 1837)	B M	<i>fahraei</i> Thomson, 1862	B M
<i>melanocarpa</i> (Hartig, 1840)	B M	<i>flavescens</i> Stephens, 1835	B M S
<i>moesta</i> (Zaddach, 1883)	M	<i>forsiusi</i> (Enslin, 1915)	B
<i>mollis</i> (Hartig, 1837)	B M S	<i>frenalis</i> Thomson, 1871	S
<i>monogyniae</i> (Hartig, 1840)	B M	<i>fuscomaculatus</i> Foerster, 1854	M
<i>moravica</i> Gregor, 1940	M S	<i>hypoxanthus</i> Foerster, 1854	B M
<i>nigripes</i> (Hartig, 1837)	B M	<i>incompletus</i> Foerster, 1854	B M
<i>pallida</i> (Konow, 1904)	M	<i>leucotrochus</i> Hartig, 1837	B M
<i>pallidiventris</i> (Fallén, 1808)	B M S	<i>lucidus</i> (Panzer, 1797)	B M
<i>pallidula</i> Konow, 1902	M	<i>melanaspis</i> Hartig, 1840	B M
<i>pseudodecipiens</i> Beneš et Kříštek, 1976	M	<i>melanocephalus</i> Hartig, 1837	M
<i>punctifrons</i> (Thomson, 1871)	B M S	<i>miliaris</i> (Panzer, 1797)	M S
<i>robusta</i> (Konow, 1895)	B	<i>mimus</i> (Konow, 1903)	B M
<i>ruficornis</i> (Olivier, 1811)	B M S	<i>myosotidis</i> (Fabricius, 1809)	B M S
<i>rufipes</i> Lepeletier, 1823	B M S	<i>nigricornis</i> (Lepeletier, 1823)	B M S
<i>saxesenii</i> (Hartig, 1837)	B M S	<i>notabilis</i> (Konow, 1903)	M
<i>subarctica</i> (Forsslund, 1936)	B M	<i>oligospilus</i> Foerster, 1854	B M S
<i>subbifida</i> (Thomson, 1871)	M S	<i>pavidus</i> (Lepeletier, 1823)	B M
<i>testacea</i> (Jurine, 1807)	B M	<i>polyspilus</i> Foerster, 1854	B M S
<i>tetrica</i> (Zaddach, 1863)	M	<i>prasinus</i> Hartig, 1837	B M
<i>thalictri</i> Kriechbaumer, 1884	B	<i>ribesii</i> (Scopoli, 1763)	B M S
<i>wesmaeli</i> (Tischbein, 1853)	B M	<i>salicis</i> (Linnaeus, 1758)	B M S
<b>Sharliphora</b> Wong, 1969		<i>similator</i> Foerster, 1854	M
<i>ambigua</i> (Fallén, 1808)	B M S	<i>spiraeeae</i> Zaddach, 1883	B
<i>amphibola</i> (Foerster, 1854)	M S	<i>tibialis</i> Newman, 1837	B M S
<i>parva</i> (Hartig, 1837)	M	<i>viridescens</i> Cameron, 1885	B M
<b>Croesus</b> Leach, 1817		<i>wahlbergi</i> Thomson, 1871	M S
<i>brischkei</i> (Zaddach, 1875)	B M	<b>Phyllocolpa</b> , Benson, 1960	
<i>latipes</i> (Villaret, 1832)	B M	<i>anglica</i> (Cameron, 1883)	M
		<i>coriacea</i> (Benson, 1953)	S
		<i>excavata</i> (Marlatt, 1896)	M
		<i>leucapsis</i> (Tischbein, 1846)	B M S

<i>leucosticta</i> (Hartig, 1837)	B	M	S
<i>piliserra</i> (Thomson, 1862)	B	M	
<i>puella</i> (Thomson, 1871)	B	M	
<i>purpureae</i> (Cameron, 1884)	B	M	S
<i>scotaspis</i> (Foerster, 1854)		M	
<b>Pontania</b> Costa, 1859			
<i>bridgmani</i> (Cameron, 1883)	B	M	S
<i>collectanea</i> (Foerster, 1854)	M	S	
<i>crassipes</i> (Thomson, 1871)	S		
<i>dolichura</i> (Thomson, 1871)	B	M	S
<i>joergenseni</i> Enslin, 1916	M		
<i>kriechbaumeri</i> Konow, 1901	M	S	
<i>pedunculi</i> (Hartig, 1837)	B	M	S
<i>proxima proxima</i> (Lepeletier, 1823)	B	M	S
<i>proxima triandrae</i> (Benson, 1940)	B	M	S
<i>reticulatae</i> Malaise, 1920	S		
<i>arctica</i> auct. nec Mac Gillivray, 1919			
<i>retusae</i> Benson, 1960	S		
<i>vesicator</i> (Bremi, 1849)	B	M	S
<i>viminalis</i> (Linnaeus, 1758)	B	M	S
<b>Euura</b> Newman, 1837			
<i>amerinae</i> (Linnaeus, 1758)	B	M	S
<i>atra</i> (Jurine, 1807)	B	M	S
<i>laeta</i> (Zaddach, 1883)	B	M	
<i>lappo</i> Malaise, 1920	B		
<i>mucronata</i> (Hartig, 1837)	B	M	S
<i>saliceti</i> auct. nec Fallén, 1808			
<i>testaceipes</i> (Zaddach, 1883)	B	M	S
<i>venusta</i> (Zaddach, 1883)	B	M	S
<b>Amauronematus</b> Konow, 1890			
<i>alpicola</i> Konow, 1890	B	M	
<i>amplus</i> Konow, 1895	B	S	
<i>distinguendas</i> Enslin, 1915	M		
<i>fallax</i> (Lepeletier, 1823)	B	M	
<i>fasciatus</i> Konow, 1907	B		
<i>histrio</i> (Lepeletier, 1823)	B	M	S
<i>humeralis</i> (Lepeletier, 1823)	B	M	
<i>leucolaenus</i> (Zaddach, 1882)	M	S	
<i>miltonotus</i> (Zaddach, 1882)		B	
<i>polaris</i> (Holmgren, 1883)		S	
<i>leucopyga</i> Lindqvist, 1948			
<i>punicetus</i> (Christ, 1791)	B	M	
<i>taeniatus</i> (Lepeletier, 1823)		M	
<i>tunicatus</i> (Zaddach, 1882)	B	M	
<i>viduatus</i> (Zetterstedt, 1838)	B	M	S
<i>vittatus</i> (Lepeletier, 1823)	B	M	
<b>Pontopristia</b> Malaise, 1921			
<i>amentorum</i> (Foerster, 1854)		M	
<i>boreoalpina</i> Lindqvist, 1961		S	
<b>Decanematus</b> Malaise, 1931			
<i>logiserra</i> (Thomson, 1871)	B		
<b>PSEUDODINEURINI</b>			
<b>Pseudodineura</b> Konow, 1885			
<i>enslini</i> (Hering, 1923)	B		
<i>fuscula</i> (Klug, 1814)	B	M	
<i>heringi</i> (Enslin, 1921)	M		
<i>mentiens</i> (Thomson, 1871)	M		
<i>parvula</i> (Klug, 1814)	M		
<b>HETERARTHINAE</b>			
<b>CALIROINI</b>			
<b>Endelomyia</b> Ashmead, 1898			
<i>aethiops</i> (Fabricius, 1781)	B	M	S
<b>Caliroa</b> Costa, 1859			
<i>annulipes</i> (Klug, 1814)	B	M	S
<i>cerasi</i> (Linnaeus, 1758)	B	M	S
<i>cinxia</i> (Klug, 1814)	B	M	S
<i>varipes</i> (Klug, 1814)	B	M	S
<b>HETERARTHINI</b>			
<b>Heterarthrus</b> Stephens, 1835			
<i>fumipennis</i> (Cameron, 1888)	B	M	
<i>leucomelus</i> (Klug, 1814)	M		
<i>microcephalus</i> (Klug, 1814)	B	M	
<i>nemoratus</i> (Fallén, 1808)	B	M	
<i>ochropodus</i> (Klug, 1814)	B	M	

<i>vagans</i> (Fallén, 1808)	B M S	<i>scutellariae</i> Cameron, 1880	
FENUSINI			M
<b>Metallus</b> Forbes, 1885			
<i>albipes</i> (Cameron, 1875)	M		
<i>gei</i> (Brischke, 1883)	B M		
<i>pumilus</i> (Klug, 1814)	B M		
<b>Messa</b> Leach, 1817			
<i>Fenusella</i> Enslin, 1914			
<i>glaukopis</i> (Konow, 1907)	M		
<i>hortulana</i> (Klug, 1814)	M		
<i>nana</i> (Klug, 1814)	B		
<i>wuestneii</i> (Konow, 1894)	B M		
<b>Profenusia</b> Mac Gillivray, 1914			
<i>Gunnea</i> Malaise, 1964			
<i>pygmaea</i> (Klug, 1814)	B M S		
<i>thomsoni</i> (Konow, 1886)	B		
<b>Scolioneura</b> Konow, 1890			
<i>betuleti</i> (Klug, 1814)	B M		
<b>Fenusia</b> Leach, 1817			
<i>dohrnii</i> (Tischbein, 1846)	B M		
<i>pusilla</i> (Lepeletier, 1823)	B M		
<i>ulmi</i> Sundevall, 1844	M		
<b>Fenella</b> Westwood, 1840			
<i>minuta</i> Thomson, 1871	B M		
<i>nigrita</i> Westwood, 1840	B M		
<b>Parna</b> Benson, 1936			
<i>tenella</i> (Klug, 1814)	B M		
<b>Hinatara</b> Benson, 1936			
<i>excisa</i> (Konow, 1885)	M		
<i>recta</i> (Thomson, 1871)	M		
BLENNOCAMPINAE			
ATHALIINI			
<b>Athalia</b> Leach, 1817			
<i>bicolor</i> Lepeletier, 1823	B M S		
<i>circularis</i> (Klug, 1813)	B M S		
<i>cordata</i> Lepeletier, 1823	B M S		
<i>glabricollis</i> Thomson, 1870	B M S		
<i>liberta</i> (Klug, 1813)	B M S		
<i>lugens lugens</i> (Klug, 1813)	B M S		
<i>rosae</i> (Linnaeus, 1758)	B M S		
<i>rufoscutellata</i> Mocsáry, 1879	B M		
LYCAOTINI			
<b>Hoplocampoides</b> Enslin, 1914			
<i>xylostei</i> (Graud, 1863)	B M S		
TOMOSTETHIINI			
<b>Tomostethus</b> Konow, 1886			
<i>nigritus</i> (Fabricius, 1804)	B M		
PHYMATOCERINI			
<b>Phyamatocera</b> Dahlbom, 1835			
<i>aterrima</i> (Klug, 1814)	B M S		
Paracharactus Mac Gillivray, 1908			
<i>gracilicornis</i> (Zaddach, 1859)	B M		
<i>longicornis</i> (Hartig, 1837)	B M		
Rhadinoceraea Konow, 1886			
<i>brensoni</i> Beneš, 1961	B S		
<i>micans</i> (Klug, 1814)	B M		
<i>nodicornis</i> Konow, 1906	B		
<i>reitteri</i> Konow, 1890	M S		
<b>Monophadnus</b> Hartig, 1837			
<i>monticola</i> Hartig, 1837	B M S		
<i>pallescens</i> (Gmelin in Linnaeus, 1790)	B M S		
<i>semicinctus</i> Hartig, 1837	B		
<i>spinolai</i> (Klug, 1814)	M S		
<b>Stethomostus</b> Benson, 1952			
<i>fuliginosus</i> (Schrank, 1781)	B M S		
<i>funereus</i> (Klug, 1814)	B M		
<b>Eutomostethus</b> Enslin, 1914			
<i>ephippium</i> (Panzer, 1798)	B M S		
<i>gagathinus</i> (Klug, 1814)	B M S		
<i>luteiventris</i> (Klug, 1814)	B M S		
<i>punctatus</i> (Konow, 1887)	M		
BLENNOCAMPINI			
<b>Blennocampa</b> Hartig, 1837			
<i>pusilla</i> (Klug, 1814)	B M S		
<b>Ardis</b> Konow, 1886			

<i>brunniventris</i> (Hartig, 1837)	B M		<i>immersa</i> (Klug, 1814)	M S
<i>sulcata</i> (Cameron, 1882)	M		<i>klugi</i> (Stephens, 1835)	B M S
<b>Monardis</b> Enslin, 1914			<i>liturata</i> (Gmelin in Linnaeus, 1790)	B M S
<i>plana</i> (Klug, 1814)	B M S		<i>longicornis</i> (Thomson, 1871)	B M
<b>Periclista</b> Konow, 1886			<i>parvula</i> (Konow, 1884)	B M
<i>albida</i> (Klug, 1814)	B M		<i>pumila</i> (Konow, 1896)	B M S
<i>lineolata</i> (Klug, 1814)	B		<i>testaceipes</i> (Konow, 1896)	M
<b>Monophadnoides</b> Ashmead, 1898			<i>tridens</i> (Konow, 1896)	B M
<i>Claremontia</i> Rohwer, 1909			<b>Monosoma</b> Mac Gillivray, 1908	
<i>alternipes</i> (Klug, 1814)	B M S		<i>pulverata</i> (Retzius, 1783)	B M S
<i>confusa</i> (Konow, 1886)	B M S		<b>Monostegia</b> Costa, 1859	
<i>geniculata</i> (Hartig, 1837)	B M		<i>abdominalis</i> (Fabricius, 1798)	B M S
<i>puncticeps</i> (Konow, 1886)	B M		<b>Ametastegia</b> Costa, 1882	
<i>ruficruris</i> (Brullé, 1832)	B M		<i>Protomphytus</i> Rohwer, 1909	
<i>tenuicornis</i> (Klug, 1814)	B M		<i>albipes</i> (Thomson, 1871)	B
<i>waldheimii</i> (Gimmerthal, 1847)	B M		<i>carpini</i> (Hartig, 1837)	B M S
<b>Pareophora</b> Konow, 1886			<i>equiseti</i> (Fallén, 1808)	B M S
<i>pruni</i> (Linnaeus, 1758)	B M		<i>glabrata</i> (Fallén, 1808)	B M S
<b>Cladardis</b> Benson, 1952			<i>pallipes</i> (Spinola, 1808)	B M S
<i>elongatula</i> (Klug, 1814)	B M		<i>perla</i> (Klug, 1814)	B M
WALDHEIMIINI			<i>stictica</i> (Klug, 1814)	B M S
<b>Halidamia</b> Benson, 1952			<i>tener</i> (Fallén, 1808)	B M S
<i>affinis</i> (Fallén, 1807)	B M		<b>Eopsis</b> Benson, 1959	
ALLANTINAE			<i>beaumonti</i> Benson, 1959	B S
ERIOCAMPINI			ALLANTINI	
<b>Eriocampa</b> Hartig, 1837			<b>Allantus</b> Panzer, 1801	
<i>ovata</i> (Linnaeus, 1761)	B M S		<i>Emphytus</i> Klug, 1813	
<i>umbratica</i> (Klug, 1814)	B M S		<i>balteatus</i> (Klug, 1814)	B M S
<b>Siobia</b> Cameron, 1877			<i>basalis</i> (Klug, 1814)	B M S
<i>sturmi</i> (Klug, 1814)	B M		<i>calceatus</i> (Klug, 1814)	B M S
EMPRIINI			<i>cinctus</i> (Linnaeus, 1758)	B M S
<b>Harpiphorus</b> Hartig, 1837			<i>cingillum</i> (Klug, 1814)	B M S
<i>lepidus</i> (Klug, 1814)	B M		<i>cingulatus</i> (Scopoli, 1763)	B M S
<b>Empria</b> Lepeletier, 1823			<i>coxalis</i> (Klug, 1814)	B
<i>alboscutellata</i> Konow, 1894	M		<i>melanarius</i> (Klug, 1814)	B M S
<i>alector</i> Benson, 1938	B S		<i>rufocinctus</i> (Retzius, 1783)	B M S
<i>baltica</i> Conde, 1937	B S		<i>togatus</i> Panzer, 1801	B M
<i>candidata</i> (Fallén, 1807)	B M S		<i>truncatus</i> (Klug, 1814)	B M
<i>excisa</i> (Thomson, 1871)	B M		<i>viennensis</i> (Schrank, 1781)	B M
<i>hungarica</i> (Konow, 1895)	B M		TAXONUS Hartig, 1837	

<i>agrorum</i> (Fallén, 1808)	B M S	<i>costalis</i> (Fabricius, 1775)	B M S
<b>Apethymus</b> Benson, 1939		<b>Elinora</b> Benson, 1946	
<i>abdominalis</i> (Lepeletier, 1823)	B	<i>flaveola</i> (Gmelin in Linnaeus, 1790)	B M
<i>braccatus</i> (Gmelin in Linnaeus, 1790)	B M	<i>sabariensis</i> (Mocsáry, 1880)	M S
<i>serotinus</i> (Müller, 1776)	B M		
TENTHREDINIINAE		TENTHREDININI	
PERINEURINI			
<b>Perineura</b> Hartig, 1837		<b>Rhogogaster</b> Konow, 1884	
<i>rubi</i> (Panzer, 1805)	B M	<i>chlorosoma</i> (Benson, 1943)	B
TENTHREDOPSINI		<i>dryas</i> (Benson, 1943)	B
<b>Aglaostigma</b> Kirby, 1882		<i>genistae</i> Benson, 1949	B
<i>aucupariae</i> (Klug, 1814)	B M S	<i>picta</i> (Klug, 1814)	B M S
<i>discolor</i> (Klug, 1814)	B	<i>punctulata</i> (Klug, 1814)	B M S
<i>fulvipes</i> (Scopoli, 1763)	B M S	<i>viridis</i> (Linnaeus, 1758)	B M S
<i>lichtwardti</i> (Konow, 1890)	M	<b>Tenthredo</b> Linnaeus, 1758	
<i>nebulosa</i> (André, 1881)	M	<i>albicornis</i> Fabricius, 1781	B M S
<i>pinguis</i> (Klug, 1814)	S	<i>amoena</i> Gravenhorst, 1807	B M
<b>Tenthredopsis</b> Costa, 1859		<i>arcuata</i> <i>arcuata</i>	
<i>annuligera</i> (Tischbein, 1846)	M S	Foerster, 1771	B M S
<i>arrogans</i> Konow, 1890	B M	<i>arcuata korabica</i> (Czíki, 1923)	S
<i>austriaca</i> Konow, 1890	B M	<i>atra</i> Linnaeus, 1758	B M S
<i>carbonaria</i> (Linnaeus, 1767)	B M S	<i>balteata</i> Klug, 1814	B M S
<i>coqueberti</i> (Klug, 1814)	B M S	<i>bifasciata rossii</i> (Panzer, 1805)	S
<i>excisa</i> Thomson, 1870	B M S	<i>bipunctula</i> Klug, 1814	B M
<i>floricola</i> Costa, 1859	B	<i>campestris</i> Linnaeus, 1758	B M S
<i>friesei</i> Konow, 1887	B M S	<i>colon</i> Klug, 1814	B M S
<i>hungarica</i> (Klug, 1814)	M	<i>costata</i> Klug, 1814	S
<i>inornata</i> Cameron, 1881	B M	<i>cunyi</i> Konow, 1886	B M S
<i>lactiflua</i> (Klug, 1814)	B M	<i>distinguenda</i> Stein, 1885	B S
<i>laticeps</i> Konow, 1884	M	<i>excelens</i> (Konow, 1886)	S
<i>nassata</i> (Linnaeus, 1767)	B M S	<i>fagi</i> Panzer, 1798	B M S
<i>parvula</i> Konow, 1890	B M	<i>ferruginea</i> Schrank, 1776	B M S
<i>scutellaris</i> (Fabricius, 1798)	B M S	<i>frivaldszkyi</i> Mocsáry, 1880	S
<i>sordida</i> (Klug, 1814)	B M S	<i>ignobilis</i> Klug, 1814	M
<i>stigma</i> (Fabricius, 1798)	B M S	<i>limbata</i> Klug, 1814	B M S
<i>tarsata</i> (Fabricius, 1804)	B M S	<i>livida</i> Linnaeus, 1758	B M S
<i>tesselata</i> (Klug, 1814)	B M	<i>maculata maculata</i>	
SCIAPTERYGINI		Geoffroy, 1762	B M S
<b>Sciapteryx</b> Stephens, 1835		<i>maculata semseyi</i>	M S
<i>consobrina</i> (Klug, 1814)	B M S	<i>mandibularis</i> Fabricius, 1804	B M S
		<i>marginella</i> Fabricius, 1793	B M S

<i>mesomelas</i>	Linnaeus, 1758	B M S	<i>postica</i> (Brullé, 1832)	S
<i>mioceras</i>	Enslin, 1912	B M S	<i>punctumalbum</i>	
<i>moniliata</i>	Klug, 1814	B M S	(Linnaeus, 1767)	B M S
<i>neobesa</i>	Zombori, 1980	S	<i>ribis</i> (Schrank, 1781)	B M S
<i>nitidior</i> {Konow, 1888}		B M S	<i>rufipes</i> (Linnaeus, 1758)	B M S
<i>aegra</i>	Enslin, 1912		<i>sanguinolenta</i> (Gmelin in Linnaeus, 1790)	B M S
<i>notha notha</i>	Klug, 1814	B M S	<i>tenella</i> Mocsáry, 1881	S
<i>perkinsi</i>	Morice, 1919		<i>teutona</i> (Panzer, 1799)	B M S
<i>obsoleta</i>	Klug, 1814	B M S	<b>Pachyprotasis</b> Hartig, 1837	
<i>olivacea</i>	Klug, 1814	B M S	<i>antennata</i> (Klug, 1814)	B M S
<i>omissa</i>	Foerster, 1844	B M S	<i>nigronotata</i>	
<i>procera</i>	Klug, 1814	B M S	(Kriechbaumer, 1874)	B M
<i>pseudorossii</i>	Taeger, 1984	B M S	<i>rapae</i> (Linnaeus, 1767)	B M S
<i>rossii</i> auct. non Panzer, 1805			<i>simulans</i> (Klug, 1814)	B M
<i>rubricoxis</i> {Enslin, 1912}	B M		<i>variegata</i> (Fallén, 1808)	B M S
<i>schaefferi</i>	Klug, 1814	B M S		
<i>scrophulariae</i>	Linnaeus, 1758	B M S	<b>SIRICOIDEA</b>	
<i>solitaria</i>	Scopoli, 1763	M S	<b>SIRICIDAE</b>	
<i>sulphuripes</i>			<b>SIRICINAE</b>	
(Kriechbaumer, 1869)	B M		<b>Sirex</b> Linnaeus, 1761	
<i>temula</i>	Scopoli, 1763	B M S	<i>juvencus juvencus</i>	
<i>trabeata</i>	Klug, 1814	B M S	(Linnaeus, 1758)	B M S
<i>velox</i>	Fabricius, 1798	B M S	<i>noctilio</i> Fabricius, 1793	B M S
<i>vespa</i>	Retzius, 1783	B M S	<b>Urocerus</b> Geoffroy, 1762	
<i>vespiformis</i>	Schrank, 1781	B M S	<i>augur augur</i> (Klug, 1803)	B
<i>zona</i>	Klug, 1814	B M S	<i>gigas gigas</i> (Linnaeus, 1758)	B M S
<i>zonula</i>	Klug, 1814	B M S	<i>tardigradus</i> (Cederhjelm, 1798)	B M S
<b>Cuneala</b> Zirngiebl, 1956			<b>Xeris</b> Costa, 1894	
<i>dahli</i> {Klug, 1814}	S		<i>spectrum spectrum</i>	
<i>koehleri</i> {Klug, 1814}	B M S		(Linnaeus, 1758)	B M S
<b>MACROPHYINI</b>			<b>TREMECINAE</b>	
<b>Macrophya</b> Dahlbom, 1835			<b>Tremex</b> Jurine, 1807	
<i>albicincta</i> {Schrank, 1776}	B M S		<i>fuscicornis</i> (Fabricius, 1787)	B M
<i>albimacula</i>	Mocsáry, 1881	S	<i>magus</i> (Fabricius, 1787)	M S
<i>albipuncta</i> {Fallén, 1808}	B M		<b>XIPHYDRIIDAE</b>	
<i>annulata</i> {Geoffroy in Fourcroy, 1785}	B M S		<b>XIPHYDRIINAE</b>	
<i>blanda</i> {Fabricius, 1775}	B M S		<b>Xiphydria</b> Latreille, 1802	
<i>carinthiaca</i> {Klug, 1814}	B M		<i>camelus</i> (Linnaeus, 1758)	B M S
<i>crassula</i> {Klug, 1814}	B M S		<i>longicollis</i> (Geoffroy in Fourcroy, 1785)	S
<i>diversipes</i> {Schrank, 1782}	B M S			
<i>duodecimpunctata</i>				
(Linnaeus, 1758)	B M S			
<i>erythrocnema</i>	Costa, 1859	M S		
<i>militaris</i> {Klug, 1814}	B M S			
<i>montana</i> {Scopoli, 1763}	B M S			
<i>pallidilabris</i> Costa, 1899	B M			

<i>prolongata</i> (Geoffroy in Fourcroy, 1785)	B M S	CEPHINI
ORUSSOIDEA		<b>Cephus</b> Latreille, 1802
ORUSSIDAE		<i>Pseudocephus</i> Dovnar-Zapolskij, 1931
ORUSSINAE		<i>brachycercus</i> Thomson, 1871 B M S
<b>Orussus</b> Latreille, 1796		<i>culturatus</i> Eversmann, 1847 B M S
<i>abietinus</i> (Scopoli, 1763)	B M	<i>fumipennis</i> Eversmann,, 1847 B M
CEPHOIDEA		<i>nigrinus</i> Thomson, 1871 B M S
CEPHIDAE		<i>pulcher</i> Tischbein, 1852 B
CEPHINAE		<i>pygmaeus</i> (Linnaeus, 1767) B M S
HARTIGIINI		<b>Calameuta</b> Konow, 1896
<b>Hartigia</b> Schioedte, 1838		<i>filiformis</i> (Eversmann, 1847) B M
<i>linearis</i> (Schrink, 1781)	B S	<i>haemorrhoidalis</i> (Fabricius, 1781) M S
<i>nigra</i> (Harris, 1776)	B M S	<i>pallipes</i> (Klug, 1803) B M S
<b>Janus</b> Stephens, 1835		<b>Trachelus</b> Jurine, 1807
<i>compressus</i> (Fabricius, 1793)	B M S	<i>tabidus</i> (Fabricius, 1775) S
<i>femoratus</i> (Curtis, 1830)	B M S	<i>troglodyta</i> (Fabricius, 1787) B M S
<i>luteipes</i> (Lepeletier, 1823)	B M	



APOCRITA  
PARASITICA  
ICHNEUMONOIDEA

BRACONIDAE

MIROSLAV ČAPEK, JOZEF LUKĀŠ

The Braconidae represent the second most numerous family of parasitic Hymenoptera. Several of its subfamilies, tribes or genera lack modern revisions and their species are not easily determinable. That is why the Braconid material from the collections of Czechoslovakian museums was not studied in detail and this check-list was made up on the basis of published data (first of all by Shenefelt, 1969—1978, and Shenefelt & Marsh, 1976) and original knowledge of the authors. Hence it follows that probably still one third or fourth part of species awaits the discovery of their occurrence in Czechoslovakia and a number of these will be described as new for science.

The development of Czechoslovakian braconidology began as late as the nineteen thirties when several outstanding Czech amateur hymenopterologists focussed their attention on parasitic Hymenoptera. They collected very intensively and namely in the warmer steppe localities of southern Moravia and Slovakia (Gregor sen., Hoffer, and Šnoflák), of which the last named specialized in Braconidae. His collection of Braconidae, deposited in the Moravian Museum in Brno, is among the largest in Czechoslovakia. Kolubajiv should be mentioned from among applied entomologists, as he extended the knowledge of Braconidae parasitizing forest insect pests. The family was also studied by Starý for several years and finally, at the present time, by the authors of this check-list.

The braconid fauna of southern Moravia and southern Slovakia as well as of broadleaved and coniferous forests in the whole of Czechoslovakia are rather well-known, while the fauna of mountain regions is less known.

The Braconidae are solitary or gregarious, internal or external parasitoids, first of all of Coleoptera, Lepidoptera, and Diptera, but also of Copeognatha, Heteroptera, Hymenoptera, Raphidioptera, Planipennia, and Mecoptera. Their larvae develop mostly in the host larvae or adults, but the larval development of few braconid groups begins in the host eggs already, while in some others it is finished in the host pupae

or adults. Pupation in a typical cocoon or within the mummified host skin.

Essentially the nomenclature and classification of the family as given in the "Hymenopterorum Catalogus n.ed." (Shenefelt, 1969—1978, Shenefelt & Marsh, 1976) with several modifications after Čapek (1969) and van Achterberg (1984) has been followed; for the subfamily Opiinae the catalogue by Fischer (1971), and for the subfamily Microgastrinae the revision by Mason (1981).

The subfamilies Opiinae and Alysiinae were arranged by the junior author, who complemented also further subfamilies with his findings. All other subfamilies as well as the whole check-list were adjusted by the senior author.

References: van Achterberg, C., 1984: Essay on the phylogeny of Braconidae (Hymenoptera: Ichneumonoidea). *Ent. Tidskr.* 105: 41—58. — Čapek, M., 1969: An attempt at a natural classification of the family Braconidae based on various unconventional characters. *Proc. Ent. Soc. Washington* 71: 304—312. — Fischer, M., 1971: Index of world Opiinae, in: Index of entomophagous insects, 5: 189 pp., Le Francois, Paris. — Mason, W. R. M., 1981: The polyphyletic nature of Apanteles Foerster (Hymenoptera: Braconidae): a phylogeny and reclassification of Microgastrinae. *Mem. Entomol. Soc. Canada* 115: 147 pp. — Shenefelt, R. D., 1969—1978: Braconidae 1—8: 1262 pp. and 10: 1425—1872, in: Hymenopterorum Catalogus n.ed., W. Junk, 's-Gravenhage. — Shenefelt, R. D. & Marsh, P. M., 1976: Braconidae 9: 1263—1424, l. c. — Tobias, V. I., 1986: Semejstvo Braconidae — brakonidy, in: Medvedev, G. S. [red.]: Opredelitel' nasekomykh Jevropejskoj časti SSSR, T. III, Perepončatokrylyje, 4. č.: 500 pp. — Tobias, V. I. & Jakimavičius, A. B.: Semejstvo Braconidae — brakonidy, l. c. 5. č.: 231 pp.

BRACONIDAE				
DORYCTINAE				
DORYCTINI				
<b>Doryctes</b> Haliday, 1836				
<i>fulviceps</i> Reinhard, 1865	S			
<i>heydenii</i> Reinhard, 1865	M S			
<i>leucogaster</i> (Nees, 1834)	B M S			
<i>mutillator</i> (Thunberg, 1822)	B M S			
<i>planiceps</i> Reinhard, 1865	*B M			
<i>pomarius</i> Reinhard, 1865	S			
<i>striatelloides</i> Strand, 1918	*S			
<i>undulatus</i> (Ratzeburg, 1852)	B M S			
<b>Ontsira</b> Cameron, 1900				
<i>antica</i> (Wollaston, 1858)	M S			
<i>igneus</i> (Ratzeburg, 1852)	M S			
<i>imperator</i> (Haliday, 1836)	B M S			
<b>Rhoptrocentrus</b> Marshall, 1897				
<i>piceus</i> Marshall, 1897	*S			
<b>Heterospilus</b> Haliday, 1836				
		ater Fischer, 1960		M S
		<i>cephi</i> Rohwer, 1925		*S
		<i>incompletus</i> (Ratzeburg, 1844)	B	S
		<i>leptosoma</i> Fischer, 1960		S
		<i>minimus</i> Fischer, 1960		*S
		<i>planus</i> (Ratzeburg, 1848)		*S
		<i>separatus</i> Fischer, 1960		*S
		<b>Rhaconotus</b> Ruthe, 1854		
		<i>aciculatus</i> Ruthe, 1854	M S	
		<b>Neurocrassus</b> Šnoflák, 1945		
		<i>tesari</i> Šnoflák, 1945	B	
		<b>Dendrosoter</b> Wesmael, 1838		
		<i>hartigii</i> (Ratzeburg, 1848)	B M S	
		<i>middendorffii</i> (Ratzeburg, 1848)	B M S	
		<b>SPATHIINI</b>		
		<b>Spathius</b> Nees, 1819		
		<i>brevicaudis</i> Ratzeburg, 1844	B	S

<i>curvicaudis</i> Ratzeburg, 1844	B M S	<b>Xenarcha</b> Foerster, 1862	
<i>dentatus</i> Telenga, 1941	S	<i>lustrator</i> (Haliday, 1836)	M S
<i>depressus</i> Hedqvist, 1976	*S	<b>Acrisis</b> Foerster, 1862	
<i>erythrocephalus</i>		<i>clavipes</i> Marshall, 1888	S
Wesmael, 1838	S	<b>Rhyssalus</b> Haliday, 1833	
<i>exarator</i> (Linnaeus, 1758)	B M S	<i>clavator</i> Haliday, 1833	*B S
<i>melanophilae</i> Fischer, 1966	*M	<b>Dolopsidea</b> Hincks, 1944	
<i>pedestris</i> Wesmael, 1838	M	<i>indagator</i> (Haliday, 1836)	S
<i>phyamatodis</i> Fischer, 1966	*M S	<b>Rhysipolis</b> Foerster, 1862	
<i>radzayanus</i> Ratzeburg, 1848	*M S	<i>decorator</i> (Haliday, 1836)	B *S
<i>rubidus</i> (Rossi, 1794)	B M S	<i>hariolator</i> (Haliday, 1836)	B M S
HECABOLINI		<i>intermedius</i> (Wesmael, 1838)	*B S
<b>Polystenus</b> Foerster, 1862		<i>meditator</i> (Haliday, 1836)	*S
<i>rugosus</i> Foerster, 1862	B M S	<i>variabilis</i> (Szépligeti, 1896)	S
<b>Pareucorystes</b> Tobias, 1961		<b>Oncophanes</b> Foerster, 1862	
<i>varinervis</i> Tobias, 1961	M S	<i>laevigatus</i> (Ratzeburg, 1852)	B M S
<b>Doryctosoma</b> Picard, 1938		<i>minutus</i> (Wesmael, 1838)	S
<i>transcaucasicum</i> (Tobias, 1976)	*S	<b>Noserus</b> Foerster, 1862	
<b>Hecabolus</b> Curtis, 1834		<i>flavicoxa</i> (Tobias, 1964)	*S
<i>sulcatus</i> Curtis, 1834	B M S	<i>similaris</i> (Szépligeti, 1896)	*S
<b>Ecphylus</b> Foerster, 1862		HORMIINI	
<i>silesiacus</i> (Ratzeburg, 1848)	B M S	<b>Lysitermus</b> Foerster, 1862	
<b>Monolexis</b> Foerster, 1862		<i>Rogadinaspis</i> Bouček, 1956	
<i>foersteri</i> Marshall, 1897	M	<i>pallidus</i> Foerster, 1862	B
<i>fuscicornis</i> Foerster, 1862	S	<i>tritoma</i> (Bouček, 1956)	
HISTEROMERINAE		<b>Hormius</b> Nees, 1818	
<b>Histeromerus</b> Wesmael, 1838		<i>moniliatus</i> (Nees, 1811)	B M S
<i>mystacinus</i> Wesmael, 1838	M	PAMBOLINI	
EXOTHECINAE		<b>Chremylus</b> Haliday, 1833	
EXOTHECINI		<i>elaphus</i> Haliday, 1833	B M S
<b>Colastes</b> Haliday, 1833		<b>Pambolus</b> Haliday, 1836	
<i>braconius</i> Haliday, 1833	B M S	<i>pallipes</i> Foerster, 1862	*M
<i>flavitarsis</i> (Thomson, 1891)	S	<i>rugulosus</i> (Hellén, 1927)	*S
<i>gracilis</i> Papp, 1975	*S	<i>tricolor</i> (Ruthe, 1854)	*B
<i>subquadratus</i> Papp, 1975	*S	GNAPDOTONTINAE	
<b>Shawiana</b> Achterberg, 1983		<b>Gnaptodon</b> Haliday, 1837	
<i>catenator</i> (Haliday, 1836)	*B	<i>breviradialis</i> Fischer, 1959	*S
<i>laevis</i> (Thomson, 1892)	*S	<i>decoris</i> (Foester, 1862)	*S
<i>lapponica</i> (Thomson, 1892)	S	<i>pumilio</i> (Nees, 1834)	B S

BRACONINAE				
IPHIAULACINI				
<b>Iphiaulax</b> Foerster, 1862				
<i>impeditor</i> (Kokujev, 1898)	S			
<i>impostor</i> (Scopoli, 1763)	B	S		
<i>mactator</i> (Klug, 1817)		S		
<i>umbraculator</i> (Nees, 1834)	M	S		
ISOMECINI				
<b>Pseudovipio</b> Szépligeti, 1896				
<i>castrator</i> (Fabricius, 1798)	M	S		
<i>inscriptor</i> (Nees, 1834)		S		
<i>variegata</i> (Bohemian, 1851)	B	S		
<b>Cyanopterus</b> Haliday, 1835				
<i>curvatus</i> (Telenga, 1936)	*B			
<i>extricator</i> (Nees, 1834)		*M	S	
<i>flavator</i> (Fabricius, 1793)		S		
<i>nigrator</i> (Zetterstedt, 1838)		*S		
<i>rector</i> (Thunberg, 1822)	M	S		
<i>samedovi</i> (Abdinbekova, 1973)		*S		
<b>Bracomorpha</b> Papp, 1971				
<i>torkai</i> Papp, 1971		*S		
<b>Isomecus</b> Kriechbaumer, 1895				
<i>appellator</i> (Nees, 1834)	B	M	S	
<i>insectator</i> (Kokujev, 1898)		M	S	
<i>intermedius</i> (Szépligeti, 1896)		S		
<i>nominator</i> (Fabricius, 1787)	B	M	S	
<i>nomioides</i> (Shestakov, 1926)		*S		
<i>siculus</i> (Marshall, 1888)		*S		
<i>simulator</i> (Kokujev, 1898)		*S		
<i>tentator</i> (Rossi, 1790)	B	M	S	
<i>terrefactor</i> (Villers, 1789)		M	S	
<b>Atanycolus</b> Foerster, 1862				
<i>denigrator</i> (Linnaeus, 1758)		S		
<i>fulviceps</i> Kriechbaumer, 1898		M		
<i>initiator</i> (Fabricius, 1793)	B	M	S	
<i>neesii</i> Marshall, 1898			B	M
<i>sculpturatus</i> (Thomson, 1892)			S	
BRACONINI			B	M
<b>Baryproctus</b> Ashmead, 1900			S	
<i>barypus</i> (Marshall, 1885)	*B			
<i>hungaricus</i> Szépligeti, 1901		B	S	
<b>Coeloides</b> Wesmael, 1838				
<i>abdominalis</i> (Zetterstedt, 1840)		B	M	S
<i>bostrichorum</i> Giraud, 1872	B	M	S	
<i>filiformis</i> Ratzeburg, 1852	B	M	S	
<i>forsteri</i> Haeselbarth, 1967		S		
<i>melanotus</i> Wesmael, 1838	B	S		
<i>rossicus</i> (Kokujev, 1902)	M	S		
<i>scolyticida</i> Wesmael, 1838	B	S		
<i>sordidator</i> (Ratzeburg, 1844)		S		
<i>ungularis</i> Thomson, 1892	B	M		
<b>Bracon</b> Fabricius, 1804				
<i>abbreviator</i> Nees, 1834	B			
<i>aestivalis</i> Szépligeti, 1901		S		
<i>anthracinus</i> Nees, 1834	M	S		
<i>atrator</i> Nees, 1834		*S		
<i>brachycerus</i> Thomson, 1892		S		
<i>carbonarius</i> Szépligeti, 1901		*S		
<i>caudatus</i> Ratzeburg, 1848		*B		
<i>caudiger</i> Nees, 1834	B			
<i>colpophorus</i> Wesmael, 1838		*M		
<i>coniferarum</i> Fahringer, 1928		M		
<i>conjugellae</i> Bengtsson, 1924		*S		
<i>dichromus</i> Wesmael, 1838		S		
<i>discoideus</i> Wesmael, 1838	B	S		
<i>epitriptus</i> Marshall, 1885		*S		
<i>erraticus</i> Wesmael, 1838		*S		
<i>exhilarator</i> Nees, 1834		*S		
<i>fulvipes</i> Nees, 1834		*S		
<i>fuscicoxis</i> Wesmael, 1838		*S		
<i>guttiger</i> Wesmael, 1838	*B			
<i>hungaricus</i> (Szépligeti, 1896)		*S		

<i>hylobii</i> Ratzeburg, 1848	B M	<i>palpebrator</i> (Ratzeburg, 1844)	B
<i>immutator</i> Nees, 1834	B M	<i>stabilis</i> (Wesmael, 1838)	B M S
<i>intercessor</i> Nees, 1834	*S	<i>variegator</i> (Spinola, 1808)	M S
<i>laetus</i> Wesmael, 1838	B	<i>viktorovi</i> Tobias, 1961	*S
<i>larvicida</i> Wesmael, 1838	*S		
<i>leptus</i> Marshall, 1897	*S	<b>Rostrobracon</b> Tobias, 1957	
<i>luteator</i> Spinola, 1808	*S	<i>urinator</i> (Fabricius, 1798)	*S
<i>maculiger</i> Wesmael, 1838	B S	<b>Pigeria</b> van Achterberg, 1985	
<i>mariae</i> Dalla Torre, 1898	*S	<i>piger</i> (Wesmael, 1838)	*S
<i>mediator</i> Nees, 1834	B M S		
<i>megapterus</i> Wesmael, 1838	M		
<i>minutator</i> (Fabricius, 1798)	*M S		
<i>monticola</i> Kokujev, 1899	S		
<i>nigratus</i> Wesmael, 1838	B S	<b>Aleiodes</b> Wesmael, 1838	
<i>nigriventris</i> Wesmael, 1838	*S	<i>alpinus</i> (Thomson, 1891)	?
<i>nitidifrons</i> Nieuzeitowski, 1910	*S	<i>armatus</i> Wesmael, 1838	B
<i>obscuator</i> Nees, 1834	B S	<i>bicolor</i> (Spinola, 1808)	B M S
<i>osculator</i> Nees, 1812	B M S	<i>cantherius</i> (Lyle, 1919)	S
<i>pectoralis</i> Wesmael, 1838	*S	<i>carbonarius</i> Giraud, 1857	M S
<i>pellucidus</i> Ratzeburg, 1852	B	<i>caudalis</i> Hellén, 1927	S
<i>picticornis</i> Wesmael, 1838	S	<i>circumscripitus</i> (Nees, 1834)	B M S
<i>pineti</i> Thomson, 1892	B S	<i>cruentus</i> (Nees, 1834)	B M S
<i>praetermissus</i> Marshall, 1885	*S	<i>dimidiatus</i> (Spinola, 1808)	B M S
<i>rhyngchiti</i> Grese, 1928	S	<i>dispar</i> (Curtis, 1834)	M S
<i>romani</i> Fahringer, 1927	*S	<i>dissector</i> (Nees, 1834)	B M
<i>rufipalpis</i> Szépligeti, 1901	*S	<i>ductor</i> (Thunberg, 1822)	B M S
<i>sphaerocephalus</i> Szépligeti, 1901	*S	<i>excavatus</i> (Telenga, 1941)	B M S
<i>sulphurator</i> Szépligeti, 1896	S	<i>farmakena</i> Maláč, 1941	
<i>tener</i> Szépligeti, 1904	*S	<i>gasterator</i> (Jurine, 1807)	M S
<i>terebella</i> (Wesmael, 1838)	*S	<i>geniculator</i> (Nees, 1834)	M S
<i>titubans</i> Wesmael, 1838	*S	<i>grandis</i> Giraud, 1857	M
<i>triangularis</i> Nees, 1834	B S	<i>medianus</i> (Thomson, 1896)	*S
<i>variator</i> Nees, 1812	B S	<i>miniatus</i> (Herrich et Schaeffer, 1838)	B M S
<i>xanthogaster</i> Nees, 1834	B	<i>modestus</i> (Reinhard, 1863)	S
<b>Foveobracon</b> Tobias, 1961		<i>nigricornis</i> Wesmael, 1838	M S
<i>biimpressus</i> (Telenga, 1936)	S	<i>pallidator</i> (Thunberg, 1822)	S
<b>Habrobracon</b> Ashmead, 1895		<i>pallidicornis</i> (Herrich et Schaeffer, 1838)	M
<i>brevicornis</i> (Wesmael, 1838)	B M S	<i>periscelis</i> (Reinhard, 1863)	M
<i>hebetor</i> (Say, 1836)	B M S	<i>praerogator</i> (Nees, 1812)	S
<i>nigricans</i> Szépligeti, 1901	*S	<i>pulchripes</i> Wesmael, 1838	B
		<i>rossicus</i> Kokujev, 1898	*S
		<i>rugulosus</i> (Nees, 1812)	*S
		<i>testaceus</i> (Spinola, 1808)	S
		<i>tristis</i> Wesmael, 1838	B M S
		<i>ungularis</i> Thomson, 1891	*S
		<i>unipunctator</i> (Thunberg, 1822)	M S

<i>vittiger</i> Wesmael, 1838	S	<i>cubitalis</i> Fischer, 1959	S
<b>Petalodes</b> Wesmael, 1838		<i>curtifemur</i> Fischer, 1961	B S
<i>unicolor</i> Wesmael, 1838	B M S	<i>curtipectus</i> Fischer, 1958	B *S
<b>Clinocentrus</b> Haliday, 1833		<i>curvatus</i> Fischer, 1957	S
<i>cunctator</i> Haliday, 1836	B M S	<i>decoratus</i> Stelfox, 1949	
<i>excubitor</i> Haliday, 1836	*S	<i>depeculator</i> (Foerster, 1862)	*B S
<i>exsertor</i> (Nees, 1812)	B M S	<i>docilis</i> Haliday, 1837	S
<i>tenuicornis</i> Thomson, 1819	?	<i>exiguus</i> Westmael, 1835	S
<i>umbratilis</i> Haliday, 1833	*S	<i>exilis</i> Haliday, 1837	S
<b>Rogas</b> Nees, 1818		<i>eximius</i> Fischer, 1957	S
<i>luteus</i> Nees, 1834	B M S	<i>exsertus</i> Thomson, 1895	S
<i>tricolor</i> (Wesmael, 1838)	S	<i>fasciatus</i> Thomson, 1895	?
<b>OPIINAE</b>		<i>ficedus</i> Papp, 1979	S
<b>Ademon</b> Haliday, 1833		<i>filicornis</i> Thomson, 1895	*S
<i>decrescens</i> (Nees, 1812)	?	<i>funebris</i> Wesmael, 1835	*S
<b>Cephaloplites</b> Szépligeti, 1897		<i>fuscipennis</i> Wesmael, 1835	B
<i>mocsaryi</i> Szépligeti, 1897	M	<i>geniculatus</i> Thomson, 1895	S
<b>Opius</b> Wesmael, 1835		<i>gracilis</i> Fischer, 1957	S
<i>abditiformis</i> Fischer, 1984	S	<i>helios</i> Fischer, 1959	M
<i>aemulus</i> Haliday, 1837	*S	<i>hilaris</i> Fischer, 1963	*S
<i>aethiops</i> Haliday, 1837	B	<i>ilicis</i> Nixon, 1939	S
<i>altimontanus</i> Fischer, 1969	*S	<i>impatientis</i> Fischer, 1957	M
<i>ambiguus</i> Wesmael, 1835	S	<i>instabilis</i> Wesmael, 1835	*S
<i>attributus</i> Fischer, 1962	S	<i>irregularis</i> Wesmael, 1835	S
<i>aureliae</i> Fischer, 1957	*S	<i>kostolnaensis</i> Fischer, 1984	S
<i>austriacus</i> Fischer, 1958	S	<i>laetatorius</i> Fischer, 1958	M S
<i>bajulus</i> Haliday, 1837	?	<i>latipes</i> Fischer, 1958	B S
<i>basalis</i> Fischer, 1958	*S	<i>leptostigma</i> Wesmael, 1835	?
<i>basirufus</i> Fischer, 1958	B	<i>levis</i> Wesmael, 1835	S
<i>bouceki</i> Fischer, 1958	B	<i>longicornis</i> Thomson, 1895	*S
<i>breviscapus</i> Thomson, 1895	*S	<i>lonicerae</i> Fischer, 1958	M S
<i>caesus</i> Haliday, 1837	S	<i>lucidus</i> Szépligeti, 1896	B
<i>campanariae</i> Fischer, 1959	*S	<i>lugens</i> Haliday, 1837	S
<i>capeki</i> Fischer, 1963	S	<i>lukasi</i> Fischer, 1984	S
<i>caudatus</i> Wesmael, 1835	*S	<i>macrocerus</i> Thomson, 1895	S
<i>caudifer</i> Fischer, 1958	B	<i>maculipes</i> Wesmael, 1835	S
<i>cingulatus</i> Wesmael, 1835	S	<i>magnicauda</i> Fischer, 1958	B S
<i>circulator</i> (Nees, 1834)	*S	<i>minor</i> Fischer, 1957	S
<i>clausus</i> Fischer, 1958	*S	<i>moravicus</i> Fischer, 1960	M
<i>comatus</i> Wesmael, 1835	*S	<i>mundus</i> Foerster, 1862	?
<i>compar</i> Marshall, 1894	B	<i>nitidulator</i> (Nees, 1834)	S
<i>connivens</i> Thomson, 1895	*S	<i>ocellatus</i> Wesmael, 1835	S
<i>coracinus</i> Thomson, 1895	S	<i>ochrogaster</i> Wesmael, 1835	B M S
<i>crassicrus</i> Thomson, 1895	M S	<i>pactus</i> Haliday, 1837	S
<i>crassipes</i> Wesmael, 1835	S	<i>pallipes</i> Wesmael, 1835	M S
		<i>parvungula</i> Thomson, 1895	*S

<i>pendulus</i> Haliday, 1837	S	<i>haemorrhoeus</i> (Haliday, 1837)	S
<i>piceus</i> Thomson, 1895	S	<i>longicauda</i> (Thomson, 1895)	S
<i>polyzonius</i> Wesmael, 1835	?	<i>punctiscuta</i> (Thomson, 1895)	*S
<i>propodealis</i> Fischer, 1958	B	<i>rusticus</i> (Haliday, 1837)	M S
<i>pulchriventris</i> Fischer, 1958	B	<i>scabriculus</i> (Wesmael, 1835)	?
<i>pulicariae</i> Fischer, 1969	*S	<i>spinaciae</i> (Thomson, 1895)	M S
<i>pygmaeator</i> (Nees, 1811)	S	<i>wesmaeli</i> (Haliday, 1837)	S
<i>pygmaeus</i> Fischer, 1962	*S	<b>Diachasma</b> Foerster, 1862	
<i>radialis</i> Fischer, 1957	S	<i>caffer</i> (Wesmael, 1835)	S
<i>reconditor</i> Wesmael, 1835	S	<i>cephalotes</i> (Wesmael, 1835)	*S
<i>rex</i> Fischer, 1958	M S	<b>Pokomandy</b> Fischer, 1959	
<i>rhaagoleticolus</i> Sachtleben, 1934	B	<i>curticornis</i> Fischer, 1959	S
<i>rotundiventris</i> Thomson, 1895	S	ALYSIINAE	
<i>rudiformis</i> Fischer, 1958	B	ALYSINI	
<i>rudis</i> Wesmael, 1835	M S	<b>Alloea</b> Haliday, 1833	
<i>rufipes</i> Wesmael, 1835	S	<i>bonessi</i> Fischer, 1966	*S
<i>similis</i> Szépligeti, 1896	S	<i>lonchopterae</i> Fischer, 1966	*S
<i>similoides</i> Fischer, 1962	*S	<b>Alysia</b> Latreille, 1804	
<i>singularis</i> Wesmael, 1835	S	<i>curtungula</i> Thomson, 1895	*S
<i>snoflaki</i> Fischer, 1959	M	<i>diversiceps</i> Fischer, 1967	*S
<i>soenderupianus</i> Fischer, 1967	*S	<i>frigida</i> Haliday, 1838	*S
<i>speciosus</i> Fischer, 1959	*S	<i>fuscipennis</i> Haliday, 1838	*S
<i>spretus</i> Haliday, 1837	S	<i>incongrua</i> Nees, 1834	*S
<i>staryi</i> Fischer, 1958	M S	<i>lucicola</i> Haliday, 1838	*S
<i>subaffinis</i> Fischer, 1962	*S	<i>luciella</i> Stelfox, 1941	*S
<i>tersus</i> (Foerster, 1862)	S	<i>manducator</i> (Panzer, 1799)	S
<i>testaceus</i> Wesmael, 1838	?	<i>obscuripes</i> Thomson, 1895	*S
<i>thomsoni</i> Fischer, 1971	B	<i>rudis</i> Tobias, 1962	S
<i>tirolensis</i> Fischer, 1958	*S	<i>rufidens</i> Nees, 1834	*S
<i>trencensis</i> Fischer, 1984	S	<i>sophia</i> Haliday, 1835	*S
<i>trisulcus</i> Thomson, 1895	S	<b>Anisocyrt</b> Foerster, 1862	
<i>truncatus</i> Wesmael, 1835	?	<i>alpinicola</i> van Achtenberg, 1986	B
<i>tuberculifer</i> Fischer, 1958	*S	<i>perdita</i> (Haliday, 1838)	S
<i>turcicus</i> Fischer, 1960	*S	<b>Aphaereta</b> Foerster, 1862	
<i>variegatus</i> Szépligeti, 1896	S	<i>brevis</i> Tobias, 1962	S
<i>viennensis</i> Fischer, 1959	M	<i>major</i> Thomson, 1895	*S
<i>zelotes</i> Marshall, 1891	S	<i>minuta</i> (Nees, 1812)	S
<b>Biosteres</b> Foerster, 1862	S	<i>tenuicornis</i> Nixon, 1939	*S
<i>blandus</i> (Haliday, 1837)	S	<b>Chasmodon</b> Haliday, 1838	
<i>brevipalpis</i> (Thomson, 1895)	S		
<i>brevisulcus</i> (Thomson, 1895)	*S		
<i>carbonarius</i> (Nees, 1834)	B M S		
<i>caudatulus</i> (Thomson, 1895)	S		

<i>apterus</i> (Nees, 1814)	M	<i>semicompressa</i> Stelfox et Graham, 1949	*S
<b>Asobara</b> Foerster, 1862		<i>variabilis</i> Tobias, 1962	*S
<i>tabida</i> (Nees, 1834)	S	<i>varipes</i> Tobias, 1962	*S
<b>Aspilota</b> Foerster, 1862		<i>venusta</i> Tobias, 1962	*S
<i>areolata</i> Stelfox et Graham, 1950	*S	<b>Cratospila</b> Foerster, 1862	
<i>betae</i> Bengtsson, 1926	*S	<i>circe</i> (Haliday, 1838)	S
<i>brevicauda</i> Tobias, 1962	*S	<b>Adelurola</b> Strand, 1928	
<i>brevicornis</i> (Nees, 1814)	B	<i>florimela</i> (Haliday, 1838)	*S
<i>brevissimicornis</i> Stelfox et Graham, 1948	*S	<b>Dapsilarthra</b> Foerster, 1862	
<i>carinata</i> Tobias, 1962	*S	<i>apii</i> (Curtis, 1826)	*S
<i>caudata</i> Thomson, 1895	*S	<i>dictynna</i> (Marshall, 1895)	B S
<i>concinna</i> (Haliday, 1838)	*S	<i>fuscula</i> Griffiths, 1968	*S
<i>contracticornis</i> Fischer, 1973	*S	<i>gahani</i> (Baume-Pluvine, 1915)	*S
<i>costulata</i> Thomson, 1895	*S	<i>phoridarum</i> (Goidanich, 1936)	*S
<i>cratocera</i> Thomson, 1895	*S	<i>rufiventris</i> (Nees, 1814)	M S
<i>daemon</i> Stelfox et Graham, 1948	*S	<i>sylvia</i> (Haliday, 1839)	*S
<i>denticulata</i> Stelfox et Graham, 1951	*S	<b>Idiasta</b> Foerster, 1862	
<i>dimidiata</i> (Thomson, 1895)	*S	<i>annulicornis</i> Thomson, 1895	*S
<i>divisa</i> Stelfox et Graham, 1950	*S	<i>nephela</i> (Haliday, 1838)	*S
<i>efoveolata</i> Thomson, 1895	*S	<i>paramaritima</i> Königsmann, 1960	*S
<i>erythropa</i> (Foerster, 1862)	*S	<i>picticornis</i> (Ruthe, 1854)	*S
<i>fuscicornis</i> (Haliday, 1838)	*S	<b>Idiolexis</b> Foerster, 1862	
<i>hirticornis</i> Thomson, 1895	S	<i>punctigera</i> (Haliday, 1838)	S
<i>inops</i> (Foerster, 1862)	S	<b>Orthostigma</b> Ratzeburg, 1844	
<i>insidiatrix</i> Marshall, 1895	*S	<i>antennatum</i> Tobias, 1962	*S
<i>insolita</i> (Tobias, 1962)	*S	<i>breviradiale</i> Königsmann, 1969	B M S
<i>jaculans</i> (Haliday, 1838)	S	<i>cratospilum</i> Thomson, 1895	B S
<i>laevinota</i> Tobias, 1962	*S	<i>laticeps</i> Thomson, 1895	B M S
<i>macrocera</i> Thomson, 1895	*S	<i>longicorne</i> Königsmann, 1969	M
<i>macrops</i> Stelfox et Graham, 1951	*S	<i>longicubitale</i> Königsmann, 1969	B S
<i>macrura</i> Thomson, 1895	*S	<i>lucidum</i> Königsmann, 1969	*S
<i>microcera</i> Thomson, 1895	*S	<i>mandibulare</i> (Tobias, 1962)	M S
<i>naevia</i> Tobias, 1962	*S	<i>pseudolaticeps</i> Königsmann, 1969	B S
<i>nervosa</i> (Haliday, 1833)	*S	<i>pumilum</i> (Nees, 1834)	B M S
<i>nigricornis</i> (Thomson, 1895)	*S	<i>sculpturatum</i> Tobias, 1962	*S
<i>oleracea</i> Tobias, 1962	*S		
<i>propodealis</i> Tobias, 1962	*S		
<i>pulvinata</i> Stelfox et Graham, 1949	*S		
<i>ruficornis</i> (Nees, 1834)	*S		

<i>sordipes</i> Thomson, 1895	B	S	<i>clandestina</i> (Haliday, 1839)	?
<b>Pentapleura</b> Foerster, 1862			<b>Aristelix</b> Nixon, 1943	
<i>fuliginosa</i> (Haliday, 1838)	*S		<i>phaenicura</i> (Haliday, 1839)	S
<i>laevipleuris</i> (Tobias, 1962)	*S		<b>Chaenusa</b> Haliday, 1839	
<i>pumilio</i> (Nees, 1814)	*S		<i>conjugens</i> (Nees, 1812)	?
<b>Phaenocarpa</b> Foerster, 1862			<b>Chorebus</b> Haliday, 1833	
<i>angustiptera</i> Papp, 1968	S		<i>gracilis</i> (Nees, 1834)	*S
<i>collaris</i> Papp, 1968	*S		<i>petiolatus</i> (Nees, 1834)	S
<i>conspurcator</i> (Haliday, 1838)	*S		<i>rufipes</i> (Nees, 1814)	B
<i>eugenia</i> (Haliday, 1838)	*S		<i>talaris</i> (Haliday, 1839)	?
<i>eunice</i> (Haliday, 1838)	*S		<b>Coelinius</b> Nees, 1818	
<i>flavipes</i> (Haliday, 1838)	S		<i>circulator</i> (Gravenhorst, 1807)	M
<i>laticellula</i> Papp, 1968	*S		<i>gracilis</i> (Curtis, 1829)	*S
<i>livida</i> (Haliday, 1838)	*S		<i>niger</i> (Nees, 1812)	*S
<i>plicinervis</i> (Haliday, 1838)	*S		<b>Chaenon</b> Curtis, 1829	
<i>pullata</i> (Haliday, 1838)	S		<i>anceps</i> Curtis, 1829	S
<i>ruficeps</i> (Nees, 1814)	S		<b>Coloneura</b> Foerster, 1862	
<i>tacita</i> Stelfox, 1941	*S		<i>dice</i> (Nixon, 1943)	?
<i>theodori</i> (Snellen van Vollenhoven, 1878)	*S		<i>stylata</i> Foerster, 1862	?
<b>Prosapha</b> Foerster, 1862			<b>Dacnusa</b> Haliday, 1833	
<i>speculum</i> (Haliday, 1838)	*S		<i>adducta</i> Haliday, 1839	S
<b>Synaldis</b> Foerster, 1862			<i>annulata</i> (Nees, 1834)	*S
<i>concolor</i> (Nees, 1814)	*S		<i>discolor</i> (Foerster, 1862)	?
<i>distracta</i> (Nees, 1834)	*S		<i>faeroensis</i> (Roman, 1917)	B
<i>globipes</i> Fischer, 1962	*S		<i>lugens</i> Haliday, 1839	?
<i>maxima</i> Fischer, 1962	*S		<i>maculipes</i> Thomson, 1895	*S
<i>sulcata</i> Fischer, 1962	*S		<i>pubescens</i> (Curtis, 1826)	B
<b>Trisynaldis</b> Fischer, 1958			<i>temula</i> Haliday, 1839	*M
<i>confuncta</i> Fischer, 1958	?		<b>Epimicta</b> Foerster, 1862	
<b>Tanycarpa</b> Foerster, 1862			<i>marginalis</i> (Haliday, 1839)	?
<i>bicolor</i> (Nees, 1814)	*S		<b>Exotela</b> Foerster, 1862	
<i>gracilicornis</i> (Nees, 1814)	S		<i>gilvipes</i> (Haliday, 1839)	B
<i>mitis</i> Stelfox, 1941	*S		<b>Laotris</b> Nixon, 1943	
<b>Trachyusa</b> Ruthe, 1854			<i>striatula</i> (Haliday, 1839)	?
<i>aurora</i> (Haliday, 1838)	S		<b>Polemochartus</b> Schulz, 1911	
<b>Apronopa</b> Achterberg, 1980			<i>liparae</i> (Giraud, 1863)	?
<i>haeselbarthi</i> van Achterberg, 1980	*S		<i>melas</i> (Giraud, 1863)	?
<b>DACNUSINI</b>			<b>Protodacnusa</b> Griffiths, 1964	
<b>Amyras</b> Nixon, 1943			<i>aridula</i> (Thomson, 1895)	*S
			<i>tristis</i> (Nees, 1834)	*S

<b>Symphya</b> Foerster, 1862						
<i>hians</i> (Nees, 1816)	*S					*S
<i>mandibularis</i> (Nees, 1814)	*S					*S
<b>Synelix</b> Foerster, 1862						
<i>rea</i> (Nixon, 1942)	?					
HELCONINAE						
HELCONINI						
<b>Helcon</b> Nees, 1814						
<i>angustator</i> Nees 1814	B M					
<i>tardator</i> Nees, 1814	B M S					
<b>Helconidea</b> Viereck, 1914						
<i>dentator</i> (Fabricius, 1804)	B M S					
<i>ruspator</i> (Linnaeus, 1758)	S					
<b>Aspicolpus</b> Wesmael, 1838						
<i>carinator</i> (Nees, 1814)	B M S					
<i>maximus</i> (Szépligeti 1900)	M S					
CENOCOELIINI						
<b>Cenocoelius</b> Westwood, 1840						
<i>analis</i> (Nees, 1834)	B M S					
<b>Lestricus</b> Reinhard, 1865						
<i>secalis</i> (Linnaeus, 1758)	B M S					
DIOSPILINI						
<b>Baeacis</b> Foerster, 1878						
<i>abietis</i> (Ratzeburg, 1844)	M S					
<b>Diospilus</b> Haliday, 1833						
<i>capito</i> (Nees, 1834)	S					
<i>dilatatus</i> Thomson, 1895	*S					
<i>molorchicola</i> Fischer, 1966	M S					
<i>morosus</i> Reinhard, 1862	*S					
<i>oleraceus</i> Haliday, 1833	*S					
<i>robustus</i> Reinhard, 1862	S					
<i>rufipes</i> Reinhard, 1862	*S					
<i>rugosus</i> Thomson, 1895	M					
<b>Dyscoletes</b> Haliday, 1840						
<i>lancifer</i> (Haliday, 1836)	*B	S				
<b>Taphaeus</b> Wesmael, 1835						
<i>hiator</i> (Thunberg, 1822)	*S					
BRACHISTINI						
<b>Eubazus</b> Nees, 1814						
<i>atricornis</i> (Ratzeburg, 1848)			S			
<b>augustinus</b> (Ruthe, 1867)						*S
<b>caudatus</b> (Niezabitowski, 1910)						*S
<b>claviventris</b> (Ruthe, 1867)						*S
<b>corrugatus</b> (Ruthe, 1867)						*S
<b>cracoviensis</b>						
(Niezabitowski, 1910)						*S
<b>exsertor</b> (Reinhard, 1867)						S
<b>firmus</b> (Ratzeburg, 1844)	B					
<b>fuscipalpis</b> (Wesmael, 1835)						*S
<b>gallicus</b> (Reinhard, 1867)						S
<b>longicaudis</b> (Ratzeburg, 1844)	B					S
<b>macrocephalus</b> Nees, 1814	B					S
<b>minutus</b> (Ratzeburg, 1848)						*S
<b>mucronatus</b> (Thomson, 1892)	B					
<b>opacus</b> (Reinhard, 1867)						*S
<b>parvulus</b> (Ruthe, 1867)						*S
<b>politus</b> (Ratzeburg, 1852)						*S
<b>ruficoxis</b> (Wesmael, 1835)						*S
<b>rufipes</b> (Herrich et Schaeffer, 1838)						*S
<b>rugosus</b> (Ratzeburg, 1848)						*S
<b>tibialis</b> (Haliday, 1835)						*S
<b>vitripennis</b> (Herrich et Schaeffer 1838)						*S
<b>Aliolus</b> Say, 1836						
<b>breviseta</b> (Šnoflák, 1952)	M					
<b>glypturus</b> (Thomson, 1874)						S
<b>hofferi</b> (Šnoflák, 1952)						M S
<b>lepidus</b> (Haliday, 1835)						*S
<b>obtusus</b> (Šnoflák, 1952)						M
<b>semirugosus</b> (Nees, 1816)						*S
<b>sudeticus</b> (Šnoflák, 1952)	B					
<b>Foersteria</b> Szépligeti, 1896						
<b>flavipes</b> Szépligeti, 1896						*S
<b>Polydegmon</b> Foerster, 1862						
<b>foveolatus</b> (Herrich et Schaeffer, 1838)						*S
<b>sinuatus</b> Foerster, 1862						S
<b>Dicyrtaspis</b>						
van Achterberg, 1980						
<b>cavifrons</b> (Šnoflák, 1952)	B					
<b>Schizoprymnus</b> Foerster, 1862						

<i>acataphractus</i> Šnoflák, 1952	M	<i>flavipalpis</i> (Wesmael, 1835)	M
<i>ambiguus</i> (Nees, 1816)	B	<i>flavipes</i> (Ivanov, 1899)	S
<i>angustatus</i> (Herrick et Schaeffer 1838)	S	<i>floricola</i> (Wesmael, 1835)	S
<i>angustissimus</i> Šnoflák, 1952	B M S	<i>gibberosus</i> (Szépligeti, 1901)	S
<i>bidentulus</i> (Szépligeti, 1901)	S	<i>lugubris</i> Šnoflák, 1952	M
<i>cataphractus</i> Šnoflák, 1952	M S	<i>obscurellus</i> (Nees, 1816)	S
<i>crassiceps</i> (Thomson, 1892)	*S	<i>pallipes</i> (Nees, 1816)	*S
<i>cylindricus</i> Šnoflák, 1952	M	<i>sekerai</i> Šnoflák, 1952	M
<i>emarginatus</i> Šnoflák, 1952	M S	<i>semilissus</i> Šnoflák, 1952	M
<i>eurygaster</i> (Šnoflák, 1952)	B	<i>striatulus</i> (Nees, 1816)	B M S
<i>glaberrimus</i> (Šnoflák, 1952)	B M	<i>thomsoni</i> Fahringer, 1934	M
<i>globosus</i> (Szépligeti, 1898)	*S	<i>thoracicus</i> (Curtis, 1860)	M S
<i>glypturus</i> (Thomson, 1874)	S		
<i>gregori</i> Šnoflák, 1952	M	BLACINI	
<i>hilaris</i> (Herrick et Schaeffer, 1838)	S	<b>Blacus</b> Nees, 1818	
<i>luteipalpis</i> Šnoflák, 1952	B M	<i>achterbergi</i> Haeselbarth, 1976	M
<i>nigripes</i> (Thomson, 1892)	S	<i>ambulans</i> Haliday, 1835	B M S
<i>obscurus</i> (Nees, 1816)	*S	<i>armatus</i> Ruthe, 1861	S
<i>opacus</i> (Thomson, 1891)	S	<i>bovistae</i> Haeselbarth, 1973	*S
<i>pallidipennis</i> (Herrick et Schaeffer 1838)	S	<i>capeki</i> Haeselbarth, 1973	S
<i>parvus</i> (Thomson, 1892)	S	<i>diversicornis</i> (Nees, 1834)	B M S
<i>rufipes</i> (Herrick et Schaeffer, 1838)	S	<i>errans</i> (Nees, 1812)	B M S
<i>sculpturatus</i> Šnoflák, 1952	M S	<i>exilis</i> (Nees, 1812)	B M S
<i>stenopygus</i> Šnoflák, 1952	M	<i>filicornis</i> Haeselbarth, 1973	M
<i>telengai</i> Tobias, 1976	*S	<i>fischeri</i> Haeselbarth, 1973	*S
<i>terebralis</i> Šnoflák, 1952	M S	<i>hastatus</i> Haliday, 1835	B M S
<b>Triaspis</b> Haliday, 1835		<i>humilis</i> (Nees, 1812)	B M S
<i>aciculatus</i> (Ratzeburg, 1848)	B M S	<i>instabilis</i> Ruthe, 1861	B S
<i>brevicornis</i> (Herrick et Schaeffer 1838)	S	<i>interstitialis</i> Ruthe, 1861	M S
<i>caledonicus</i> (Marshall, 1888)	S	<i>longipennis</i> (Gravenhorst, 1809)	S
<i>caudatus</i> (Nees, 1816)	B M S	<i>maculipes</i> Wesmael, 1835	B M S
<i>coniungens</i> Šnoflák, 1952	M	<i>mamillanus</i> Ruthe, 1861	B S
<i>devinensis</i> Šnoflák, 1952	M	<i>nigricornis</i> Haeselbarth, 1973	S
<i>excisus</i> Šnoflák, 1952	B	<i>paganus</i> Haliday, 1835	S
		<i>pallipes</i> Haliday, 1835	B M S
		<i>pectinatus</i> Haeselbarth, 1973	B
		<i>robustus</i> Haeselbarth, 1973	S
		<i>rufescens</i> Ruthe, 1861	*M S
		<i>ruficornis</i> (Nees, 1812)	B M S

<i>tripudians</i> Haliday, 1835	M S	<i>Orgilus</i> s. str.	
<b>Neoblacus</b> Ashmead, 1900		<i>anurus</i> Thomson, 1895	*S
<i>koenigi</i> Fischer, 1962	M S	<i>dovnari</i> Tobias, 1986	*B
<b>Blacometeorus</b> Tobias, 1976		<i>grunini</i> Tobias, 1986	*S
<i>intermedius</i> Tobias, 1976	M S	<i>ischnus</i> Marshall, 1898	*B*M*S
MACROCENTRINAE		<i>leptocephalus</i> (Hartig, 1838)	B M S
<b>Macrocentrus</b> Curtis, 1833		<i>minutus</i> Szépligeti, 1898	*S
<i>bicolor</i> Curtis, 1833	S	<i>moldavicus</i> Tobias, 1986	*S
<i>blandus</i> Eady et Clark, 1964	B	<i>nigripennis</i> (Dahl, 1912)	M
<i>buolianae</i> Eady et Clark, 1964	S	<i>obscurator</i> (Nees, 1812)	B M S
<i>cingulum</i> Reinhard, 1882	*S	<i>parvipennis</i> Thomson, 1895	*S
<i>collaris</i> (Spinola, 1808)	M S	<i>pimpinellae</i> Niezabitowski, 1910	B M S
<i>flavus</i> Snellen van Vollenhoven, 1878	S	<i>punctulator</i> (Nees, 1812)	M S
<i>gibber</i> Eady et Clark, 1964	S	<i>rubrator</i> (Ratzeburg, 1852)	S
<i>hungaricus</i> Marshall, 1893	S	<i>rudolphae</i> Tobias, 1976	*M*S
<i>infirmus</i> (Nees, 1834)	*B	<i>ruficornis</i> Szépligeti, 1896	*B
<i>intricator</i> (Ratzeburg, 1852)	B	<i>rufifemur</i> Fahringer, 1937	*B
<i>kurnakovi</i> Tobias, 1976	S	<i>Ischiolus</i> Hellén, 1958	
<i>linearis</i> (Nees, 1812)	B M S	<i>rugosus</i> (Nees, 1834)	M S
<i>marginator</i> (Nees, 1812)	B M S	<i>punctatus</i> (Beyr, 1959)	
<i>nidulator</i> (Nees, 1834)	S	<b>Microtypus</b> Ratzeburg, 1848	
<i>nitidus</i> (Wesmael, 1835)	B S	<i>wesmaelii</i> Ratzeburg, 1848	M S
<i>pallipes</i> (Nees, 1811)	M S	AGATHIDINAE	
<i>resinellae</i> (Linnaeus, 1758)	B S	<b>Disophrys</b> Foerster, 1862	
<i>thoracicus</i> (Nees, 1812)	B S	<i>inculcator</i> (Linnaeus, 1758)	*S
HOMOLOBINAE		<b>Vipio</b> Latreille, 1804	
HOMOLOBINI		<i>desertor</i> (Linnaeus, 1758)	B M S
<b>Homolobus</b> Foerster, 1862		<b>Earinus</b> Wesmael, 1837	
<i>annulicornis</i> (Nees, 1834)	M	<i>elator</i> (Fabricius, 1804)	S
<i>discolor</i> (Wesmael, 1835)	M S	<i>gloriosus</i> (Panzer, 1809)	*S
<i>flagitator</i> (Curtis, 1837)	M	<b>Agathis</b> Latreille, 1804	
<i>infumator</i> (Lyle, 1914)	M S	<i>brevicaudis</i> (Reinhard, 1867)	*B
<i>truncator</i> (Say, 1828)	*S	<i>breviseta</i> Nees, 1814	S
CHARMONTINI		<i>calculator</i> (Fabricius, 1798)	B S
<b>Charmon</b> Haliday, 1833		<i>cinctulator</i> (Ratzeburg, 1852)	B
<i>cruentatus</i> Haliday, 1833	*B M S	<i>clausthaliana</i> (Ratzeburg, 1844)	S
<i>extensor</i> (Linnaeus, 1758)	B M S	<i>dimidiator</i> (Nees, 1834)	S
ORGILINAE			
<b>Orgilus</b> Haliday, 1833			

<i>fortipes</i> (Reinhard, 1867)	B	<i>lineipes</i> (Wesmael, 1837)	B M S
<i>lugubrator</i> (Ratzeburg, 1852)	B S	<i>longicauda</i> (Wesmael, 1837)	B M S
<i>malvacearum</i> Latreille, 1805	S	<i>longipalpalis</i> (Reinhard, 1880)	S
<i>minuta</i> Niezabitowski, 1910	*B	<i>murinanae</i> (Čapek et Zwölfer, 1957)	M S
<i>nigra</i> Nees, 1814	*S	<i>mycale</i> (Nixon, 1972)	S
<i>pumila</i> (Ratzeburg, 1844)	S	<i>obstans</i> (Papp, 1971)	*S
<i>rufipalpis</i> Nees, 1814	*S	<i>phaloniae</i> (Wilkinson, 1940)	*S
<i>rufipes</i> (Nees, 1814)	M S	<i>praetor</i> (Marshall, 1885)	*S
<i>tibialis</i> Nees, 1814	*B M S	<i>princeps</i> (Wilkinson, 1941)	*S
<i>tumidula</i> (Nees, 1814)	B S	<i>punctiger</i> (Wesmael, 1837)	S
<b>Baeognatha</b> Kokujev, 1903		<i>seriphia</i> (Nixon, 1972)	S
<i>nigra</i> Telenga, 1955	S	<i>sicaria</i> (Marshall, 1885)	S
MICROGASTRINAE		<i>sophiae</i> (Papp, 1972)	S
APANTELINI		<i>ultor</i> (Reinhard, 1880)	S
<b>Dolichogenidea</b> Viereck, 1911		<b>Pholetesor</b> Mason, 1981	
<i>breviventris</i> (Ratzeburg, 1848)	M S	<i>arisba</i> (Nixon, 1973)	*S
<i>coleophorae</i> (Wilkinson, 1938)	S	<i>circumscriptus</i> (Nees, 1834)	B M S
<i>credne</i> (Nixon, 1973)	S	<i>elpis</i> (Nixon, 1973)	S
<i>cytherea</i> (Nixon, 1972)	S	<i>exiguus</i> (Haliday, 1834)	S
<i>dilecta</i> (Haliday, 1834)	B M S	<i>maritimus</i> (Wilkinson, 1941)	S
<i>drusilla</i> (Nixon, 1972)	*S	<i>nanus</i> (Reinhard, 1880)	B M S
<i>emarginata</i> (Nees, 1834)	B M S	<i>pedias</i> (Nixon, 1973)	*B S
<i>ensiformis</i> (Ratzeburg, 1844)	S	<i>viminetorum</i> (Wesmael, 1837)	B S
<i>erasmi</i> (Nixon, 1972)	S	<b>Apanteles</b> Foerster, 1862	
<i>gracilariae</i> (Wilkinson, 1940)	B S	<i>ater</i> (Ratzeburg, 1852)	S
<i>iarbas</i> (Nixon, 1972)	B M	<i>brunnistigma</i>	
<i>imperator</i> (Wilkinson, 1939)	B	Abdinbekova, 1969	*S
<i>impurus</i> (Nees, 1834)	B	<i>lenea</i> Nixon, 1976	S
<i>infima</i> (Haliday, 1834)	*S	<i>metacarpalis</i> Thomson, 1895	*S
<i>lactea</i> (Nees, 1834)	B S	<i>obscurus</i> (Nees, 1834)	S
<i>lacteicolor</i> (Viereck, 1911)	B S	<i>weitenweberi</i> (Amerling, 1862)	?
<i>lacteipennis</i> (Curtis, 1830)	B M S	<i>xanthostigma</i> (Haliday, 1834)	M S
<i>laevigata</i> (Ratzeburg, 1848)	B M S	<b>Illidops</b> Mason, 1981	
<i>laevigatooides</i> (Nixon, 1972)	B	<i>butalidis</i> (Marshall, 1885)	S
<i>laevissima</i> (Ratzeburg, 1848)	*M	<i>cloelia</i> (Nixon, 1965)	S
<i>temariei</i> (Nixon, 1961)	M S	<i>naso</i> (Marshall, 1885)	*S

<i>suevus</i> (Reinhard, 1880)	M	S	<i>compressiventris</i>	
<i>szaboi</i> (Papp, 1972)		*S	[Muesebeck, 1921]	*S
<i>urgo</i> (Nixon, 1965)		*S	<i>eugeni</i> (Papp, 1972)	S
LISSOGASTRINI			<i>fraternus</i> (Reinhard, 1881)	S
<b>Iconella</b> Mason, 1981			<i>fulvipes</i> (Haliday, 1834)	S
<i>lacteoides</i> (Nixon, 1965)	S		<i>inclusus</i> (Ratzeburg, 1844)	S
<i>merula</i> (Reinhard, 1880)		*S	<i>lateralis</i> (Haliday, 1834)	S
<b>Choeras</b> Mason, 1981			<i>liparidis</i> (Bouché, 1834)	B S
<i>dorsalis</i> (Spinola, 1808)		*S	<i>mygdonia</i> (Papp, 1973)	S
<i>parasitellae</i> (Bouché, 1834)		S	<i>octonarius</i> (Ratzeburg, 1852)	S
<i>ruficornis</i> (Nees, 1834)		*S	<i>pallipes</i> (Reinhard, 1880)	S
<i>tedellae</i> (Nixon, 1961)	B	M	<i>pinicola</i> (Lyle, 1917)	S
<i>tiro</i> (Reinhard, 1880)		*S	<i>porthetriae</i> (Muesebeck, 1928)	S
<i>validus</i> (Thomson, 1895)		S	<i>ripus</i> (Papp, 1983)	S
<b>Sathon</b> Mason, 1981			<i>vitripennis</i> (Curtis, 1830)	S
<i>falcatus</i> (Nees, 1834)		*S	<b>Protapanteles</b> Ashmead, 1898	
<b>Lissogaster</b> Bengtsson, 1926			<i>anchisiades</i> (Nixon, 1973)	*S
<i>consors</i> (Nixon, 1968)		*S	<i>andromica</i> (Nixon, 1976)	S
<i>famulus</i> (Nixon, 1968)		*S	<i>caberae</i> (Marshall, 1885)	S
<i>fulvicrus</i> (Thomson, 1895)		*S	<i>delitutus</i> (Papp, 1984)	S
<i>hospes</i> (Marshall, 1885)	M	S	<i>enephes</i> (Nixon, 1965)	S
<i>laeviscuta</i> (Thomson, 1895)	B	M	<i>immunis</i> (Haliday, 1834)	B S
<i>parvistriga</i> (Thomson, 1895)		S	<i>popularis</i> (Haliday, 1834)	*S
<i>sticticus</i> (Ruthe, 1858)		S	<i>triangulator</i> (Wesmael, 1837)	S
<i>subcompleta</i> (Nees, 1834)	B	S	<b>Cotesia</b> Cameron, 1891	
COTESIINI			<i>abjecta</i> (Marshall, 1885)	S
<b>Rasivalva</b> Mason, 1981			<i>acuminata</i> (Reinhard, 1880)	
<i>calceata</i> (Haliday, 1834)	S		<i>affinis</i> (Nees, 1834)	M S
<b>Distatrix</b> Mason, 1981			<i>amesis</i> (Nixon, 1974)	*S
<i>pompelon</i> (Nixon, 1965)	S		<i>ancilla</i> (Nixon, 1974)	B S
<b>Venanides</b> Mason, 1981			<i>arctica</i> (Thomson, 1895)	*S
<i>moldavicus</i> (Tobias, 1975)	S		<i>brevicornis</i> (Wesmael, 1837)	S
<b>Glyptapanteles</b> Ashmead, 1905			<i>cajae</i> (Bouché, 1834)	M S
<i>acasta</i> (Nixon, 1973)	*S		<i>callimone</i> (Nixon, 1974)	S
<i>aletta</i> (Nixon, 1973)	S		<i>chares</i> (Nixon, 1965)	*S
<i>aliphera</i> (Nixon, 1973)	S		<i>cleora</i> (Nixon, 1974)	B
<i>callidus</i> (Haliday, 1834)	S		<i>cupreus</i> (Lyle, 1925)	*S
<i>capeki</i> (Györfi, 1955)	S		<i>ferruginea</i> (Marshall, 1885)	*B
			<i>gastropachae</i> (Bouché, 1834)	B M S
			<i>geryonis</i> (Marshall, 1885)	*S

<i>glomerata</i> (Linnaeus, 1758)	B M S		
<i>hyphantriae</i> (Riley, 1887) introd.	S		
<i>isolde</i> (Nixon, 1974)	S		
<i>jucunda</i> (Marshall, 1885)	S		
<i>juniperatae</i> (Bouché, 1834)	S		
<i>limbata</i> (Marshall, 1885)	S		
<i>melanoscelus</i> (Ratzeburg, 1844)	B M S		
<i>melitaearum</i> (Wilkinson, 1937)	*S		
<i>memnon</i> (Nixon, 1974)	*S		
<i>nothus</i> (Marshall, 1885)	*S		
<i>ofella</i> (Nixon, 1974)	*S		
<i>onaspis</i> (Nixon, 1974)	*S		
<i>ordinaria</i> (Ratzeburg, 1844)	*S		
<i>pieridis</i> (Bouché, 1834)	S		
<i>pilicornis</i> (Thomson, 1895)	S		
<i>plutellae</i> (Kurdjumov, 1912)	S		
<i>praepotens</i> (Haliday, 1834)	S		
<i>risilis</i> (Nixon, 1974)	S		
<i>rubecula</i> (Marshall, 1885)	B		
<i>ruficrus</i> (Haliday, 1834)	S		
<i>saltator</i> (Thunberg, 1822)	S		
<i>scabriculus</i> (Reinhard, 1880)	S		
<i>sibyllarum</i> (Wilkinson, 1936)	B		
<i>spurius</i> (Wesmael, 1837)	M S		
<i>tetrica</i> (Reinhard, 1880)	M S		
<i>tibialis</i> (Curtis, 1830)	B M S		
<i>villana</i> (Reinhard, 1880)	S		
<i>zygaenarum</i> (Marshall, 1885)	B S		
<b>Diolcogaster</b> Ashmead, 1890			
<i>abdominalis</i> (Nees, 1834)	S		
<i>alvearia</i> (Fabricius, 1798)	S		
<i>spretus</i> (Marshall, 1885)	S		
<b>Deuterixys</b> Mason, 1981			
<i>carbonaria</i> (Wesmael, 1837)	S		
<i>rimulosa</i> (Niezabitowski, 1910)	S		
			<b>Microgaster</b> Latreille, 1804
			<i>capeki</i> (Nixon, 1970) M
			<i>cebes</i> (Nixon, 1970) S
			<i>deprimator</i> (Fabricius, 1798) B M S
			<i>flavipalpis</i> Brullé, 1832 S
			<i>heterocera</i> Ruthe, 1860 S
			<i>impressa</i> Wesmael, 1837 S
			<i>mandibularis</i> Thomson, 1895 B S
			<i>mediator</i> Haliday, 1834 M S
			<i>naenia</i> (Nixon, 1970) S
			<i>ocellatae</i> Bouché, 1834 M
			<i>spectabilis</i> Haliday, 1834 S
			<i>spinolae</i> Nees, 1834 S
			<i>tristis</i> Nees, 1834 B S
			<i>tuberculifer</i> Wesmael, 1837 M S
			<i>varipes</i> Ruthe, 1860 S
			CARDIOCHILINAE
			CARDIOCHILINI
			<b>Cardiochiles</b> Nees, 1818
			<i>saltator</i> (Fabricius, 1781) M S
			MIRACINI
			<b>Mirax</b> Haliday, 1833
			<i>dryochares</i> Marshall, 1898 ?
			<i>rufilabris</i> Haliday, 1833 B S
			ADELIINI
			<b>Adelius</b> Haliday, 1833
			<i>clandestinus</i> Foerster, 1851 B
			<i>determinatus</i> Foerster, 1851 S
			<i>dubius</i> Foerster, 1851 B
			<i>erythronotus</i> Foerster, 1851 B
			<i>hyalinipennis</i> Foerster, 1851 B
			<i>parvulus</i> Foerster, 1851 B
			<i>subfasciatus</i> Haliday, 1833 B M S
			<i>viator</i> Foerster, 1851 B S
			CHELONINAE
			SIGALPHINI
			<b>Acampsis</b> Wesmael, 1835

<i>alternipes</i> (Nees, 1816)	M S	<i>hungaricus</i> (Szépligeti, 1908)	*S
<b>Sigalphus</b> Latreille, 1802		<i>latrunculus</i> (Marshall, 1885)	M*S
<i>irrorator</i> (Fabricius, 1775)	S	<i>microphthalmus</i> (Wesmael, 1838)	*S
<b>CHELONINAE</b>		<i>nitens</i> (Reinhard, 1867)	*S
<b>Chelonus</b> Jurine, 1801		<i>paricornis</i> (Herrich et Schaeffer, 1838)	M
<i>annulatus</i> (Nees, 1816)	S	<i>pictipes</i> (Wesmael, 1838)	M S
<i>annulipes</i> Wesmael, 1835	S	<i>retusus</i> (Nees, 1816)	M S
<i>asiaticus</i> Telenga, 1941	S	<i>rimatus</i> (Szépligeti, 1896)	M S
<i>bidens</i> Tobias, 1972	S	<i>risorius</i> (Reinhard, 1867)	M
<i>bimaculatus</i> Szépligeti, 1896	S	<i>rugicollis</i> (Thomson, 1874)	S
<i>brachyurus</i> Thomson, 1874	S	<i>scabrosus</i> Szépligeti, 1896	S
<i>canescens</i> Wesmael, 1835	M S	<i>starki</i> Telenga, 1953	S
<i>carbonator</i> Marshall, 1885	M	<i>sulcatus</i> (Jurine, 1807)	M
<i>corvulus</i> Marshall, 1885	M S	<b>Ascogaster</b> Wesmael, 1835	
<i>dauricus</i> Telenga, 1941	S	<i>abdominator</i> (Dahlbom, 1833)	*B S
<i>foveolatus</i> Niezabitowski, 1910	M	<i>albitarsus</i> Reinhard, 1867	B *S
<i>gravenhorsti</i> (Nees, 1816)	S	<i>annularis</i> (Nees, 1816)	S
<i>hiemalis</i> (Gautier et Cleu, 1930)	*S	<i>armata</i> Wesmael, 1835	B S
<i>inanitus</i> (Linnaeus, 1767)	M S	<i>bidentula</i> Wesmael, 1835	B S
<i>intermedius</i> Thomson, 1874	M	<i>canifrons</i> Wesmael, 1835	B S
<i>obscuratus</i> Herrich et Schaeffer, 1838	M S	<i>caucasica</i> Kokuvjev, 1895	*S
<i>oculator</i> (Panzer, 1779)	M	<i>consobrina</i> Curtis, 1837	*M S
<i>pannonicus</i> Szépligeti, 1896	M S	<i>contracta</i> Ratzeburg, 1848	S
<i>pulchricornis</i> Szépligeti, 1898	S	<i>dentifer</i> Tobias, 1976	*S
<i>scabrator</i> (Fabricius, 1793)	M S	<i>dispar</i> Fahringer, 1934	*B S
<i>sculpturatus</i> Szépligeti, 1898	*S	<i>klugii</i> (Nees, 1816)	B *S
<i>seticornis</i> Thomson, 1891	*S	<i>quadridentata</i> Wesmael, 1835	B M S
<i>submuticus</i> Wesmael, 1835	*S	<i>reticulata</i> Watanabe, 1967	*S
<i>subseticornis</i> Tobias, 1972	*S	<i>rufidens</i> Wesmael, 1835	B M S
<i>zimini</i> Tobias, 1972	*S	<i>rufipes</i> (Latreille, 1809)	B S
<b>Microchelonus</b> Szépligeti, 1908		<i>scabricula</i> (Dahlbom, 1833)	*S
<i>alboannulatus</i> Szépligeti, 1896	S	<i>similis</i> (Nees, 1816)	S
<i>cingulipes</i> Niezabitowski, 1910	B M	<i>varipes</i> Wesmael, 1835	B M S
<i>contractus</i> (Nees, 1816)	M S	<b>Phanerotoma</b> Wesmael, 1838	
<i>exilis</i> (Marshall, 1885)	*S	<i>acuminata</i> Szépligeti, 1908	M
<i>fenestratus</i> (Nees, 1816)	M	<i>antennalis</i> Šnoflák, 1950	B M S
		<i>atra</i> Šnoflák, 1950	M S
		<i>dentata</i> (Panzer, 1805)	S
		<i>gregori</i> Šnoflák, 1950	M
		<i>minor</i> Šnoflák, 1950	M S

<i>moravica</i> Šnoflák, 1950	M	S	<i>pulchricornis</i> (Wesmael, 1835)		S
<i>obscura</i> Šnoflák, 1950	M		<i>rubens</i> (Nees, 1811)		M S
<i>picta</i> Šnoflák, 1950	M		<i>rubriceps</i> (Ratzeburg, 1844)	B	S
<i>planifrons</i> (Nees, 1816)	M	S	<i>ruficeps</i> (Nees, 1811)		M S
<i>platypyga</i> Šnoflák, 1950	M		<i>salicorniae</i>		
<b>Phanerotomella</b> Szépligeti, 1900			Schmiedeknecht, 1897	*B	S
<i>flavipes</i> Šnoflák, 1950	M		<i>sulcatus</i> Szépligeti, 1896		S
<i>nigra</i> Szépligeti, 1900		?	<i>tabidus</i> (Wesmael, 1835)	B	
ICHNEUTINAE			<i>unicolor</i> (Wesmael, 1835)	B	S
<b>Proterops</b> Wesmael, 1835			<i>versicolor</i> (Wesmael, 1835)	B	S
<i>nigripennis</i> Wesmael, 1835	M	S	<i>vexator</i> (Haliday, 1835)		*S
<b>Ichneutes</b> Nees, 1816			NEONEURINAE		
<i>brevis</i> Wesmael, 1835	B		<b>Neoneurus</b> Haliday, 1838		
<i>reunitor</i> Nees, 1816	B	M S	<i>auctus</i> (Thomson, 1895)	M*S	
METEORINAE			<i>viennensis</i> (Giraud, 1870)	M S	
<b>Zele</b> Curtis, 1832	B	S	<b>Elasmosoma</b> Ruthe, 1858		
<i>albiditarsus</i> Curtis, 1832	B	S	<i>berolinense</i> Ruthe, 1858	B	
<i>chlorophthalmus</i> (Spinola, 1808)	B	M S	EUPHORINAE		
<b>Meteorus</b> Haliday, 1835			<b>Allurus</b> Foerster, 1862		
<i>abdominator</i> (Nees, 1811)		*S	<i>muricatus</i> (Haliday, 1833)	*S	
<i>abscissus</i> Thomson, 1895		S	<b>Centistes</b> Haliday, 1835		
<i>affinis</i> (Wesmael, 1835)		S	<i>cuspidatus</i> (Haliday, 1833)	?	
<i>cinctellus</i> (Spinola, 1808)	*M	S	<i>edentatus</i> (Haliday, 1835)	*S	
<i>colon</i> Haliday, 1835		S	<i>fuscipes</i> (Nees, 1834)	?	
<i>eadyi</i> Huddleston, 1980		*S	<b>Pygostolus</b> Haliday, 1833		
<i>filator</i> (Haliday, 1835)		*S	<i>falcus</i> (Nees ab Esenbeck, 1834)	*S	
<i>gyrator</i> (Thunberg, 1822)	B	S	<i>multiarticulatus</i> (Ratzeburg, 1852)	B	
<i>heliophilus</i> Fischer, 1970		*S	<i>sticticus</i> (Fabricius, 1798)	M	
<i>ictericus</i> (Nees, 1811)	B	M S	<b>Cosmophorus</b> Ratzeburg, 1848		
<i>jaculator</i> Haliday, 1835		S	<i>cembrae</i> Ruschka, 1925	B	S
<i>lionotus</i> Thomson, 1895		S	<i>klugii</i> Ratzeburg, 1848	B	S
<i>longicornis</i> (Ratzeburg, 1844)		S	<i>regius</i> Niezabitowski, 1910	B	S
<i>micropterus</i> Haliday, 1835		*S	<i>roubali</i> Čapek, 1958	B	
<i>obfuscatus</i> (Nees, 1811)	B	S	<b>Aridelus</b> Marshall, 1887		
<i>obsoletus</i> (Wesmael, 1835)		*B	<i>egregius</i> (Schmiedeknecht, 1907)	B	S
<i>oculatus</i> Ruthe, 1862		*S			
<i>pallidus</i> (Nees, 1811)		S			
<i>pallipes</i> (Wesmael, 1835)		S			
<i>pendulator</i> (Latreille, 1799)		?			

<b>Wesmaelia</b> Foerster, 1862				
<i>pendula</i> Foerster, 1862		M		
<b>Perilitus</b> Nees, 1818				
<i>areolaris</i> Gerdin et Hedqvist, 1984	*B			
<i>dubius</i> (Wesmael, 1838)		*S		
<i>falciger</i> (Ruthe, 1856)		*S		
<i>pallidipes</i> (Costa, 1883)		*S		
<i>rutilus</i> (Nees, 1811)	B			
<i>strenuus</i> Marshall, 1887		S*		
<b>Peristenus</b> Foerster, 1862				
<i>orchesiae</i> (Curtis, 1833)		*S		
<i>pallipes</i> (Curtis, 1833)		*S		
<i>picipes</i> (Curtis, 1833)		*S		
<b>Dinocampus</b> Foerster, 1862				
<i>coccinellae</i> (Schrank, 1802)	B	S		
<b>Microctonus</b> Wesmael, 1835				
<i>aethiops</i> (Nees, 1834)		*S		
<i>brevicollis</i> (Haliday, 1835)		*S		
<i>labilis</i> Ruthe, 1856		*S		
<i>melanopus</i> Ruthe, 1856		*S		
<i>secalis</i> (Haliday, 1833)		S		
<b>Townesilitus</b> Haeselbarth et Loan, 1983				
<i>bicolor</i> (Wesmael, 1835)	*B			
<b>Chrysopophthorus</b> Goidanich, 1948				
			<b>hungaricus</b> (Zilahi-Kiss, 1927]	M
			<b>Syntretus</b> Foerster, 1862	
			<i>conterminus</i> (Nees, 1834)	*S
			<i>elegans</i> (Ruthe, 1856)	*M S
			<i>testaceus</i> (Capron, 1887)	*S
			<i>vernalis</i> (Wesmael, 1835)	*S
			<b>Loxocephalus</b> Foerster, 1862	
			<i>boops</i> (Wesmael, 1835)	?
			<b>Leiophron</b> Nees, 1818	
			<i>apicalis</i> Curtis, 1833	*S
			<i>fascipennis</i> (Ruthe, 1856)	*S
			<i>similis</i> Curtis, 1833	*S
			<b>Streblocera</b> Westwood, 1833	
			<i>longiscapha</i> Westwood, 1882	M
			<i>macroscapa</i> (Ruthe, 1856)	M S
			<b>Ropalophorus</b> Curtis, 1837	
			<i>clavicornis</i> (Wesmael, 1835)	B S
			PAXYLOMMATINAE	
			<b>Hybrizon</b> Fallen, 1813	
			<b>Paxylomma</b> , de Brébisson, 1825	
			<i>buccatus</i> de Brébison, 1825	B M S

## APHIDIIDAE

PETR STARÝ

The history of research on this group in Czechoslovakia is very brief. In fact, it started in the nineteen sixties when several young hymenopterists became specialised in the particular group. The first summarising paper was published by Starý (1966).

The collections deposited in various museums and institutions are generally poor and involve prevalently netted material which is of low value as for the mounting requirements and identification procedures. The greatest collection is temporarily deposited in the Institute of Entomology, Czechoslovak Academy of Sciences, in České Budějovice.

Most of the references can be found in the comprehensive paper by Mackauer et Starý (1967). Almost all information on the group was obtained from the collection of Aphidiidae deposited in the Institute of Entomology (see above).

The aphidiids are solitary endophagous parasitoids of aphids.

The territory of Czechoslovakia is among the areas where information on the aphidiid fauna is very good. The species composition seems to be prevalently determined but more information on the host range of the species is required.

The nomenclature used by Mackauer et Starý (1967, see above) has largely been adopted.

References: Mackauer M. et Starý P. (in Delucchi V. et Remaudière G. Eds.), 1967, Hym. Ichneumonoidea World Aphidiidae, Index of entomophagous insects, Le François, Paris: 197 p. Starý P., 1966: Aphid parasites of Czechoslovakia. Academia Praha et Dr. W. Junk N. V. The Hague: 242 p.

### APHIDIIDAE

<b>Aclitus</b> Foerster, 1862		<b>Aphidius</b> Nees, 1819
<i>obscuripennis</i> Foerster, 1862	B	<i>absinthii</i> Marshall, 1896 B M S
<b>Adialytus</b> Foerster, 1862		<i>aquilus</i> Mackauer, 1961 B S
<i>ambiguus</i> (Haliday, 1834)	B M S	<i>cingulatus</i> Ruthe, 1859 B M S
<i>salicaphis</i> (Fitch, 1855)	B M S	<i>colemani</i> Viereck, 1912
<i>thelaxis</i> (Starý, 1961)	B S	introd.,? B M
		<i>transcaspicus</i> Telenga, 1958
		<i>eadyi</i> Starý et Gonzalez, Hall, 1980 B M S

<i>eglantiae</i> Haliday, 1834	B	<i>planiceps</i> (Marshall, 1896)	B M S
<i>ervi</i> Haliday, 1834	B M S	<b>Ephedrus</b> Haliday, 1833	
<i>funebris</i> Mackauer, 1961	B M S	<i>blatnyi</i> Starý, 1973	S
<i>hieracorum</i> Starý, 1962	B M S	<i>brevis</i> Stelfox, 1941	B
<i>hortensis</i> Marshall, 1896	B M S	<i>cerasicola</i> Starý, 1962	B S
<i>matricariae</i> Haliday, 1834	B M S	<i>lacertosus</i> (Haliday, 1833)	B M S
<i>megourae</i> Starý, 1965	B	<i>minor</i> Stelfox, 1941	B M S
<i>phalangomyzi</i> Starý, 1963	B S	<i>nacheri</i> Quilis, 1934	B M
<i>picipes</i> (Nees, 1811)	B M S	<i>niger</i> Gautier et Bonnamour, Gaumont, 1929	
<i>rhopalosippi</i> DeStefani, 1902	B M S	<i>persicae</i> Froggatt, 1904	B M S
<i>ribis</i> Haliday, 1834	B M S	<i>plagiator</i> (Nees, 1811)	B M S
<i>rosae</i> Haliday, 1834	B M S	<i>prociphili</i> Starý, 1982	B M S
<i>salicis</i> Haliday, 1834	B M S	<i>salicicola</i> Takada, 1968	B
<i>schimitscheki</i> (Starý, 1960)	B	<b>Falciconus</b> Mackauer, 1959	
<i>setiger</i> Mackauer, 1961	B M	<i>pseudoplatani</i> (Marshall, 1896)	B M S
<i>smithi</i> Sharma et Subba Rao, 1959		<b>Lipolexis</b> Foerster, 1862	
introd.,?	B M S	<i>gracilis</i> Foerster, 1862	B M S
<i>sonchi</i> Marshall, 1896	B	<b>Lysaphidus</b> Smith, 1944	
<i>tanacetarius</i> Mackauer, 1962	B M S	<i>arvensis</i> Starý, 1960	B M S
<i>urticae</i> Haliday, 1834	B M S	<i>erysimi</i> Starý, 1960	B S
<i>uzbekistanicus</i> Luzhetszki, 1960	B M S	<b>Lysephedrus</b> Starý, 1958	
<b>Areopraon</b> Mackauer, 1959		<i>validus</i> (Haliday, 1833)	B M
<i>lepelleyi</i> (Waterston, 1926)	B	<b>Lysiphlebus</b> Foerster, 1862	
<b>Betuloxys</b> Mackauer, 1960		<i>Lysiphlebus</i> s. str.	
<i>compressicornis</i> (Ruthe, 1859)	B	<i>dissolutus</i> (Nees, 1811)	B
<i>hortorum</i> (Starý, 1960)	B M	<i>Phlebus</i> Starý, 1975	
<b>Calaphidius</b> Mackauer, 1961		<i>cardui</i> (Marshall, 1896)	B M S
<i>elegans</i> Mackauer, 1961	B	<i>confusus</i> Tremblay et Eady, 1978	B M S
<b>Diaeretellus</b> Starý, 1960		<i>fabarum</i> (Marshall, 1896)	B M S
<i>ephippium</i> (Haliday, 1833)	B	<i>fritzmuelleri</i> Mackauer, 1960	B S
<i>heinzei</i> (Mackauer, 1959)	B	<i>hirticornis</i> Mackauer, 1960	B S
<i>macrocarpus</i> Mackauer, 1961	B	<i>melandriicola</i> Starý, 1961	B M
<b>Diaeretiella</b> Starý, 1960		<i>testaceipes</i> (Cresson, 1880) introd.,?	B
<i>rapae</i> (M'Intosh, 1855)	B M S		
<b>Diaeretus</b> Foerster, 1862			
<i>leucopterus</i> (Haliday, 1834)	B M S		
<b>Dyscritulus</b> Hincks, 1943			

<b>Metaphidius</b> Starý et Sedlag, 1959		<i>exsoletum</i> (Nees, 1811)	B	M	S
<i>aterrimus</i> (Fahringer, 1935)	B	<i>flavinode</i> (Haliday, 1833)	B	M	S
<b>Monoctonia</b> Starý, 1962		<i>grossum</i> Starý, 1971	B	M	S
<i>pistaciaecola</i> Starý, 1962	B	<i>lemantinum</i> Gautier, 1922	B	M	
<b>Monoctonus</b> Haliday, 1833		<i>megourae</i> Starý, 1971	M		
<i>caricis</i> (Haliday, 1833)	B	<i>pubescens</i> Starý, 1961	B	M	S
<i>cerasi</i> (Marshall, 1896)	B	<i>rosaecola</i> Starý, 1961	B		
<i>crepidis</i> (Haliday, 1834)	B	<i>silvestre</i> Starý, 1971	B	M	
<i>nervosus</i> (Haliday, 1833)	B	<i>volucre</i> (Haliday, 1833)	B	M	S
<b>Paralipsis</b> Foerster, 1862		<b>Protaphidius</b> Ashmead, 1900			
<i>enervis</i> (Nees, 1834)	B	<i>wissimanni</i> (Ratzeburg, 1848)	B	S	
<b>Paramonoctonus</b> Starý, 1959		<b>Pseudopraon</b> Starý, 1975			
<i>angustivalvus</i> (Starý, 1959)	B	<i>mindariphagum</i> Starý, 1975	B	M	S
<b>Parapraon</b> Starý, 1982		<b>Toxares</b> Haliday, 1833			
<i>gallicum</i> (Starý, 1971)	B	<i>deltiger</i> (Haliday, 1833)	B	S	
<i>necans</i> (Mackauer, 1959)	B	<b>Trioxys</b> Haliday, 1833			
<b>Pauesia</b> Quilis, 1931		<b>Binodoxys</b> Mackauer 1960			
<b>Paraphidius</b> Starý, 1958		<i>acalephae</i> (Marshall, 1896)	B	M	S
<i>abietis</i> (Marshall, 1896)	B	<i>angelicae</i> (Haliday, 1833)	B	M	S
<i>cupressobii</i> (Starý, 1960)	M	<i>brevicornis</i> (Haliday, 1833)	B	M	S
<i>grossa</i> (Fahringer, 1937)	B	<i>centaureae</i> (Haliday, 1833)	B	M	S
<i>infulata</i> (Haliday, 1834)	B	<i>genistae</i> Mackauer, 1960	S		
<i>jezoensis</i> Watanabe, 1941	B	<i>heraclei</i> (Haliday, 1833)	B	M	
<i>juniperorum</i> (Starý, 1960)	B	<b>Trioxys</b> s. str.			
<i>maculolachni</i> (Starý, 1960)	B	<i>auctus</i> (Haliday, 1833)	B		
<i>pini</i> (Haliday, 1834)	B	<i>betulae</i> (Marshall, 1896)	B	M	
<i>silvestris</i> (Starý, 1960)	B	<i>cirsii</i> (Curtis, 1831)	B	M	S
<b>Pauesia</b> s. str.		<i>complanatus</i> Quilis, 1931	B		
<i>laricis</i> (Haliday, 1834)	B	<i>curvicaudus</i> Mackauer, 1967	B	M	S
<i>picta</i> (Haliday, 1834)	S	<i>falcatus</i> Mackauer, 1959	B	M	
<i>unilachni</i> (Gahan, 1927)	B	<i>galiobii</i> Starý, 1974	M		
<b>Praon</b> Haliday, 1833		<i>glaber</i> Starý, 1967	S		
<i>abjectum</i> (Haliday, 1833)	B	<i>humuli</i> Mackauer, 1960	B		
<i>absinthii</i> Bignell, 1894	B	<i>macroceratus</i> Mackauer, 1960	M		
<i>barbatum</i> Mackauer, 1967	B	<i>pallidus</i> (Haliday, 1834)	B	M	S
<i>bicolor</i> Mackauer, 1959	B	<i>pannonicus</i> Starý, 1960	B	M	S
<i>cavariellae</i> Starý, 1971	B	<i>parauctus</i> Starý, 1960	B	M	S
<i>dorsale</i> (Haliday, 1833)	B				

*phyllaphidis* Mackauer,  
1961

M

*spinosus* Starý, 1963 S  
*tenuicaudus* Starý, 1972 B M

## ICHNEUMONIDAE

JOSEF ŠEDIVÝ

The family Ichneumonidae is the richest group of Hymenoptera as to species. In Czechoslovakia, the species of this family account for 20 % of the total of parasitic Hymenoptera.

The first literary data on the Ichneumonidae in our territory are found in the work Kirchner (1867).

Systematic investigations into the fauna of the family Ichneumonidae in our territory were initiated by Gregor, sen. (1928, 1929). His collection of Ichneumonidae is the greatest in Czechoslovakia and is deposited in the National Museum in Prague and partly in the Moravian Museum in Brno. In 1938—1962 Kolubajiv (1962) studied the importance of ichneumonids as parasitoids of forest insect pests. His collection is deposited in the Research Institute for Forest Management and Gamekeeping in Prague-Jíloviště. In 1946—1960, R. Obrtel published papers on ichneumonids, mainly of subfamilies Pimplinae and Phygadeuontinae. His collection is deposited in the Moravian Museum in Brno. The present author evaluated faunal findings of Pimplinae, Xoridinae, Acaenitinae and Anomalinae. He also published a revision of the European species of the tribe Polysphinctini (Šedivý, 1963) and of the subfamily Cremastinae (Šedivý, 1970, 1971). The collection of the present author is private.

Most species of the genera *Triclistus*, *Trieces* and *Chorinaeus*, listed below, were determined by J. Aeschlimann, Montpellier, France. The species of the genus *Delopia* were determined by R. Hinz, Goslar, FRG, and the species of the genus *Diadegma*, subfamily Tersilochinae, part of those of other campoplegine genera were determined by K. Horstmann, Würzburg, FRG.

A major part of the collections of Ichneumonidae in Czechoslovakia has not been studied in detail as yet, and one may assume that the number of species may be at least  $\frac{1}{4}$  larger than that listed below.

The Ichneumonidae are parasitoids of eggs, larvae, pupae and rarely also adults of insects. They also parasitize eggs, cocoons of spiders (Aranei). Larvae of *Polysphincta* are external parasitoids of spiders. Larvae of *Obisiphaga stenoptera* develop in clutches of eggs of *Obisium*, *Pseudoscorpiones*. Larvae of some species develop as hyperparasitoids.

The check-list has been arranged according to the classifications used in the monographs 1962—1983 and the catalogue by H. Townes et al. (1965), in Check-list of British Insects by Kloet G. S. et al. (1978) and in the handbook edited by Kasparjan (1981).

References: Gregor, F., 1928: Příspěvek k poznání československých lomků (Ichneumonidae). *Sbor. klubu přírodnověd. v Brně*, 10: 1–11. — Gregor, F., 1929 a 1930: Tryphonidi Československé republiky I. a II. Výroční zpráva čes. stát. reál. gym. v Novém Jičíně: 1–11, 13–23. — Kasparjan D. R. et al., 1981: Opredelitel nasekomych evropskoy časti SSSR, T. 3. Perepončatokrylye III, Leningrad, 687 pp. — Kirchner J., 1867: Catalogus Hymenopterorum Europae, Vindobonae, 283 pp. — Kloet G. S. et Hincks, W. D., 1978: A Check List of British Insects. Handb. for Identi. Brit. Insects XI, Part 4, London: 12–46. — Kolubajiv S., 1962: Výsledky chovu entomofágů (cizopasníků a dravců) hmyzích škůdců — hlavně lesních — získané v období 1934–1958. *Rozpravy ČSAV*, 72: 73 pp. — Šedivý J., 1963: Die europäischen Arten der Gattungen Laufela Tosq., Polysphincta Grav. und Zatypota Först. (Hym. Ichneumonidae). *Acta ent. Mus. nat. Pragae* 35: 243–261. — Šedivý, J., 1970: Westpalaearktische Arten der Gattungen Dimorphora, Pristomerus, Eucremastus und Cremastus (Hym. Ichneumonidae). *Acta Sc. Nat. Brno* 4 (11): 38 pp. — Šedivý, J., 1971: Revision der europäischen Temelucha-Arten (Hym. Ichneumonidae). *Acta Sc. Nat. Brno* 5 (1): 34 pp. — Townes H., Momoi S., Townes M., 1965: A Catalogue and Reclassification of the Eastern Palearctic Ichneumonidae. *Mem. Amer. Ent. Soc.* Nr. 5: 661 pp.

#### ICHNEUMONIDAE

##### PIMPLINAE

##### EPHIALTINAE

##### EPHIALTINI

#### **Exeristes** Foerster, 1868

<i>Eremochila</i> Foerster 1868		
<i>arundinis</i> (Kriechbaumer, 1887)	B M S	
<i>longiseta</i> (Ratzeburg, 1844)	*B	
<i>robator</i> (Fabricius, 1793)	B M S	
<i>ruficollis</i> (Gravenhorst, 1829)	*B M S	

#### **Flavopimpla** Betrem, 1932

<i>Afrephialtes</i> Benoit, 1932		
<i>cicatricosa</i> (Ratzeburg, 1848)	M	

#### **Scambus** Hartig, 1838

<i>Endromopoda</i> Hellén, 1939		
<i>arundinator</i> (Fabricius, 1804)	B M S	
<i>melanopyga</i> (Gravenhorst 1829)		
<i>detritus</i> (Holmgren, 1860)	B M S	
<i>brunneus</i> (Brischke, 1880)		
<i>dilutus</i> (Ratzeburg, 1852)	*B	*S
<i>phragmitidis</i> Perkins, 1957	B M S	
<i>Atteleophadnus</i> Cameron, 1905		
<i>nigricans</i> (Thomson, 1877)	B M S	

#### *rufator* Aubert, 1963

S

<i>Scambus</i> s. str.		
<i>Epiurus</i> Foerster, 1868		
<i>annulatus</i> (Kiss, 1924)	B	S
<i>brevicornis</i> (Gravenhorst, 1829)	B M S	
<i>buolianae</i> (Hartig, 1838)	B	S
<i>calobatus</i> (Gravenhorst, 1829)	B M S	
<i>elegans</i> (Woldstedt, 1876)	M S	
<i>eucosmidarum</i> (Perkins, 1957)	S	
<i>planatus</i> (Hartig, 1838)	B M S	
<i>nucum</i> (Ratzeburg, 1844)		
<i>pomorum</i> (Ratzeburg, 1848)	B M S	
<i>sagax</i> Hartig, 1838	B M S	
<i>signatus</i> (Pfeffer, 1913)	*B	
<i>strobilarum</i> (Ratzeburg, 1848)	B M S	
<i>vesicarius</i> (Ratzeburg, 1844)	B M S	
<b>Epiphialtes</b> Gravenhorst, 1829		
<i>Pimpla</i> sensu Curtis, 1828		
<i>manifestator</i> (Linnaeus, 1758)	B M S	
<i>polytauma</i> Heinrich, 1949	*B	*S
<b>Liotryphon</b> Ashmead, 1900		
<i>caudatus</i> Ratzeburg, 1848	B M S	
<i>crassisetus</i> (Thomson, 1877)	*B	
<i>cydiae</i> (Perkins, 1942)		*S

<i>punctulatus</i> (Ratzeburg, 1848)	B M S	<i>pictipes</i> (Gravenhorst, 1829)	M S
<i>ruficollis</i> (Desvignes, 1856)	B M S	<b>Gregopimpla</b> Momoi, 1965	
<i>strobilellae</i> (Linnaeus, 1758)	B M S	<i>bernuthii</i> (Hartig, 1838)	B M S
<b>Townesia</b> Ozols, 1962		<i>inquisitor</i> (Scopoli, 1763)	B M S
<i>tenuiventris</i> (Holmgren, 1860)	B M S	<b>Iseropus</b> Foerster, 1868	
<b>Paraperithous</b> Haupt, 1954		<i>stercorator</i> (Fabricius, 1793)	B M S
<i>gnathaulax</i> (Thomson, 1877)	B M S	<b>Tromatobia</b> Foerster, 1868	
<b>Dolichomitus</b> Smith, 1877		<i>oculatoria</i> (Fabricius, 1798)	B M S
<i>aciculatus</i> (Hellén, 1915)	B M	<i>ornata</i> (Gravenhorst, 1829)	B M S
<i>agnoscendus</i> (Roman, 1939)	B	<i>ovivora ovivora</i> (Boheman, 1821)	B M S
<i>atratus</i> (Rudow, 1881)	B M S	<i>variabilis</i> (Holmgren, 1856)	B S
<i>brevicornis</i> (Tscheck, 1868)	B M S	<b>Zaglyptus</b> Foerster, 1868	
<i>cephalotes</i> (Holmgren, 1859)	*S	<i>multicolor</i> (Gravenhorst, 1829)	B M S
<i>cognator</i> (Thunberg, 1822)	*M	<i>varipes varipes</i> (Gravenhorst, 1829)	B M S
<i>curticornis</i> Perkins, 1943	B M	<b>Clistopyga</b> Gravenhorst, 1829	
<i>diversicostae</i> (Perkins, 1943)	M	<i>canadensis</i> Provancher, 1880	M
<i>dux</i> (Tscheck, 1868)	M	<i>incitator</i> (Fabricius, 1793)	B M S
<i>imperator</i> (Kriechbaumer, 1854)	B M S	<i>temporalis</i> (Hellén, 1949)	
<i>mastfestator</i> auct.		<i>rufator</i> Holmgren, 1856	S
<i>mesocentrus</i> (Gravenhorst, 1829)	B M S	 POLYSPHINCTINI	
<i>messor</i> <i>messor</i> (Gravenhorst, 1829)	*B M	 <b>Dreisbachia</b> Townes, 1962	
<i>populneus</i> (Ratzeburg, 1848)	B M S	<i>pictifrons</i> (Thomson, 1877)	B M
<i>pterelas</i> (Say, 1829)	*B	<b>Schizopyga</b> Gravenhorst, 1829	
<i>sericeus</i> (Hartig, 1847)	B*M S	<i>circulator circulator</i>	
<i>subglabratus</i> (Perkins, 1943)	*B M	Panzer, 1801)	M S
<i>geniculatus</i> (Kriechbaumer, 1896)		<i>frigida</i> Cresson, 1870	M
<i>terebrans terebrans</i> (Ratzeburg, 1844)	B M S	<i>varipes</i> Holmgren, 1856	B M
<i>tuberculatus tuberculatus</i> (Geoffroy in Fourcroy, 1785)	B M S	 <b>Zabrychypus</b> Cushman, 1920	
<b>Acropimpla</b> Townes, 1960		<i>primus</i> Cushman, 1920	M
<i>didyma</i> (Gravenhorst, 1829)	B M	gregori Šedivý, 1961	
		 <b>Aerodactyla</b> Haliday, 1838	
		<i>Colpomeria</i> Holmgren, 1859	
		<i>Polemophthorus</i> Schulz, 1911	

<i>degener</i> (Haliday, 1838)	*B		<i>Ephialtes</i> Schrank, 1802
<i>madida</i> (Haliday, 1838)	M S		suppressed.
<i>quadrisculpta</i> (Gravenhorst, 1820)	M S		
<b>Oxyrrhexis</b> Foerster, 1868			
<i>carbonator carbonator</i> Gravenhorst, 1807)	B M S		<i>capulifera</i> (Kriechbaumer, 1887) B M
<b>Polysphincta</b> Gravenhorst, 1829			<i>compunctor compunctor</i> Linnaeus, 1758) B M S
<i>boops</i> Tschek, 1868	B M S		<i>quadridentata</i> (Thomson, 1877) B M S
<i>clypeata</i> Holmgren, 1860	M S		<i>rufata</i> (Gmelin in Linnaeus, 1790) B M S
<i>rufipes</i> Gravenhorst, 1829	B		
<i>tuberosa tuberosa</i> (Gravenhorst, 1829)	B M		
<b>Sinarachna</b> Townes, 1960			<b>Pimpla</b> Fabricius, 1804
<i>anomala anomala</i> (Holmgren, 1860)	M S		<i>Coccygomimus</i> Saussure, 1892
<i>nigricornis</i> (Holmgren, 1860)	M		<i>aethiops</i> Curtis, 1828 *B
<i>pallipes pallipes</i> (Holmgren, 1860)	M		<i>aqilonia flavicoxis</i> Thomson, 1877 B M S
<b>Zatypota</b> Foerster, 1868			<i>arctica</i> Zetterstedt, 1838 B M
<i>bohemani</i> (Holmgren, 1860)	B M S		<i>contemplator</i> (Müller, 1776) B M S
<i>eximia eximia</i> Schmiedeknecht, 1907	M		<i>illecebrator</i> (Villers, 1789) M
<i>gracilis</i> (Holmgren, 1860)	B M S		<i>instigator</i> (Fabricius, 1793) B M S
<i>percontatoria</i> (Müller, 1776)	B M S		<i>melanacrias</i> Perkins, 1941 B M S
<i>discolor</i> (Holmgren, 1860)			<i>sodalis sodalis</i> Ruthe, 1859
<b>PIMPLINI</b>			<i>spuria</i> Gravenhorst, 1829 B M S
<b>Itoplectis</b> Foerster, 1868			<i>turionellae turionellae</i> (Linnaeus, 1758) B M S
<i>alternans alternans</i> (Gravenhorst, 1829)	B M S		<i>examinator</i> (Fabricius, 1804)
<i>clavicornis</i> (Thomson, 1889)	*B *S		<b>DELOMERISTINI</b>
<i>curticauda curticauda</i> (Kriechbaumer, 1887)	B M		<i>Theronia</i> Holmgren, 1859
<i>maculator maculator</i> (Fabricius, 1775)	B M S		<i>Pseudocoenites</i> Kriechbaumer 1892
<i>melanocephala</i> (Gravenhorst, 1829)	B S		<b>Theronia</b> s. str.
<i>viduata</i> (Gravenhorst, 1829)	B M S		<i>atalantae atalantae</i> (Poda, 1761) B M S
<b>Apechthis</b> Foerster, 1868			<i>laevigata laevigata</i> (Tschek, 1868) B M S
			<i>moravicus</i> (Kriechbaumer, 1892)
			<b>Perithous</b> Holmgren, 1859
			<b>Perithous</b> s. str.
			<i>albicinctus</i> (Gravenhorst, 1829) B
			<i>divinator</i> (Rossi, 1790) M S

<i>mediator mediator</i>			<i>curvipes</i> (Gravenhorst, 1829)
(Fabricius, 1804)	B M S		<i>obliterata</i> (Gravenhorst, 1829)
<i>Hymboischos</i> Baltazar, 1961			M S
<i>septemcinctarius</i>			
(Thunberg, 1822)	B M S		<b>Megarhyssa</b> Ashmead, 1900
<i>vartus</i> (Gravenhorst, 1829)			<i>Thalessa</i> Holmgren, 1859
<b>Delomerista</b> Foerster, 1868			<i>emarginatoria</i> (Thunberg, 1822)
<i>laevis</i> (Gravenhorst, 1829)	B M S		B
<i>mandibularis</i> (Gravenhorst, 1829)	M		<i>gigas</i> (Laxmann, 1770)      B      S
<b>Pseudorhyssa</b> Merrill, 1915			<i>perlata</i> (Christ, 1791)      B
<i>alpestris</i> (Holmgren, 1859)	M		<i>superba</i> (Schrank, 1781)      B
<b>POEMENIINI</b>			
Neoxoridini			
<b>Poemenia</b> Holmgren, 1859			<b>TRYPHONINAE</b>
<i>Calliclisis</i> Foerster, 1868			<b>SPHINCTINI</b>
<i>brachyura</i> (Holmgren, 1860)	M		
<i>collaris</i> Haupt, 1917	*B		<b>Sphinctus</b> Gravenhorst, 1829
<i>hectica</i> (Gravenhorst, 1829)	B M		<i>serotinus</i> Gravenhorst, 1829      *B M
<i>notata</i> Holmgren, 1859	B		
<b>Deuteroxorides</b> Viereck, 1914			<b>PHYTODIETINI</b>
<i>albitarsus</i> (Gravenhorst, 1829)	B M		<b>Phytodietus</b> Gravenhorst, 1829
<b>Podoschistus</b> Townes, 1957			<i>Neuchorus</i> Uchida, 1931
<i>scutellaris</i> (Desvignes, 1856)	B M S		<i>rufipes</i> Holmgren, 1860      B M
<b>Neoxorides</b> Clément, 1938			<i>Phytodietus</i> s. str.
<i>collaris</i> (Gravenhorst, 1829)	B M S		<i>albipes</i> Holmgren, 1856      B M
<i>montanus</i> Oehlke, 1966	B		<i>arcuatorius</i> (Thunberg, 1822)
<i>nitens</i> (Gravenhorst, 1829)	B M S		M
<i>varipes</i> (Holmgren, 1860)	*B		<i>crassitarsis</i> Thomson, 1877      M
<b>RHYSSINI</b>			<i>femoralis</i> Holmgren, 1860      M
<b>Rhyssa</b> Gravenhorst, 1829			<i>gelitorius</i> (Thunberg, 1822)
<i>amoena</i> Gravenhorst, 1829	M		B
<i>persuasoria</i> (Linnaeus, 1758)	B M S		<i>geniculatus</i> Thomson, 1877      M S
<b>Rhyssella</b> Rohwer, 1920			<i>griseanae</i> Kerrich, 1962      S
<i>approximator</i> (Fabricius, 1793)	B M S		<i>obscurus</i> Desvignes, 1856      M
			<i>ornatus</i> Desvignes, 1856      B S
			<i>polyzonias</i> (Foerster, 1771)
			B M S
			<i>segmentator</i> Gravenhorst, 1829      B M S
			<i>Weisia</i> Schmiedeknecht, 1907
			<i>bayeri</i> Gregor, 1935      M
			<b>Netelia</b> Gray, 1860
			<i>Paniscus</i> misident.
			<b>Paropheltes</b> Cameron, 1907
			<i>ornatus</i> (Vollenhoven, 1873)
			*B
			<i>longipes</i> (Brauns, 1889)

			TRYPHONINI
<i>tarsatus</i> (Brischke, 1880)	*B		<b>Grypocentrus</b> Ruthe, 1855
<i>thomsoni</i> (Brauns, 1889)	B		<i>albipes</i> Ruthe, 1855 M
<b>Prosthodocis</b> Enderlein, 1912			<i>basalis</i> Ruthe, 1855 M
<i>japonicus</i> Uchida, 1928	*B		<i>cinctellus</i> Ruthe, 1855 M
<i>Netelia</i> s. str.			
<i>dilatatus</i> (Thomson, 1888)	*B		<b>Polyblastus</b> Hartig, 1837
<i>fuscicornis</i> (Holmgren, 1858)	B M S		<b>Polyblastus</b> s. str.
<i>melanurus</i> (Thomson, 1888)	B		<i>conthurnatus</i>
<i>testaceus</i> (Gravenhorst, 1829)	B M S		(Gravenhorst, 1829) B M*S
<i>ocellaris</i> (Thomson, 1888)			<i>palaemon</i> Schioedte, 1839 *B M*S
<i>vinulae</i> (Scopoli, 1763)	B *S		<i>subalpinus</i> Holmgren, 1855 *B M*S
<b>Bessobates</b> Townes et Gupta, 1961			<i>carbonator</i> Kasparjan, 1970
<i>cristatus</i> (Thomson, 1888)	B S		<i>tener</i> Habermehl, 1909 *B*M
<i>latungulus</i> (Thomson, 1888)	B S		<i>wahlbergi wahlbergi</i>
<i>virgatus</i> (Geoffroy in Fourcroy, 1785)	*B S		Holmgren, 1855 *B M*S
THYMARIDINI			<i>wahlbergi varitarsus</i>
Eclytini			(Gravenhorst, 1829) B M*S
<b>Eclytus</b> Holmgren, 1855			<b>Labroctonus</b> Foerster, 1868
<b>Anoplectes</b> Kriechbaumer, 1896			<i>alternans</i> (Schioedte, 1839) *B M*S
<i>multicolor</i> Kriechbaumer, 1896	*B M		<i>melanostigmus</i>
<b>Eclytus</b> s. str.			Holmgren, 1855 M
<i>exornatus</i> (Gravenhorst, 1829)	B M		<i>pallicoxa</i> Thomson, 1888 *B
<i>ornatus</i> Holmgren, 1855	M		<i>stenocentrus</i> Holmgren, 1855 B M
<b>Hercus</b> Townes, 1969			<i>westringi</i> Holmgren, 1855 *B M*S
<i>frontalis frontalis</i> (Zetterstedt, 1838)	M		<b>Ctenochira</b> Foerster, 1855
<i>fontinalis</i> (Holmgren, 1855)	*B S		<i>angulata</i> (Thomson, 1883) *B M
<b>Thymaris</b> Foerster, 1868			<i>arcuata</i> (Holmgren, 1855) *B
<i>contaminatus</i> (Gravenhorst, 1829)	M		<i>flavicauda</i> (Roman, 1912) B M
<i>modestus</i> Schmiedeknecht, 1912	S		<i>infesta</i> (Holmgren, 1855) *B
<b>Hybophanes</b> Foerster, 1868			<i>genalis</i> (Thomson, 1883) *B M*S
<i>Oedemopsis</i> Tschek, 1868			<i>marginata</i> (Holmgren, 1855) *B*M
<i>scabriculus</i> (Gravenhorst, 1829)	M		<i>pastoralis</i> (Gravenhorst, 1829) M
			<i>pratensis</i> (Gravenhorst, 1829) M
			<i>propinqua</i> (Gravenhorst, 1829) M
			<i>rufipes</i> (Gravenhorst, 1829) B M
			<i>sanquinatoria</i> (Ratzeburg, 1852) B

<i>sphaerocephala</i>			<i>Psilosage</i> Foerster, 1868
(Gravenhorst, 1829)	B*M S		<i>abditus</i> Kasparjan, 1969
<i>validicornis</i> (Brischke, 1871)	B M S		<i>pleuralis</i> Thomson 1883
<i>vetusta</i> (Holmgren, 1855)	*B		<i>atriceps</i> Stephens, 1835
			*M*S
<b>Erromenus</b> Holmgren, 1855			<i>auricularis</i> Thomson, 1883
<i>analis</i> Brischke, 1871	*B		B M S
<i>brunnicanus</i> (Gravenhorst, 1829)	B M		<i>nigripes</i> Holmgren, 1855
<i>calcator</i> (Müller, 1776)	M		M
<i>junior</i> (Thunberg, 1822)	M		<i>relator</i> (Thunberg, 1822)
<i>melanotus</i> (Gravenhorst, 1829)	M		B M S
<i>plebejus</i> (Woldstedt, 1877)	*B M		<i>rutilator</i> (Linnaeus, 1761)
<i>punctulatus</i> Holmgren, 1855	*B M*S		B M S
<i>tarsator</i> Aubert, 1969	*M		<i>signator</i> Gravenhorst, 1829
<i>zonarius</i> (Gravenhorst, 1820)	*B*M*S		B M S
			<i>thomsoni</i> Roman, 1939
<b>Monoblastus</b> Hartig, 1837			M S
<i>Pseudopsilosage</i> Gregor, 1937			<i>trochanteratus</i> Holmgren, 1855
<i>brachyacanthus</i>			B M S
(Gmelin in Linnaeus, 1790)	B M S		<i>zavreli</i> Gregor, 1939
<i>fulvescens</i>			<i>Stenocrotaphon</i> Kasparjan 1969
Fonscolombe, 1849	S		<i>obtusator</i> (Thunberg, 1822)
<i>romani</i> (Gregor, 1937)			M S
<i>luteomarginatus</i>			<i>subsulcatus subsulcatus</i>
(Gravenhorst, 1829)	*B M S		Holmgren, 1855
<i>marginellus</i> (Gravenhorst, 1829)	*B M S		M
<i>proditor</i> (Gravenhorst, 1829)	M		<b>Symboethus</b> Foerster, 1868
			<i>bidentatus</i> Stephens, 1835
<b>Dyspetes</b> Foerster, 1868			*B M S
<i>arrogator</i> Heinrich, 1949	M S		<i>incestus</i> (Holmgren, 1855)
<b>Cosmoconus</b> Foerster, 1868			<i>brunniventris</i> Gravenhorst, 1829
<i>ceratophorus</i> (Thomson, 1888)	B M S		M
<i>elongator</i> (Fabricius, 1775)	B M S		<i>heliophilus</i> Gravenhorst, 1829
<i>meridionator</i> Aubert, 1963	*B		B M
<i>nigriventris</i> Kasparjan 1971	*M		<b>EXENTERINI</b>
			Cteniscini
<b>Tryphon</b> Fallén, 1813			<b>Kristetomus</b> Mason, 1962
<b>Tryphon</b> s. str.			<i>laetus</i> (Gravenhorst, 1829)
			M
			<i>laticeps</i> (Gravenhorst, 1829)
			B M
			<i>ridibundus</i> (Gravenhorst, 1829)
			M
			<i>triangulatorius</i>
			Gravenhorst, 1829)
			M
			<b>Cycasis</b> Townes
			in Townes, Momoi et Townes, 1965
			<i>rubiginosa</i> (Gravenhorst, 1829)
			B M S
			<b>Excavarus</b> Davis, 1897
			<i>apiarius</i> (Gravenhorst, 1829)
			B M S
			<b>Exyston</b> Schioedte, 1839
			<i>Tricamptus</i> Foerster, 1868

<i>albicinctus</i> (Gravenhorst, 1829)	M		<b>Eridolius</b> Foerster, 1868
<i>calcaratus</i> Thomson, 1883	M		<i>Antisoctenion</i> Foerster, 1868
<i>genalis</i> Thomson, 1883	B		<i>alacer</i> (Gravenhorst, 1829) B M
<i>pratorum</i> (Woldstedt, 1874)	B M S		<i>borealis</i> (Holmgren, 1855) M
<i>sponsorius</i> (Fabricius, 1781)	B M S		<i>curtisi</i> (Haliday, 1838) B M
<i>subnitidus</i> (Gravenhorst, 1829)	*B*M		<i>dorsator</i> Thunberg, 1822 M
<i>phaeorrhænus</i> (Haliday, 1838)			<i>elegans</i> (Stephens, 1835) *M
<b>Smicroplectrus</b> Thomson, 1883			<i>flavomaculatus</i> (Gravenhorst, 1829) M
<i>bohemani</i> (Holmgren, 1855)	B M		<i>gibbulus</i> (Holmgren, 1855) M
<i>erosus</i> (Holmgren, 1855)	B M		<i>gnathoxanthus</i> (Gravenhorst, 1829) B M
<i>excisus</i> Kerrich, 1952	M		<i>hofferi</i> (Gregor, 1937) M
<i>jucundus</i> (Holmgren, 1855)	*B		<i>marginatus</i> (Thomson, 1883) M
<i>perkinsorum</i> Kerrich, 1952	B M		<i>mitigiosus</i> (Gravenhorst, 1829) M
<i>quinquecinctus</i> (Gravenhorst, 1820)	B M S		<i>pictus</i> (Gravenhorst, 1829) M
<i>trianguligena</i> Kerrich, 1952			<i>pumilus</i> (Holmgren, 1855) B M
<b>Acrotomus</b> Holmgren, 1855			<i>rufilabris</i> (Holmgren, 1855) M
<i>lucidulus</i> (Gravenhorst, 1829)	M S		<i>similis</i> (Holmgren, 1855) M
<i>succinctus</i> (Gravenhorst, 1829)	B M		<b>IDIOPRAGMATINI</b>
<b>Cteniscus</b> Haliday, in Curtis, 1832			<b>Idiogramma</b> Foerster, 1868
<i>devius</i> (Mason, 1955)	M		<i>uryops</i> Schmiedeknecht, 1888 *B
<i>pedatorius</i> (Panzer, 1809)	M		<b>EUCEROTINI</b>
<b>Exenterus</b> Hartig, 1837			<b>Euceros</b> Gravenhorst, 1829
<i>abruptorius</i> (Thunberg, 1822)	B M S		<i>albitarsus</i> Curtis, 1837 M
<i>adspersus</i> Hartig, 1838	B M S		<i>pruinosus pruinosus</i> (Gravenhorst, 1829) *B M
<i>amicotorius</i> (Panzer, 1801)	B M S		<i>serricornis</i> (Haliday, 1838) M
<i>marginatorius</i> (Fabricius, 1793)			<i>superbus</i> Kriechbaumer, 1888 S
<i>claripennis</i> Thomson, 1883	B		<b>LABENINAE</b>
<i>confusus</i> Kerrich, 1952	B M S		<b>Brachycyrtus</b> Kriechbaumer, 1880
<i>ictericus</i> (Gravenhorst, 1829)	M		<i>ornatus</i> Kriechbaumer, 1880 B M S
<i>oriolus</i> Hartig, 1838	B M S		<b>ADELOGNATHINAE</b>
<i>simplex</i> Thomson, 1883	B M		<b>Adelognathus</b> Holmgren, 1855
<i>tricolor</i> , Roman, 1913	B *S		<i>chrysopygus</i> (Gravenhorst, 1829) B M S
<i>vellicatus</i> Cushman, 1940	B *S		

<i>dorsalis</i> (Gravenhorst, 1829)	B	<i>brachylabis</i> (Kriechbaumer, 1889)	B M		
<i>nigriceps</i> Thomson, 1888	M	<i>depressus</i> (Holmgren, 1860)	B		
<i>nigricornis</i> Thomson, 1888	M	<i>filiformis</i> (Gravenhorst, 1829)	B M S		
<i>nigrifrons</i> Holmgren, 1855	M	<i>fuligator</i> (Thunberg, 1822)	B M		
<i>pallipes</i> (Gravenhorst, 1829)	B	<i>gracilicornis</i> Gravenhorst, 1829	B M S		
<i>pusillus</i> Holmgren, 1855	M	<i>gravenhorsti</i> (Curtis, 1831)	B M S		
XORIDINAE					
<b>Ischnoceros</b> Gravenhorst, 1829		<i>securicornis</i> (Holmgren, 1860)			
<i>caligatus</i> (Gravenhorst, 1829)	B M S	<i>hedwigi</i> (Clément, 1938)	M		
<i>rusticus</i> (Geoffroy in Fourcroy, 1785)	B M S	<i>indicarius</i> Latreille, 1809	S		
<i>filicornis</i> Kriechbaumer, 1879		<i>irrigator</i> (Fabricius, 1793)	B M		
<b>Odontocolon</b> Cushman, 1942		<i>planus</i> Šedivý, 1958			
<i>appendiculatus</i> (Gravenhorst, 1829)	B M S	<i>minutus</i> Clément, 1938	*S		
<i>dentipes</i> (Gmelin in Linnaeus, 1790)	B M S	<i>niger</i> (Pfeffer, 1913)	B M		
<i>pinetorum</i> (Thomson, 1877)		<i>praecatorius</i> (Fabricius, 1793)	B M S		
<i>geniculatus</i> (Kriechbaumer, 1889)	M	<i>sepulchralis</i> (Holmgren, 1860)	S		
<i>punctulatus</i> (Thomson, 1877)	B M S	<b>Moerophora</b> Foerster, 1868			
<i>quercinus</i> (Thomson, 1877)	*B	<i>ephialtooides</i> (Kriechbaumer, 1882)	*M*S		
<i>rufiventris</i> (Holmgren, 1860)	M S	<i>rufipes</i> (Gravenhorst, 1829)	B S		
<i>spinipes</i> (Gravenhorst, 1829)	B M S	AGRIOTYPINAE			
<i>thomsoni</i> Clément, 1938	*S	<b>Agriotypus</b> Curtis, 1832			
<b>Xorides</b> Latreille, 1809		<i>armatus</i> Curtis, 1832	B M S		
<b>Gonophonus</b> Foerster, 1868		PHYGADEUONTINAE			
<i>annulator</i> (Fabricius, 1804)	M	Cryptinae			
<i>corycynensis</i> (Kriechbaumer, 1894)	M S	Hemitelinae			
<i>propinguus</i> Tschek, 1868	M S	Gelinae			
<b>Xorides</b> s. str.		PHYGADEUONTINI			
<i>Xylonomus</i> Gravenhorst, 1829		Hemitelini			
<i>Sichelia</i> Foerster, 1868		Gelini			
<i>Rhadinopimpla</i> Schulz, 1906		<b>Chirotica</b> Foerster, 1868			
<i>Xylonominus</i> Clément, 1938		<i>decorator</i> (Villers, 1789)	M S		
<i>alpestris</i> Habermehl, 1904	B	<i>trifasciator</i> (Thunberg, 1822)			
<i>ater</i> (Gravenhorst, 1829)	B M	<i>insignis</i> (Gravenhorst, 1829)	*B*M*S		
<i>baueri</i> (Clément, 1938)	B	<i>maculipennis</i> (Gravenhorst, 1829)	M S		
<b>Encrateola</b> Strand, 1916					
<i>laevigata</i> (Ratzeburg, 1848)				B M S	

<b>Eudelus</b> Foerster, 1868		<i>stiavnicense</i> (Čapek, 1956)	S
<i>scabriculus</i> (Thomson, 1884)	M S		
<i>simillimus</i> (Taschenberg, 1865)	*B		
<b>Acrolyta</b> Foerster, 1868		<b>Dichrogaster</b> Doumerc, 1855	
<i>distincta</i> (Bridgman, 1883)	*B*M	<i>aestivalis</i> (Gravenhorst, 1829)	B M S
<i>marginata</i> (Bridgman, 1883)	B M	<i>classicornis</i> Horstmann, 1976	B
<i>submarginata</i> (Bridgman, 1883)	B	<i>genalis</i> (Habermehl, 1925)	M
<b>Diaglyptidea</b> Viereck, 1913		<i>heteropus</i> (Thomson, 1896)	M
<i>conformis</i> (Gmelin in Linnaeus, 1790)	B M S	<i>liostylus</i> (Thomson, 1885)	B S
<b>Lysibia</b> Foerster, 1868		<i>schaffneri</i> (Schmiedeknecht, 1897)	
<i>nana</i> (Gravenhorst, 1829)	B M S	<i>longicaudatus</i> (Thomson, 1884)	B
<i>fulvipes</i> (Gravenhorst, 1829)		<i>nigrithorax</i> Horstmann, 1971	*B
<b>Obisiphaga</b> Morley, 1907		<b>Xenolytus</b> Foerster, 1868	
<i>stenoptera</i> (Marshall, 1868)	*B	<i>bitinctus</i> (Gmelin in Linnaeus, 1790)	*B *S
<b>Arotrepes</b> Townes, 1970		<b>Gelis</b> Thunberg, 1827	
<i>perfusor</i> (Gravenhorst, 1829)	M	<i>Pezomachus</i> Gravenhorst, 1829	
<i>speculator</i> (Gravenhorst, 1829)	*B	<i>Pezolochus</i> Foerster, 1850	
<b>Xiphulcus</b> Townes, 1970		<i>Thaumatotyphus</i> Foerster, 1868	
<i>floricator</i> (Gravenhorst, 1807)	B M	<i>acarorum</i> (Linnaeus, 1758)	B M
<b>Hemiteles</b> Gravenhorst, 1829		<i>agilis</i> (Fabricius, 1775)	B M S
<i>bipunctator</i> (Thunberg, 1822)	B M S	<i>alopecosae</i> Horstmann, 1986	B
<i>similaris</i> (Gmelin in Linnaeus, 1790)	B M	<i>anthracinus</i> (Foerster, 1850)	B
<i>unicolor</i> Thomson, 1884		<i>aquisgranensis</i> (Foerster, 1850)	B
<b>Aclastus</b> Foerster, 1868		<i>neesi</i> (Foerster, 1850)	
<i>gracilis</i> (Thomson, 1884)	B	<i>areator</i> (Panzer, 1804)	B M S
<i>pilosus</i> Horstmann, 1980	B	<i>pulchellus</i> (Gravenhorst, 1829)	
<i>solutus</i> (Thomson, 1884)	B	<i>astutus</i> (Foerster, 1850)	M
<b>Polyaulon</b> Foerster, 1868		<i>attentus</i> (Foerster, 1850)	M
<i>Rhacodopteron</i> Čapek, 1956		<i>balteata</i> (Thomson, 1885)	*S
<i>paradoxus</i> (Zetterstedt, 1838)	B M S	<i>brassicae</i> Horstmann, 1986	B M S
<i>billupsi</i> (Bridgman, 1882)		<i>cautus</i> (Foerster, 1850)	M
		<i>cinctus</i> (Linnaeus, 1758)	B M S
		<i>bicolorinus</i> (Gravenhorst, 1829)	
		<i>circumcinctus</i> (Foerster, 1850)	B
		<i>confusus</i> (Bridgman, 1883)	B

<i>corruptor</i> (Foerster, 1850)	B	M
<i>cursitans</i> (Fabricius, 1775)	B	M
<i>distinctus</i> (Foerster, 1850)	B	M
<i>festinans</i> (Fabricius, 1798)	M	
<i>formicarius</i> (Linnaeus, 1758)	B	
<i>fraudulentus</i> (Foerster, 1850)	M	
<i>gentilis</i> (Foerster, 1850)	B	S
<i>gonatopinus</i> (Thomson, 1884)	B	
<i>gracilis</i> (Foerster, 1850)	B	
<i>hortensis</i> (Gravenhorst, 1829)	B	M
<i>hostilis</i> (Foerster, 1850)	B	
<i>impotens</i> (Foerster, 1850)	M	
<i>inermis</i> (Foerster, 1850)	B	
<i>insolens</i> (Foerster, 1850)	M	
<i>instabilis</i> (Foerster, 1850)	B	M
<i>intermedius</i> (Foerster, 1850)	M	
<i>kiesenwetteri</i> (Foerster, 1850)	B	
<i>latrator</i> (Foerster, 1850)	*M	
<i>lichtensteinii</i> (Pfankuch, 1913)	*B	*S
<i>lucidulus</i> (Foerster, 1850)	B	
<i>mandibularis</i> (Thomson, 1884)	B	M
<i>marginatus</i> (Foerster, 1851)	B	
<i>melanarius</i> (Gravenhorst, 1829)	M	S
<i>melanocephalus</i> (Schrank, 1781)	B	M S
<i>melanogaster</i> (Thomson, 1884)	B	M
<i>micrurus</i> (Foerster, 1850)	*B	
<i>nigricornis</i> (Foerster, 1850)	*B	
<i>nigritus</i> (Foerster, 1850)	M	
<i>notabilis</i> (Foerster, 1850)	B	
<i>ochraceus</i> (Foerster, 1850)	B	
<i>pedicularius</i> (Fabricius, 1793)	B	
<i>picipes</i> (Gravenhorst, 1829)	B	
<i>proximus</i> (Foerster, 1850)	B	
<i>prudens</i> (Foerster, 1851)	*B	
<i>pulicarius</i> (Fabricius, 1793)		S
<i>pumilus</i> (Foerster, 1850)	*B	
<i>rufulus</i> (Foerster, 1850)	M	
<i>ruficornis</i> Thunberg, 1827	M	
<i>rufipes</i> (Foerster, 1850)	*B	
<i>spinulus</i> (Thomson, 1884)	*B	
<i>sulcatus</i> (Blunck, 1951)	B	M S
<i>sylvicolus</i> (Foerster, 1850)		*B
<i>terebrator</i> (Ratzeburg, 1848)	B	
<i>timidus</i> (Foerster, 1850)	M	
<i>vagabundus</i> (Foerster, 1850)		M
<i>vagans</i> (Olivier, 1792)	B	M S
<i>vigil</i> (Foerster, 1850)	B	
<i>vorax</i> (Foerster, 1850)	M	
<i>vulpinus</i> (Gravenhorst, 1815)	B	M
<i>zonatus</i> (Foerster, 1850)	*B	
<b>Catalytus</b> Foerster, 1851		
<i>fulveolatus</i> (Gravenhorst, 1829)	B	
<i>mangeri</i> (Gravenhorst, 1829)	B	
<b>Agasthenes</b> Foerster, 1868		
<i>varitarsus</i> (Gravenhorst, 1829)		*B
<b>Mastrus</b> Foerster, 1868		
<i>Isadelphus</i> Foerster, 1868		
<i>Cecidonomus</i> Bridgman, 1880		
<i>armatus</i> (Gravenhorst, 1829)	B	
<i>carbonarius</i> (Schmiedeknecht, 1905)	B	
<i>castaneus</i> (Taschenberg, 1865)	B	M
<i>coriarius</i> (Taschenberg, 1865)	*B	M
<i>inimicus</i> (Gravenhorst, 1829)	B	M S
<i>ridibundus</i> (Gravenhorst, 1829)	M	
<i>rufulus</i> Thomson, 1884	B	M
<i>luteiventris</i> (Gravenhorst, 1829)		

<i>sordipes</i> (Gravenhorst, 1829)	B M S
<i>varicoxis</i> (Taschenberg, 1865)	B
<b>Mastrulus</b> Horstmann, 1979	
<i>capra</i> (Thomson, 1884)	*M
<b>Zoophthora</b> Foerster, 1868	
<i>dodecellae</i> (Obertel et Šedivý, 1960)	M
<i>hirticeps</i> (Thomson, 1885)	*B
<i>incisus</i> Bridgman, 1883	B
<i>platygaster</i>	
(Schmiedeknecht, 1897)	M
<i>plumbeus</i> (Thomson, 1884)	*B*M
<i>punctiventris</i> (Thomson, 1884)	M
<b>Helcostizus</b> Taschenberg, 1865	
<i>restaurator restaurator</i> (Fabricius, 1775)	B M S
<i>albator</i> (Thunberg, 1822)	
<i>brachycentrus</i> (Gravenhorst, 1829)	
<b>Rhembobius</b> Foerster, 1868	
<i>nigriceps</i> (Thomson, 1883)	B M
<i>nigricollis</i> (Thomson, 1883)	M
<i>perscrutator</i> (Thunberg, 1822)	B M
<i>quadrispinus</i> (Gravenhorst, 1829)	B M S
<b>Ethelurgus</b> Foerster, 1868	
<i>sodalis</i> (Taschenberg, 1865)	B M S
<i>vulnerator</i> (Gravenhorst, 1829)	*B
<b>Charitopes</b> Foerster, 1868	
<i>chrysopae</i> (Brischke, 1890)	S
<i>melanogaster</i> (Thomson, 1884)	B M
<b>Medophron</b> Foerster, 1868	
<i>afflictor</i> (Gravenhorst, 1829)	S
<i>armatulus</i> (Thomson, 1888)	M
<b>Endasys</b> Foerster, 1868	
<i>amoenus</i> (Habermehl, 1912)	B
<i>analis</i> (Thomson, 1883)	S
<i>brevis</i> (Gravenhorst, 1829)	B M S
<i>erythrogaster</i>	
(Gravenhorst, 1829)	B M S
<i>euxestus</i> (Speiser, 1891)	M
<i>kriegeri</i> Habermehl, 1912	B M
<i>rubricator</i> (Thunberg, 1822)	M S
<i>rusticus</i> (Habermehl, 1912)	B M S
<i>testaceipes</i> (Brischke, 1891)	B M S
<b>Glyphtinemis</b> Foerster, 1868	
<i>clypealis</i> (Thomson, 1883)	*B M
<i>profligator</i> (Fabricius, 1775)	B M S
<i>vagabunda</i> (Gravenhorst, 1829)	B. M S
<b>Bathythrix</b> Foerster, 1868	
<i>Ischnurgops</i> Foerster, 1868	
<i>aereus</i> (Gravenhorst, 1829)	B
<i>argentatus</i> (Gravenhorst, 1829)	B M S
<i>lacustris</i> (Schmiedeknecht, 1902)	
<i>claviger</i> (Taschenberg, 1865)	B M S
<i>collaris</i> (Thomson, 1896)	B M S
<i>ornatus</i> Kříštek et Obertel, 1958	
<i>decipiens</i> (Gravenhorst, 1829)	B
<i>fragilis</i> (Gravenhorst, 1829)	B M S
<i>bellulus</i> (Kriechbaumer, 1892)	
<i>laminus</i> (Thomson, 1884)	B *S
<i>linearis</i> (Gravenhorst, 1829)	M
<i>maculatus</i> (Hellén, 1957)	B
<i>montanus</i>	
(Schmiedeknecht, 1902)	M

<i>pellucidator</i>			
(Gravenhorst, 1829)	B	M	S
<i>prominens</i> (Strobl, 1900)	B		
<i>rugulosus</i> (Thomson, 1884)	B		
<i>strigosus</i> (Thomson, 1884)	B	M	
<i>tenuis</i> (Gravenhorst, 1829)	M		
<i>thomsoni</i> (Kerrich, 1942)	B	S	
<b>Sulcarius</b> Townes, 1970			
<i>biannulatus</i>			
(Gravenhorst, 1829)	B		
<i>hellbachi</i> (Schmiedeknecht, 1905)	M		
<b>Tropistes</b> Gravenhorst, 1829			
<i>falcatus</i> (Thomson, 1884)	B	M	S
<i>nitidipennis</i>			
Gravenhorst, 1829	S		
<b>Tricholinum</b> Foerster, 1868			
<i>ischonoceros</i> (Thomson 1888)		*M	
<b>Orthizema</b> Foerster, 1868			
<i>hadrocerum</i> (Thomson, 1884)	M		
<i>melanopygum</i> (Gravenhorst, 1829)	B		
<i>subannulatum</i> (Bridgman, 1883)	M		
<i>triangulatum</i> (Thomson, 1884)	M		
<b>Gnotus</b> Foerster, 1868			
<i>chionops</i> (Gravenhorst, 1829)	M		
<i>tenuipes</i> (Gravenhorst, 1829)	*B		
<b>Stibeutes</b> Foerster, 1868			
<i>curvispina</i> (Thomson, 1884)	*B		
<i>bonellii</i> (Gravenhorst, 1829)	*B		
<i>gravenhorstii</i> (Foerster, 1850)	B		
<b>Theroscopus</b> Foerster, 1850			
<i>Thysiotorus</i> Foerster, 1868			
<i>annulicornis</i> (Thomson, 1884)	M		
<b>dromicus</b> (Gravenhorst, 1815)		M	
<i>cingulatus</i> (Foerster, 1851)			
<b>hemipterus</b> (Fabricius, 1793)	B	M	S
<i>melanopygus</i> (Gravenhorst, 1829)	B		
<i>micator</i> (Gravenhorst, 1807)	B	M	
<i>pedestris</i> (Fabricius, 1775)	M		
<i>subzonatus</i> (Gravenhorst, 1829)	S		
<b>Megacara</b> Townes, 1970			
<i>alteareolatus</i>			
(Schmiedeknecht, 1905)		*M	
<i>hortulana</i> (Gravenhorst, 1829)		*B	
<i>vagans</i> (Gravenhorst, 1829)		M	S
<b>Phygadeuon</b> Gravenhorst, 1829			
<i>britannicus</i> Habermehl, 1923		M	
<i>canaliculatus</i> Thomson, 1889	M		
<i>clypearis</i> Strobl, 1900	M		
<i>cylindraceus</i> Ruthe, 1859	M		
<i>dimidiatus</i> Thomson, 1884	M		
<i>dumetorum</i> Gravenhorst, 1829		*B	
<i>exiguus</i> Gravenhorst, 1829	B		
<i>flavimanus</i> Gravenhorst, 1829	B		
<i>forticornis</i> (Kriechbaumer, 1892)	B	M	
<i>fumator</i> Gravenhorst, 1829	B		
<i>geniculatus</i> (Kriechbaumer, 1892)	B	M	
<i>grandiceps</i> Thomson, 1884	B	M	
<i>hercynicus</i> Gravenhorst, 1829	B	M	
<i>inflatus</i> Thomson, 1884	M		
<i>interstitialis</i>			
(Schmiedeknecht, 1897)	S		
<i>leucostigmus</i>			
Gravenhorst, 1829	B	M	
<i>nitidus</i> Gravenhorst, 1829	B		
<i>ovaliformis</i> Dalla Torre, 1902	B	M	

<i>ovatus</i> Gravenhorst, 1829	M	<i>compressus</i> Thomson, 1884
<i>pegomyiae</i> Habermehl, 1928	B M	<i>exilis</i> Haliday, 1838 B M
<i>rozsyppali</i> Gregor, 1941	M	<i>gravidus</i> Gravenhorst, 1829 M
<i>ruficornis</i> (Gravenhorst, 1829)	M	<i>oreophilus</i> Foerster, 1876 B M
<i>rugulosus</i> Gravenhorst, 1829	B M	<i>picipes</i> Holmgren, 1860 B
<i>scapus</i> Thomson, 1884	B M S	<i>tenuipes</i> Thomson, 1884 M
<i>wiesmanni</i> Sachtleben, 1934		
<i>subtilis</i> Gravenhorst, 1829	M	<b>Cremnoides</b> Foerster, 1850
<i>thomsoni</i> Roman, 1924	B M	<i>atricapillus</i> (Gravenhorst, 1815) *B
<i>trichops</i> Thomson, 1884	B M	
<i>troglodytes</i> Gravenhorst, 1829	M	<b>HEMIGASTERINI</b>
<i>variabilis</i> Gravenhorst, 1829	B	Aptesini
<i>vexator</i> (Thunberg, 1822)	B	Echthrinini
<b>Stilpnus</b> Gravenhorst, 1829		<b>Demopheles</b> Foerster, 1868
<i>blandus</i> Gravenhorst, 1829	B M S	<i>corruptor</i> (Taschenberg, 1865) B
<i>crassicornis</i> Thomson, 1884	M	<i>Javra</i> Cameron, 1903
<i>gagates</i> (Gravenhorst, 1829)	B M S	<i>opaca</i> (Thomson, 1873) B
<i>pavoniae</i> (Scopoli, 1763)	B M	<b>Parmortha</b> Townes in Townes et Gupta, 1962
<i>tenebricosus</i> (Gravenhorst, 1829)	B M S	<i>pleuralis</i> (Thomson, 1873) M
<i>tenuipes</i> Thomson, 1884	M	<b>Cratocryptus</b> Thomson, 1873
<b>Mesoleptus</b> Gravenhorst, 1829		<i>furcator</i> (Gravenhorst, 1829) M
<i>Exolytus</i> Holmgren, 1858		<b>Cubocephalus</b> Ratzeburg, 1848
<i>filicornis</i> (Thomson, 1884)	B	<i>Chaeretimma</i> Foerster, 1868
<i>flavipes</i> (Thomson, 1884)	B	<i>anatorius</i> (Gravenhorst, 1829) M
<i>gracilis</i> Foerster, 1876	M	<i>associator</i> (Thunberg, 1822) B M
<i>laevigatus laevigatus</i> (Gravenhorst, 1820)	B M	<i>leucopsis</i> (Gravenhorst, 1829)
<i>marginatus</i> (Thomson, 1884)	M	<i>brevicornis</i> (Taschenberg, 1865) B
<i>petiolaris</i> (Thomson, 1884)	M	<i>oviventris</i> (Gravenhorst, 1829)
<i>ripiculus</i> (Thomson, 1884)	M	<i>distinctor</i> (Thunberg, 1822) B
<i>scrutator</i> (Haliday, 1838)	B M	<i>erythrinus</i> (Gravenhorst, 1829) M
<i>incertus</i> (Foerster, 1876)		<i>lacteator</i> (Gravenhorst, 1829) B
<i>strandii</i> Gregor, 1937	B M	<i>nigriventris</i> (Thomson, 1874) S
<b>Atractodes</b> Gravenhorst, 1829		<i>sternocerus</i> (Thomson, 1873) B M
<i>bicolor</i> Gravenhorst, 1829	B	
<i>croceicornis</i> Haliday, 1838	B	<b>Oresbius</b> Marshall, 1867

<i>arridens</i> (Gravenhorst, 1829)	*B M	<i>brachypterus</i> (Gravenhorst, 1815)	B M S
<i>galactinus</i> (Gravenhorst, 1829)	B M	<i>micropterus</i> Gravenhorst, 1829]	B M S
<i>septentrionalis</i> (Thomson, 1883)	B	<i>larvatus</i> (Gravenhorst, 1829)	B
<b>Echthrus</b> Gravenhorst, 1829		<i>sericans</i> (Gravenhorst, 1829)	*B
<i>reluctator</i> <i>reluctator</i> (Linnaeus, 1758)	B M	<i>sperator</i> (Müller, 1776)	*M
<b>Polytribax</b> Foerster, 1868		<i>vestigialis</i> (Foerster, 1850)	*B M
<i>arrogans</i> (Gravenhorst, 1829)	B	<b>Aptesis</b> Foerster, 1850	
<i>perspicillator</i> (Gravenhorst, 1807)	B M*S	<i>abdominator</i> (Gravenhorst, 1829)	B M
<i>rufipes</i> (Gravenhorst, 1829)	*B	<i>bifrons</i> (Gmelin in Linnaeus, 1790)	B
<b>Aconias</b> Cameron, 1904		<i>cretata</i> (Gravenhorst, 1829)	B M
<i>tarsatus</i> (Bridgman, 1881)	*B M	<i>desertor</i> (Gravenhorst, 1829)	B M
<b>Plectocryptus</b> Thomson, 1874		<i>femoralis</i> (Thomson, 1883)	B M
<i>digitatus</i> (Gmelin in Linnaeus, 1790)	B	<i>flagitator</i> (Rossi, 1794)	B M S
<b>Colocnema</b> Foerster, 1868		<i>galactinus</i> (Gravenhorst, 1829)	B M
<i>Coelocryptus</i> Thomson, 1873		<i>gracilicornis</i> (Kriechbaumer, 1891)	*B
<i>rufina</i> (Gravenhorst, 1829)	B M S	<i>nigritula</i> (Thomson, 1885)	M
<b>Giraudia</b> Foerster, 1868		<i>nigrocincta</i> (Gravenhorst, 1829)	B M
<i>grisescens</i> (Gravenhorst, 1829)	M	<i>puncticollis</i> (Thomson, 1883)	M S
<i>gyratoria</i> (Thunberg, 1822)	*B M	<i>scopulifer</i> (Obrtel, 1953)	M
<b>Schenkia</b> Foerster, 1868		<i>subguttata</i> (Gravenhorst, 1829)	B M S
<i>crassicornis</i>		<i>terminata</i> (Gravenhorst, 1829)	M
(Kriechbaumer, 1891)	M	<i>tricincta</i> (Gravenhorst, 1829)	*B
<i>graminicola</i> (Gravenhorst, 1829)	M	<b>Listrocryptus</b> Brauns, 1905	
<i>improba</i> (Gravenhorst, 1829)	M	<i>spatulatus</i> Brauns, 1905	*S
<i>opacula</i> (Thomson, 1883)	*M	<b>MESOSTENINI</b> <i>Cryptini</i>	
<b>Megaplectes</b> Foerster, 1868		<b>Apsilops</b> Foerster, 1868	
<i>monticola</i> <i>monticola</i> (Gravenhorst, 1829)	B M*S	<i>Dapanus</i> Foerster, 1868	
<b>Pleolophus</b> Townes, 1962		<i>Trichocryptus</i> Thomson, 1873	
<i>basizonus</i> (Gravenhorst, 1829)	B M S	<i>aquaticus</i> (Thomson, 1874)	*B

<i>cinctarius</i> (Fabricius, 1775)	B	S	<i>bellosa</i> (Curtis, 1837)	B	M	S
<b>Thrybius</b> Townes in Townes, Momoi et Townes, 1965			<i>signatorius</i> (Fabricius, 1793)			
<i>brevispina</i> (Thomson, 1896)		*S	<i>clementi</i> (Habermehl, 1926)	B		
<i>praedator</i> (Rossi, 1792)	B	*S	<i>confector</i> (Gravenhorst, 1829)	M		
<i>drewseni</i> (Thomson, 1873)			<i>coxator</i> (Tschech, 1870)	M		
<i>leucopygus</i> (Gravenhorst, 1829)			<i>dubia</i> (Taschenberg, 1865)	M		
<b>Agrotherutes</b> Foerster, 1868			<i>fugitiva</i> (Gravenhorst, 1829)	B	S	
<i>abbreviator</i> (Fabricius, 1793)		B M S	<i>heliofila</i> (Tschech, 1870)	B		
<i>adustus</i> (Gravenhorst, 1829)	B	S	<i>nigripes</i> (Gravenhorst, 1829)	M	S	
<i>aterrimus</i> (Gravenhorst, 1829)		*B	<b>Pycnocryptus</b> Thomson, 1873			
<i>bicinctulus</i> (Gravenhorst, 1829)			<i>director</i> (Thunberg, 1822)	B	M	S
<i>fumipennis</i> (Gravenhorst, 1829)		B M S	<i>longicaudus</i> (Kriechbaumer, 1873)	M		
<i>zygaenarum</i> (Thomson, 1873)			<b>Hydryta</b> Foerster, 1869			
<i>grossus</i> (Gravenhorst, 1829)		B M S	<i>Brachycryptus</i> Thomson, 1873			
<i>hospes</i> (Tschech, 1870)	B	M	<i>fusiventris</i> (Thomson, 1873)	*M		
<i>mandator</i> (Linnaeus, 1758)		S	<i>sordidulus</i> (Thomson, 1873)	*B		
<i>nasutus</i> (Thomson, 1873)		*S	<b>Idiolispa</b> Foerster, 1868			
<i>tibialis</i> (Thomson, 1873)	*B		<i>analisis analis</i> (Gravenhorst, 1807)	B		
<i>tricolor</i> (Gravenhorst, 1829)		S	<b>Trychosia</b> Foerster, 1868			
<b>Mesostenidea</b> Viereck, 1914			<i>glabricula</i> (Thomson, 1873)	*B	M	S
<i>ligator</i> (Gravenhorst, 1829)		B M S	<i>legator</i> (Thunberg, 1822)	B		
<b>Gambrus</b> Foerster, 1868			<i>plebejus</i> (Tschech, 1870)			
<i>Hygrocryptus</i> Thomson, 1873			<i>mesocastana</i> (Tschech, 1870)	*B	M	S
<i>carnifex</i> (Gravenhorst, 1829)		B M S	<i>neglecta</i> (Tschech, 1870)	*B	M	S
<i>incubitor</i> (Linnaeus, 1758)		*B	<i>pauper</i> (Tschech, 1870)	*B		
<i>inferus</i> Thomson, 1896	B	M	<i>tristator</i> (Tschech, 1870)	S		
<i>nigrifemur</i> Szépligeti, 1916		S	<b>Enclisia</b> Townes, 1970			
<i>opacus</i> Szépligeti, 1916		S	<i>laticrus</i> (Thomson, 1896)	M		
<i>ornatulus</i> (Thomson, 1873)		*B	<i>macilentus</i> (Gravenhorst, 1829)	B	M	
<i>superus</i> (Thomson, 1896)	*B	M	<i>vindex</i> (Tschech, 1870)	B	M	
<b>Aritranis</b> Foerster, 1868			<i>polytomi</i> (Tschech, 1872)			
<i>Hoplocryptus</i> Thomson, 1873			<b>Ischnus</b> Gravenhorst, 1829			
			<i>alternator</i> (Gravenhorst, 1829)	M		

<i>facialis</i> (Szepligeti, 1916)	S	<i>spinosus</i> Gravenhorst, 1829	B M*S
<i>inquisitorius inquisitorius</i> (Müller, 1776)	B M S	<i>armatorius</i> (Fabricius, 1787)	
<i>assertorius</i> (Fabricius, 1793)		<i>spiralis</i> (Geoffroy in Fourcroy, 1785)	B M
<i>insulans</i> (Krieger, 1897)	M	<i>triguttatus</i> Gravenhorst, 1829	B M
<i>migrator</i> (Fabricius, 1775)	M S	<i>viduatorius</i> Fabricius, 1804	B M S
<i>minutorius</i> (Fabricius, 1804)	*B M	<b>Xylophrurus</b> Foerster, 1868	
<i>punctiger</i> (Thomson, 1896)	M S	<i>apum</i> (Thomson, 1873)	B
<b>Caenocryptus</b> Thomson, 1873		<i>dispar</i> (Thunberg, 1822)	B M S
<i>rufiventris</i> (Gravenhorst, 1829)	B	<b>Myrmeleonostenus</b> Uchida, 1936	
<i>striolatus</i> Thomson, 1896	M	<i>italicus</i> (Gravenhorst, 1829)	M
<b>Nippocryptus</b> Uchida, 1936		<b>Meringopus</b> Foerster, 1868	
<i>vittatorius</i> (Jurine, 1807)	B M S	<i>calescens calescens</i> (Gravenhorst, 1829)	*B*M
<i>sexannulatus</i> (Gravenhorst, 1829)		<i>cyanator</i> (Gravenhorst, 1829)	B M
<b>Buathra</b> Cameron, 1903		<i>nigerrimus murorum</i> (Tschek, 1872)	*B*M
<i>divisoria</i> (Tschek, 1872)	M	<i>titillator</i> (Linnaeus, 1758)	B
<i>laborator</i> (Thunberg, 1822)	B M	<b>Synechocryptus</b> Smiedeknecht, 1904	
<i>tarsoleuca</i> (Schrank, 1781)	B M	<i>mactator</i> (Tschek, 1870)	*S
<b>Itamoplex</b> Foerster, 1868		<i>erberi</i> (Tschek, 1871)	
<i>Cryptus</i> Fabricius, 1804		<b>Mesostenus</b> Gravenhorst, 1829	
<i>apparitorius</i> (Villers, 1789)	B	<i>albinotatus</i> Gravenhorst, 1829	B M S
<i>arenicola</i> Thomson, 1873	B S	<i>crassifemur</i> (Thomson, 1888)	*S
<i>armator</i> Fabricius, 1804	B M S	<i>funebris</i> Gravenhorst, 1829	S
<i>attentorius</i> (Panzer, 1804)	S	<i>grammicus</i> Gravenhorst, 1829	B
<i>dianae dianae</i> Gravenhorst, 1829	B M S	<i>notatus</i> Gravenhorst, 1829	B M S
<i>dianae obscuripes</i> Zetterstedt, 1840	S	<i>obnoxius</i> Gravenhorst, 1829	B M S
<i>inculcator</i> (Linnaeus, 1758)	B M S	<i>transfuga</i> Gravenhorst, 1829	B M S
<i>leucocheir</i> Ratzeburg, 1844	B M	<b>Listrognathus</b> Tschek, 1870	
<i>luctuosus subquadratus</i> Thomson, 1873	B	<b>Listrognathus</b> s. str.	
<i>lugubris lugubris</i> Gravenhorst, 1829	B		
<i>minator</i> (Gravenhorst, 1829)	B		
<i>moschator moschator</i> (Fabricius, 1787)	*B*M*S		

<i>compressicornis</i>			<b>Teleutea</b> Foerster, 1868
(Gravenhorst, 1829)	M		<i>brischkei</i> (Holmgren, 1860)
<i>pubescens</i> (Fonscolombe, 1850)	M S	B	<i>striata</i> (Gravenhorst, 1829)
<i>mactator</i> (Thunberg, 1822)	*S	B M*S	
<b>Stenarella</b> Szépligeti, 1916			<b>Apophua</b> Morley, 1913
<i>gladiator</i> (Scopoli, 1763)	B M S		<i>bipunctoria</i> (Thunberg, 1822)
<b>Nematopodius</b> Gravenhorst, 1829			<i>flavolineata</i> (Gravenhorst, 1829)
<b>Nematopodius</b> s. str.			<i>cubitoria</i> (Thunberg, 1822)
<i>formosus</i> Gravenhorst, 1829	B M S	B M S	<i>cicatricosa</i> (Ratzeburg, 1848)
<b>Acroricnus</b> Ratzeburg, 1852			<i>evanescens</i> (Ratzeburg, 1848)
<i>stylator stylator</i> (Thunberg, 1822)	M	B	<i>genalis</i> (Möller, 1883)
<b>Sphecophaga</b> Westwood, 1840		B M	<b>Glypta</b> Gravenhorst, 1829
<i>vesparum vesparum</i> (Curtis, 1828)	B M		<b>Glypta</b> s. str.
<i>thuringiaca</i> Schmiedeknecht, 1914			<i>arreptans</i> Hellén, 1915
<b>Latibulus</b> Gistel, 1848			<i>biauriculata</i> Strobl, 1901
<i>argiolus</i> (Rossi, 1790)	M S	S	<i>bifoveolata</i> Gravenhorst, 1829
<b>LYCORINAE</b>			M S
<b>Lycorina</b> Holmgren, 1859			<i>brevipetiolata</i> Thomson, 1889
<i>triangulifera</i> Holmgren, 1859	B M S	M	<i>crassitarsis</i> Thomson, 1889
<b>NEORHACODINAE</b>			<i>dentifera</i> Thomson, 1889
<b>Neorhacodes</b> Hedicke, 1922			<i>ephippigera</i>
<i>enslini</i> (Ruschka, 1922)	B M	S	Kriechbaumer, 1895
<b>STILBOPINAE</b>			<i>femorator</i> Desvignes, 1856
<b>Stilbops</b> Foerster, 1868		B M S	<i>filicornis</i> Thomson, 1889
<i>Aphanoroptrum</i> Foerster, 1868			<i>haesitator</i> Gravenhorst, 1829
<i>abdominalis</i> (Gravenhorst, 1829)	B M S	S	
<i>ruficornis</i> (Gravenhorst, 1829)			<i>heterocera</i> Thomson, 1889
<i>vetula</i> (Gravenhorst, 1829)	B M	M S	<i>incisa</i> Gravenhorst, 1829
<i>chrysostomus</i> (Gravenhorst, 1829)			<i>lineata</i> Desvignes, 1856
<b>BANCHINAE</b>			<i>longicauda</i> Hartig, 1838
Lissonotinae			<i>macropyga</i> Hellén, 1915
<b>GLYPTINI</b>			<i>mensurator</i> (Fabricius, 1775)
			B M S
			<i>microcera</i> Thomson, 1889
			B S
			<i>nigricornis</i> Thomson, 1889
			B
			<i>nigrina</i> Desvignes, 1856
			B M S
			<i>fractigena</i> Thomson, 1889
			M
			<i>nigripes</i> Strobl, 1901
			<i>parvicornuta</i> Bridgman, 1886
			M S
			<i>pedata</i> Desvignes, 1856
			B
			B M S

<i>rostrata</i> Holmgren, 1860	S	<i>bellator</i> Gravenhorst, 1829	B
<i>rufata</i> Bridgman, 1888	S	<i>biguttata</i> Holmgren, 1860	M S
<i>rufipes</i> Brischke, 1864	S	<i>bistrigata</i> Holmgren, 1860	B S
<i>rufoclypeata</i> Kiss, 1924	S	<i>bivittata</i> Gravenhorst, 1829	B
<i>salicis</i> Thomson, 1889	M	<i>breviventris</i> (Hellén, 1915)	B
<i>salsolicola</i> Schmiedeknecht, 1915	B M S	<i>buccator</i> (Thunberg, 1822)	B S
<i>scalaris</i> Gravenhorst, 1829	B	<i>buolianae</i> Hartig, 1838	B M S
<i>sculpturata</i> Gravenhorst, 1829	B M	<i>carbonaria</i> Holmgren, 1860	M S
<i>macrura</i> Habermehl, 1918		<i>catenator</i> (Panzer, 1804)	B M S
<i>scutellaris</i> Thomson, 1889	B M	<i>clypealis</i> Thomson, 1877	M
<i>similis</i> Bridgman, 1886	B S	<i>clypeator</i> (Gravenhorst, 1829)	B M S
<i>teres</i> Gravenhorst, 1829	B	<i>coxator</i> Smith v. Burgst, 1914	M
<i>tenuicornis</i> Thomson, 1889	M	<i>culiciformis</i> Gravenhorst, 1829	B M
<i>vulnerator</i> Gravenhorst, 1829	*M	<i>deversor</i> Gravenhorst, 1829	B M S
<b>Conoblasta</b> Foerster, 1868		<i>distincta</i> Bridgman, 1889	B
<i>caudata</i> (Thomson, 1889)	B S	<i>dubia</i> Holmgren, 1855	B M S
<i>ceratites</i> [Gravenhorst, 1829]	B M S	<i>errabunda</i> Holmgren, 1860	B M S
<i>consimilis</i> Holmgren, 1860	B S	<i>excelsa</i> Schmiedeknecht, 1900	M
<i>xanthognatha</i> Thomson, 1889		<i>fletcheri</i> Bridgman, 1882	M
<i>elongata</i> Holmgren, 1860	M	<i>freyi</i> (Hellén, 1915)	B
<i>extincta</i> Ratzeburg, 1852	B M S	<i>folii</i> Thomson, 1877	B M S
<i>fronticornis</i> [Gravenhorst, 1829]	M	<i>transversa</i> Bridgman, 1889	
<i>monoceros</i> Gravenhorst, 1829	S	<b>fundator</b> (Thunberg, 1822)	B M S
<b>Diblastomorpha</b> Foerster, 1868		<i>sulphurifera</i> Gravenhorst, 1829	
<i>cylindrator</i> Fabricius, 1787	B M	<i>gracilenta</i> Holmgren, 1860	M
<i>bicornis</i> Boie, 1850		<i>humarella</i> Thomson, 1877	B M S
<b>Cephaloglypta</b> Obertel, 1956		<i>impressor</i> Gravenhorst, 1829	B M
<i>murinanae</i> (Bauer, 1941)	M S	<i>basalis</i> Brischke, 1880	
<b>LISSONOTINI</b>		<i>impressifrons</i> Thomson, 1889	M S
<b>Lissonota</b> Gravenhorst, 1829		<i>inconspicua</i> Schmiedeknecht, 1935	M
<i>Lampronota</i> Curtis, 1832		<i>insignita</i> Gravenhorst, 1829	B M S
<i>Meniscus</i> Schioedte, 1839		<i>lineata</i> Gravenhorst, 1829	B S
<i>Pimplopterus</i> Ashmead, 1900		<i>maculata</i> Brischke, 1864	M
<i>Campocineta</i> Schmiedeknecht, 1900		<i>nigra</i> Brischke, 1880	B S
<i>agnata</i> Gravenhorst, 1829	B M	<i>nigricoxis</i> Habermehl, 1916	M
<i>antennalis</i> Thomson, 1877	B M	<i>nigridens</i> Thomson, 1889	B
<i>agricola</i> Gravenhorst, 1829	M	<i>nitida</i> Bridgman, 1886	B M

<i>palmalis</i> Thomson, 1889	*B		<i>caligata</i> (Gravenhorst, 1829)	M
<i>parallela</i> Gravenhorst, 1829	B M S		<i>errabunda</i> (Gravenhorst, 1829)	M
<i>picticoxis</i> Schmiedeknecht 1900	*M		<i>quadrilineata</i> (Gravenhorst, 1829)	M
<i>implator</i> (Zetterstedt 1840)	B M		<i>blandus</i> (Gravenhorst, 1829)	
<i>quadrinotata</i> Gravenhorst, 1829	S		<i>solitaria</i> Schmiedeknecht, 1900	*B
<i>rubricoxa</i> Brischke, 1880	*M			
<i>rufitarsis</i> Szépligeti, 1900	S			
<i>rusticator</i> (Thunberg, 1822)	B M S		<b>Syzeuctus</b> Foerster, 1868	
			<i>Diceratops</i> Foerster, 1868	
<i>segmentellator</i> Aubert, 1967			<i>apicalis</i> (Gravenhorst, 1829)	*S
<i>saturator</i> (Thunberg, 1822)	B	S	<i>bicornis</i> (Gravenhorst, 1829)	B M S
<i>setosa</i> (Geoffroy in Fourcroy 1785)	*B		<i>decoratus</i> (Costa, 1890)	*S
<i>similis</i> (Habermehl, 1917)	B M S		<i>irrisorius</i> (Rossi, 1794)	*B*M*S
<i>strigifrons</i> Schmiedeknecht, 1900	*M		<i>maculatorius</i> (Fabricius, 1787)	B S
<i>subaciculata</i> Bridgman, 1886	B		<i>tenuifasciatus</i>	
<i>tolli</i> (Heinrich, 1929)	*B		Schmiedeknecht, 1900	B M S
<i>unicincta</i> Holmgren, 1860	B M			
<i>vaga</i> Szépligeti, 1900	B M		BANCHINI	
<i>variabilis</i> Holmgren, 1860	B M S		<b>Exetastes</b> Gravenhorst, 1829	
			<i>adpressoarius</i> <i>adpressoarius</i>	
<i>trochanteralis</i> Dalla Torre, 1901			(Thunberg, 1822)	B M S
<i>versicolor</i> Holmgren, 1860	*S		<i>guttatorius</i> (Gravenhorst, 1829)	
<b>Alleplasta</b> Foerster, 1868			<i>atrator</i> (Foerster, 1771)	B M S
<i>grisea</i> (Gravenhorst, 1829)	*B		<i>cinctipes</i> (Retzius in Degeer, 1783)	
<i>piceator</i> (Thunberg, 1822)	B M S		<i>bicoloratus</i> Gravenhorst, 1829	M
<b>Arenetra</b> Holmgren, 1859			<i>braunsi</i> Dalla Torre, 1901	*B
<i>pilosella</i> (Gravenhorst, 1829)	B M S		<i>calobatus</i> Gravenhorst, 1829	B M
<b>Cryptopimpla</b>			<i>crassus</i> Gravenhorst, 1829	M S
<i>Taschenberg</i> , 1863			<i>femorator</i> (Desvignes, 1856)	M
<i>Xenakis</i> Foerster, 1868			<i>fornicator</i> <i>fornicator</i>	
<i>anomala</i> (Holmgren, 1860)	M		(Fabricius, 1781)	B M S
<i>arvicola</i> (Gravenhorst, 1829)	B M		<i>geniculosus</i> Holmgren, 1858	*B
			<i>gracilicornis</i> Gravenhorst, 1829	B *S
<i>brachycentra</i> (Gravenhorst, 1829)			<i>guttifer</i> (Thomson, 1897)	*B
<i>calceolata</i> (Gravenhorst, 1829)	B M S		<i>illusor</i> Gravenhorst, 1829	B
			<i>illyricus</i> Strobl, 1903	*B

<i>laevigator</i> (Villers, 1789)	B M S	<i>fulvipes</i> (Gravenhorst, 1829)	B M S
<i>nigripes</i> Gravenhorst, 1829	B M S	<i>mordax</i> (Thomson, 1883)	B M
<i>notatus</i> Holmgren, 1858	B M	<i>nigripes</i>	*M
<i>robustus</i> Gravenhorst, 1829	B M S	<i>Polycinetis</i> Foerster, 1868	
<i>rufipes</i> (Gmelin in Linnaeus, 1790)	*B	<i>fulvicornis</i> (Kriechbaumer, 1891)	B M
<b>Rhynchobanchus</b>		<i>resplendens</i> (Holmgren, 1855)	M S
Kriechbaumer, 1894		<b>Notopygus</b> Holmgren, 1855	
<i>bicolor</i> Kriechbaumer 1894	B S	<i>emarginatus</i> Holmgren, 1855	B M
<b>Banchus</b> Fabricius, 1798		<i>flavicornis</i> Holmgren, 1855	M
<i>dilatatorius</i> (Thunberg, 1822)	M	<i>megerlei</i> (Tschech, 1868)	B M S
<i>compressus</i> (Fabricius, 1787)		<i>nigricornis</i> Kriechbaumer, 1891	B M S
<i>falcatorius</i>		<b>Homaspis</b> Foerster, 1868	
Fabricius, 1775)	B M S	<i>Nehomaspis</i> Heinrich, 1949	
<i>hastator</i> (Fabricius, 1793)	B S	<i>alpina</i> (Heinrich, 1949)	*B*M
<i>femoralis</i> Thomson, 1897		<i>narrator</i> (Gravenhorst, 1829)	B M
<i>pictus</i> Fabricius, 1798	B M S	<i>robustus</i> (Thomson, 1894)	*B M
<i>volutatorius</i> (Linnaeus, 1758)	M	<i>rufinus</i> (Gravenhorst, 1829)	B M S
<i>monileatus</i> (Gravenhorst, 1829)		<i>subalpina</i> Schmiedeknecht, 1913	B M
<b>CTENOPELMATINAE</b>		<i>varicolor</i> (Thomson, 1894)	*M
Mesoleiinae		<b>OLETHRODOTINI</b>	
Scolobatinae		<b>Olethrodotis</b> Foerster, 1868	
<b>CTENOPELMATINI</b>		<i>Taschenbergia</i> Schmiedeknecht, 1888	
<b>Ctenopelma</b> Holmgren, 1855		<i>modestus</i> (Gravenhorst, 1829)	*M
<i>boreale</i> Holmgren, 1855	M	<b>PIONINI</b>	
<i>funebris</i> (Habermehl, 1923)	B M	<b>Lethades</b> Davis, 1897	
<i>lucifer</i> (Gravenhorst, 1829)	M	<i>facialis</i> (Brischke, 1871)	B
<i>luteum</i> Holmgren, 1855	*M	<i>laricis</i> Hinz 1976	M
<i>gagatum</i> (Kriechbaumer, 1894)		<i>scabriculus</i> (Thomson, 1883)	B M
<i>nigripenne</i> (Gravenhorst, 1829)	M	<b>Trematopygus</b> Holmgren, 1855	
<i>nigrum</i> Holmgren, 1855	B M	<i>dictator</i> (Thunberg, 1822)	S
<i>sericans</i> Tschech, 1868		<i>dubitator</i> Hinz, 1982	B
<i>xanthostigmum</i> Holmgren, 1855	B M	<i>melanocerus</i> (Gravenhorst, 1829)	M
<b>Xenoschesis</b> Foerster, 1868			
<i>Xenoschesis</i> s. str.			

<i>ruficornis</i> Holmgren, 1855	M S	<b>Lathiponus</b> Foerster, 1868
<i>vellicans</i> (Gravenhorst, 1829)	*M	<i>Ceratosoatis</i> Gregor, 1939
<b>Glyptorhaestus</b> Thomson, 1894		<i>frigidus</i> (Woldstedt, 1872) M
<i>punctatus</i> (Thomson, 1890)	M	<i>ornatus</i> (Gregor, 1939)
<b>Rhorus</b> Foerster, 1868		<b>Synoectes</b> Foerster, 1868
<i>angulatus</i> Thomson, 1888	M	<i>tenuicornis</i> (Gravenhorst, 1829) *B*M
<i>chrysopygus</i> Roman, 1909	M	
<i>extirpatorius</i> (Gravenhorst, 1829)	M	<b>Zaplethocornia</b>
<i>lapponicus</i> Roman, 1909	*M	Schmiedeknecht, 1912
<i>longicornis</i> (Holmgren, 1856)	M	<i>procurator</i> (Gravenhorst, 1820) M
<i>mesoxanthus</i> (Gravenhorst, 1829)	M	<b>Oetophorus</b> Foerster, 1868
<i>neustriæ</i> (Schrank, 1802)	M	<i>naevius</i> (Gmelin in Linnaeus, 1790) B M S
<i>palustris</i> (Holmgren, 1855)	*B	<b>Perilissus</b> Holmgren, 1855
<i>substitutor</i> (Thunberg, 1822)	*M	<i>abdominalis</i> Brischke, 1878 *S
<b>Syntactus</b> Foerster, 1868		<i>erythrocephalus</i> (Gravenhorst, 1829) B M S
<i>delusor</i> (Linnaeus, 1758)	M	<i>filicornis</i> (Gravenhorst, 1820) B M S
<i>varius</i> (Holmgren, 1855)	S	<i>longicornis</i> Brischke, 1871 *M
<b>Pion</b> Schioedte, 1838		<i>luteolator</i> (Gravenhorst, 1829) *B M
<i>Catoglyptus</i> Holmgren, 1855		<i>lutescens</i> Holmgren, 1855 *M
<i>crassipes</i> (Holmgren, 1855)	*B	<i>moravicus</i> Habermehl, 1923 M
<i>fortipes</i> (Gravenhorst, 1829)	B M S	<i>orbitalis</i> (Gravenhorst, 1829) M
<b>Rhaestus</b> Thomson, 1883		<i>pallidus</i> (Gravenhorst, 1829) B M
<i>rufipes</i> (Holmgren, 1855)	*M	<i>pictilis</i> Holmgren, 1855 *S
<b>Symphera</b> Foerster, 1868		<i>rufoniger rufoniger</i> Gravenhorst, 1820) M S
<i>Stiphrosomus</i> Foerster, 1868		<i>sericeus</i> (Gravenhorst, 1829) M
<i>ambulator</i> (Thunberg, 1822)	M*S	<i>spilonotus</i> (Stephens, 1835) *B
<i>antilope</i> (Gravenhorst, 1829)	*M	<i>thuringiacus</i> Schmiedeknecht, 1912
<i>fuscicornis</i> (Gmelin in Linnaeus, 1790)	B M S	<i>subcinctus</i> (Holmgren, 1855)
<i>montana</i> (Gravenhorst, 1829)	M	<b>Absyrtus</b> Holmgren, 1859
<i>ullrichi</i> (Tschek, 1868)	*M	<i>vicinator</i> (Thunberg, 1822) B M S
PERILISSINI		<i>luteus</i> Holmgren, 1859
		<b>Lophyroplectus</b> Thomson, 1883

<i>luteator</i> (Thunberg, 1822)	B M S		
<b>Opheltes</b> Holmgren, 1859			
<i>glaucopterus glaucopterus</i> (Linnaeus, 1758)	B M S		
<b>Priopoda</b> Holmgren, 1855			
<i>stictica</i> (Fabricius, 1798)	*B M*S		
<b>Lathrolestes</b> Foerster, 1868			
<i>ensator</i> (Brauns, 1898)	*B		
<i>luteolus</i> Thomson, 1883	B		
<i>marginatus</i> (Thomson, 1883)	*M		
SCOLOBATINI			
<b>Scolobates</b> Gravenhorst, 1829			
<i>auriculatus</i> (Fabricius, 1804)	B M S		
<i>hylotomae</i> Kriechbaumer, 1877			
MESOLEIINI			
<b>Otlophorus</b> Foerster, 1868			
<i>hypomelas</i> (Thomson, 1894)	M		
<i>italicus</i> (Gravenhorst, 1829)	*S		
<i>pulverulentus</i> (Holmgren, 1855)	M		
<i>verpetorum</i> (Gravenhorst, 1829)	B M		
<b>Azelus</b> Foerster, 1868			
<i>erythropalpus</i> (Gmelin in Linnaeus, 1790)	M		
<b>Lagarotis</b> Foerster, 1868			
<i>Lagarotus</i> Thomson, 1892			
<i>debitor</i> (Thunberg, 1822)	*M		
<i>erythrocerus</i> (Gravenhorst, 1829)	M		
<i>pubescens</i> (Holmgren, 1855)	M		
<i>semicaligatus</i> (Gravenhorst, 1820)	B M		
<b>Barytarbes</b> Foerster, 1868			
<i>colon</i> (Gravenhorst, 1829)	M		
		<i>hilarellus</i>	
		Schmiedeknecht, 1914	M
		<i>laeviusculus</i> (Thomson, 1883)	*B
		<i>superbus</i> Schmiedeknecht, 1914	*M
		<i>ubusinensis</i> Gregor, 1937	M
		<b>Alexeter</b> Foerster, 1868	
		<i>albilabris</i> (Thomson, 1894)	*M
		<i>fallax</i> (Holmgren, 1855)	M
		<i>gracilentus</i> (Holmgren, 1855)	*M
		<i>multicolor</i> (Gravenhorst, 1829)	M*S
		<i>nebulator</i> (Thunberg, 1822)	B M S
		<i>niger</i> (Gravenhorst, 1829)	M
		<i>rapinator</i> (Gravenhorst, 1829)	M
		<i>sectator</i> (Thunberg, 1822)	B M
		<i>inconspicuus</i> Schmiedeknecht, 1914	
		<i>testaceator</i> (Thunberg, 1822)	B M
		<b>Protarchus</b> Foerster, 1868	
		<i>melanurus</i> (Thomson, 1894)	*B
		<i>sorbi</i> (Ratzeburg, 1844)	M
		<i>testatorius</i> (Thunberg, 1822)	B M
		<b>Perispuda</b> Foerster, 1868	
		<i>Genarches</i> Foerster, 1868	
		<i>facialis</i> (Gravenhorst, 1829)	M S
		<i>sulphurata</i> (Gravenhorst, 1807)	*M
		<b>Lamachus</b> Foerster, 1868	
		<i>coalitorius</i> (Thunberg, 1822)	B M S
		<i>marginatus</i> (Brischke, 1871)	
		<i>ophthalmicus</i> (Holmgren, 1855)	
		<i>consimilis</i> (Holmgren, 1855)	B M
		<i>eques</i> (Hartig, 1838)	B M S
		<i>frutetorum</i> (Hartig, 1838)	B M
		<i>lophirorum</i> (Hartig, 1838)	
		<i>pini</i> Bridgman, 1882	*B

<i>thomsoni</i> Fitton, 1982	*M		<i>liosternus</i> (Thomson, 1894)	*B*M
<i>longiventris</i> (Thomson, 1895)				
<i>transiens</i> (Ratzeburg, 1852)	B M S		<i>scapularis</i> (Stephens, 1835)	M
<i>intermedius</i> (Ratzeburg, 1848)			<i>trochanteratus</i> (Kriechbaumer, 1896)	M S
<i>virgultorum</i> (Gravenhorst, 1829)	*B		<i>viduus</i> (Holmgren, 1855)	M
<b>Scopesis</b> Foerster, 1868			<b>Mesoleius</b> Holmgren, 1855	
<i>bicolor</i> (Gravenhorst, 1829)	M		<i>armillatorius</i> (Gravenhorst, 1807)	B M
<i>depressa</i> (Thomson, 1894)	B M		<i>aulicus aulicus</i> (Gravenhorst, 1829)	M
<i>fraterna</i> (Holmgren, 1855)	*M		<i>dubius</i> Holmgren, 1855	B M
<i>frontator</i> (Thunberg, 1822)	M		<i>melanoleucus</i> (Gravenhorst, 1829)	B M
<i>gesticulator</i> (Thunberg, 1822)	B M		<i>opticus</i> (Gravenhorst, 1829)	*S
<i>macropa</i> (Thomson, 1894)	M		<i>pyriformis</i> (Ratzeburg, 1852)	M
<i>obscura</i> (Holmgren, 1855)	*M		<i>tibialis</i> (Holmgren, 1855)	*B
<i>politia</i> (Holmgren, 1855)	M		<i>variegatus</i> (Jurine, 1807)	M
<i>rufonotata</i> (Holmgren, 1876)	*M		<b>Hyperbatus</b> Foerster, 1868	
<i>rubrocincta</i> Schmiedeknecht, 1914	*M		<i>segmentator</i> (Holmgren, 1855)	*B
<i>tegularis</i> (Thomson, 1894)	M		<b>Saotis</b> Foerster, 1868	
<i>thomsoni</i> Habermehl, 1925	M		<i>bilineatus</i> (Gravenhorst, 1829)	M
<b>Himerta</b> Foerster, 1868			<i>brevispina</i> (Thomson, 1883)	*M
<i>defectiva</i> (Gravenhorst, 1820)	M		<i>compressiusculus</i> (Thomson, 1883)	*M
<b>Rhinotorus</b> Foerster, 1868			<i>liopleuris</i> (Thomson, 1888)	M
<i>Spudaea</i> Foerster, 1868			<b>Anoncus</b> Townes, 1970	
<i>compactor</i> (Thunberg, 1822)	*M		<i>femorator</i> (Thomson, 1894)	*M
<i>leucostomus</i> (Gravenhorst, 1829)	*M		<b>EURYPROCTINI</b>	
<b>Campodorus</b> Foerster, 1868			<b>Synomeiix</b> Foerster, 1868	
<i>astutus</i> (Holmgren, 1876)	*S		<i>Zemiophora</i> Foerster, 1868	
<i>caligatus</i> (Gravenhorst, 1829)	M		<i>scutulata</i> (Hartig, 1838)	B *S
<i>dorsalis</i> (Gravenhorst, 1829)	M		<b>Synodites</b> Foerster, 1868	
<i>haematodes</i> (Gravenhorst, 1829)	M		<i>carinatus</i> (Holmgren, 1855)	M
<i>hamulus</i> (Gravenhorst, 1829)	*M		<i>contractus</i> (Holmgren, 1855)	M
<i>ignavus</i> (Holmgren, 1855)	M		<i>lineiger</i> (Thomson, 1894)	M
<i>incidens</i> (Thomson, 1894)	M			

<i>notatus</i> (Gravenhorst, 1829)	M	<b>Hypsantyx</b> Pfankuch, 1906
<b>Pantorhaestes</b> Foerster, 1868		<i>lituratorius</i> (Linnaeus, 1761) S
<i>Thophaconus</i> Foerster, 1868		<b>Hypamblys</b> Foerster, 1868
<i>xanthostomus</i> (Gravenhorst, 1829)	M	<i>albopictus</i> (Gravenhorst, 1829) *M
<b>Mesoleptidea</b> Viereck, 1912		<i>slaviceki</i> Gregor, 1935 M
<i>bipunctata</i> (Gravenhorst, 1829)	B M S	<i>transfuga</i> (Holmgren, 1855) *B
<i>cinctula</i> (Gravenhorst, 1829)	B M S	<b>Phobetes</b> Foerster, 1868
<i>prosoleuca</i> (Gravenhorst, 1829)	*B M	<i>Ipoctonus</i> Foerster, 1868
<i>xanthostigma</i> (Gravenhorst, 1829)	B M*S	<i>Phobetus</i> Thomson, 1889
<b>Gunomeria</b> Schmiedeknecht, 1907		<i>atomator</i> (Müller, 1776) *M
<i>macrodactylus</i> (Holmgren, 1855)	M S	<i>chrysostomus</i> (Gravenhorst, 1829) M
<b>Anisotacrus</b> Schmiedeknecht, 1913		<i>femorator</i> (Thomson, 1894) M
<i>tenellus</i> (Holmgren, 1855)	*M	<i>leptocerus</i> (Gravenhorst, 1820) M
<b>Hadrodactylus</b> Foerster, 1868		<i>nigriceps</i> (Gravenhorst, 1829) *B*M
<i>confusus</i> (Holmgren, (1856)	*M	<b>Euryproctus</b> Holmgren, 1855
<i>fasciator</i> (Thunberg, 1822)	B M	<i>Sychnoleter</i> Foerster, 1868
<i>vulneratus</i> (Zetterstedt, 1838)		<i>alpinus</i> Holmgren, 1855 M
<i>femoralis</i> (Holmgren, 1855)	M	<i>annulatus</i> (Gravenhorst, 1829) *M
<i>flavifrontator</i> (Thunberg, 1822)	M	<i>crassicornis</i> Thomson, 1889 M
<i>genalis</i> Thomson, 1883	*M	<i>geniculatus</i> (Gravenhorst, 1829) *M
<i>gracilipes</i> Thomson, 1883	*B	<i>luteicornis</i> (Gravenhorst, 1829) M
<i>larvatus</i> Kriechbaumer, 1891	*S	<i>mundus</i> (Gravenhorst, 1829) *M
<i>tiphae</i> (Geoffroy in Fourcroy, 1785)	B M	<i>nemoralis</i> (Geoffroy in Fourcroy, 1785) M
<b>Syndipnus</b> Foerster, 1868		<i>parvulus</i> (Thomson, 1883) *B*M
<i>discolor</i> (Holmgren, 1855)	B M	<i>regenerator</i> (Fabricius, 1804) *M
<i>lateralis</i> (Gravenhorst, 1829)	M	<i>tuberculatus</i> Holmgren, 1855 M
<i>pectoralis</i> Thomson, 1894	*M	PHRUDINAE
		<b>Seleucus</b> Holmgren, 1858
		<i>cuneiformis</i> Holmgren, 1858 *B

<b>Earobia</b> Townes, 1951				
<i>Echthrolaricobius</i> Perkins, 1953				
<i>paradoxa</i> (Perkins, 1953)	S			
<b>Phrudus</b> Foerster, 1868				
<i>Mengersenia</i>				
Schmiedeknecht, 1907				
<i>monilicornis</i> (Bridgman, 1886)	*B			
CAMPOPLEGINAE				
Parizontinae				
HELLWIGIINI				
<b>Hellwigia</b> Gravenhorst, 1823				
<i>elegans</i> Gravenhorst, 1823	M S			
CAMPOPLEGINI				
<b>Sinophorus</b> Foerster, 1868				
<i>Eulimneria</i> Schmiedeknecht, 1907				
<i>albidus</i> (Gmelin in Linneaus, 1790)	B			
<i>crassifemur</i> (Thomson, 1887)	B*M S			
<i>geniculatus</i> (Gravenhorst, 1829)	B M			
<i>fuscicarpus</i> (Thomson, 1887)	S			
<i>rufifemur</i> (Thomson, 1887)	B			
<i>xanthostomus</i> (Gravenhorst, 1829)	B *S			
<b>Campoplex</b> Gravenhorst, 1829				
<i>Omorgus</i> Foerster, 1868				
<i>abbreviatus</i> (Brischke, 1880)	M S			
<i>atramentarius</i> (Schmiedeknecht, 1909)	S			
<i>angulatus</i> (Thomson, 1887)	*S			
<i>bilobus</i> (Thomson, 1887)	M			
<i>borealis</i> (Zetterstedt, 1838)	B M S			
<i>caudatus</i> Szépligeti, 1900	S			
<i>coracinus</i> (Thomson, 1887)	M			
<i>cursitans</i> (Holmgren, 1860)	*M S			
			<i>difformis</i> (Gmelin in Linnaeus, 1790)	B S
			<i>ensator</i> Gravenhorst, 1829	*M
			<i>excavatus</i> (Brischke, 1880)	S
			<i>faunus</i> Gravenhorst, 1829	B M S
			<i>fusciplica</i> (Thomson, 1887)	*M S
			<i>gastroides</i> Gravenhorst, 1829	*M S
			<i>gracilis</i> Ulbricht, 1910	S
			<i>holmgreni</i> (Tschech, 1871)	*B
			<i>liogaster</i> (Thomson, 1887)	M S
			<i>lugubrinus</i> (Holmgren, 1855)	M S
			<i>lyratus</i> (Thomson, 1887)	S
			<i>maculipes</i> (Tschech, 1874)	*B
			<i>melanostictus</i> Gravenhorst, 1829	B M
			<i>mutabilis</i> (Holmgren, 1860)	M S
			<i>nigricanae</i> Horstmann, 1980	S
			<i>ornatus</i> (Gregor, 1940)	M
			<i>ovatus</i> (Brischke, 1880)	B M S
			<i>ramidulus</i> (Brischke, 1880)	M
			<i>restrictor</i> Aubert, 1960	*S
			<i>rothii</i> (Holmgren, 1855)	B M S
			<i>rufinator</i> Aubert, 1969	*M*S
			<i>ruficoxa</i> (Thomson, 1887)	B M S
			<i>rufipes</i> (Bridgman, 1883)	
			<i>scaposus</i> (Thomson, 1887)	S
			<i>solitarius</i> (Brischke, 1880)	B
			<i>tricoloripes</i> (Schmiedeknecht, 1909)	B M S
			<i>tumidulus</i> Gravenhorst, 1829	S
			<b>Lathroplex</b> Foerster, 1868	
			<i>caudatus</i> (Gregor, 1941)	M
			<i>clypearis</i> Thomson, 1887	B
			<b>Sesioplex</b> Viereck, 1912	
			<i>cerophagus</i> (Gravenhorst, 1829)	B M
			<i>picticrus</i> (Thomson, 1887)	

<b>Venturia</b> Schrottky, 1902		LIMNERIINI
<i>Devorgilla</i> Cameron, 1907		<i>Macrini</i>
<i>Idechthis</i> Foerster, 1868		
<i>canescens</i> (Gravenhorst, 1829)	B M S	
<i>investigator</i> (Habermehl, 1922)	M	<b>Rhimphoctona</b> Foerster, 1868
<i>moderator</i> (Linnaeus, 1758)	B M	<i>lucida</i> (Clément, 1924) M
<i>transfuga</i> (Gravenhorst, 1829)	*B	<i>megacephala</i> (Gravenhorst, 1829) B M
<i>siringellae</i> (Hedwig, 1944)		<i>austriacus</i> Tschek, 1871
<b>Casinaria</b> Holmgren, 1859		<i>hungaricus</i> Kiss, 1926
<i>Horogenes</i> Foerster, 1868		<i>melanura</i> (Holmgren, 1860) *B
<i>Alcima</i> Foerster, 1868		<i>obscuripes</i> (Holmgren, 1860) B M
<i>Phaedroctonus</i> Foerster, 1868		<i>rufocoxalis</i> (Clément, 1924) B *S
<i>affinis</i> Tschek, 1871	*M	<i>xoridiformis</i> (Holmgren, 1860) B M
<i>albipalpis</i> (Gravenhorst, 1829)	M	<i>xoridoidea</i> (Strobl, 1904) B M S
<i>claviventris</i> Holmgren, 1858	B M	<b>Nemeritis</b> Holmgren, 1860
<i>ischiofaster</i> Thomson, 1887	M S	<i>caudatula</i> Thomson, 1887 M
<i>mesozosta</i> (Gravenhorst, 1829)	B	<i>raphidiae</i> Krichbaumer, 1892
<i>moesta</i> (Gravenhorst, 1829)	S	<i>elegans</i> Szépligeti, 1916 M
<i>monticola</i> Thomson, 1887	*M	<i>batai</i> (Gregor, 1941)
<i>morionella</i> Holmgren, 1860	M	<i>graeca</i> Horstmann, 1975 ?
<i>nigripes</i> (Gravenhorst, 1829)	B M S	<i>lissonotoides</i> (Schmiedeknecht, 1909) *M
<i>orbitalis</i> (Gravenhorst, 1829)	B S	<i>macrocentra</i> (Gravenhorst, 1829) M S
<i>protensa</i> Thomson, 1887	M	<i>sordidus</i> (Gravenhorst, 1829)
<i>punctiventris</i> Woldstedt, 1876	S	<b>Bathyplectes</b> Foerster, 1868
<i>rufimanus</i> (Gravenhorst, 1829)	B M S	<i>Canidia</i> Holmgren, 1860
<i>scabra</i> (Thomson, 1887)	B M	<i>curculionis</i> (Thomson, 1887) *B M
<i>scutellaris</i> Tschek, 1871	M	<i>balteatus</i> (Thomson, 1887) *B
<i>stygia</i> Tschek, 1871	B M S	<i>rostratus</i> (Thomson, 1887) *B
<i>tenuiventris</i> (Gravenhorst, 1829)	M S	<i>stenostigma</i> (Thomson, 1887) B M
<i>varians</i> Tschek, 1871	B *S	<b>Callidora</b> (Foerster, 1868)
<i>vidua</i> (Gravenhorst, 1829)	M S	<i>albovincta</i> Holmgren, 1858) *B
<b>Charops</b> Holmgren, 1859		<i>annellata</i> Thomson, 1887
<i>cantator</i> (Degeer, 1778)	B M S	<b>Biolysia</b> Schmiedeknecht, 1907
<i>decipiens</i> (Gravenhorst, 1829)		<i>tristis</i> (Gravenhorst, 1829) B

<i>trochantella</i> (Thomson, 1887)			
<b>Nepiesta</b> Foerster, 1868			
<i>mandibularis</i> Holmgren, 1860	B M S		
<i>subclavata</i> Thomson, 1887	B M		
<b>Eripternus</b> Foerster, 1868			
<i>tarsalis</i> Szépligeti, 1905	*B	S	
<b>Leptocampoplex</b> Horstmann, 1970			
<i>cremastooides</i> (Holmgren, 1858)	B M S		
<b>Gonoty whole</b> Foerster, 1868			
<i>melanostoma</i> (Thomson, 1887)	B M		
<b>Pyraemon</b> Holmgren, 1859			
<i>Synetaeris</i> Foerster, 1868			
<i>carbonellum</i> Thomson, 1887	*B		
<i>fumipennis</i> (Zetterstedt, 1838)	S		
<i>truncicola</i> Thomson, 1887	M		
<b>Campoletis</b> Foerster, 1868			
<i>Sagaritis</i> Holmgren, 1859			
<i>Anilastus</i> Foerster, 1868			
<i>agilis</i> (Holmgren, 1860)	S		
<i>annulatus</i> (Gravenhorst, 1829)	M S		
<i>brachycera</i> (Thomson, 1887)	*M		
<i>cognatus</i> (Tschek, 1871)	M*		
<i>congestus</i> (Holmgren, 1858)	M		
<i>crassicornis</i> (Tschek, 1871)	M		
<i>dilatator</i> (Thunberg, 1822)	*M		
<i>disclusa</i> (Foerster, 1868)	M		
<i>erythropus</i> (Thomson, 1887)	M		
<i>latrator</i> (Gravenhorst, 1829)	M		
<i>maculipes</i> (Tschek, 1871)	M		
<i>posticus</i> (Bridgman et Fitch, 1885)	M		
<i>punctatus</i> (Bridgman, 1886)	*M		
<i>raptor</i> (Zetterstedt, 1838)	M		
<i>varians</i> (Thomson, 1887)	*S		
<i>vimmeri</i> (Gregor, 1937)	M		
<i>zonatus</i> (Gravenhorst, 1829)	M S		
<b>Delopia</b> Cameron, 1903			
<i>Dusona</i> Cameron, 1900			
<i>aemula</i> (Foerster, 1868)	*M		
<i>americana</i> (Ashmead, 1890)	B M		
<i>mariae</i> (Dalla Torre, 1901)			
<i>anceps</i> (Holmgren, 1860)	B M *S		
<i>angustata</i> (Thomson, 1887)	*S		
<i>alticola</i> (Gravenhorst, 1829)	B M		
<i>angustifrons</i> (Foerster, 1868)	M		
<i>aurita</i> (Kriechbaumer, 1883)	*M		
<i>bifida</i> (Thomson, 1887)	*B		
<i>bucculenta</i> (Holmgren, 1860)	B M S		
<i>carinifrons</i> (Holmgren, 1860)	B M		
<i>castanipes</i> (Thomson, 1887)	*B		
<i>circumcincta</i> (Foerster, 1868)	*B *M		
<i>confusa</i> (Foerster, 1868)	*B M		
<i>contumax</i> (Foerster, 1868)	B *S		
<i>crassipes</i> (Thomson, 1887)	*M		
<i>cultrator</i> (Gravenhorst, 1829)	B M S		
<i>delusor</i> (Thunberg, 1822)	*B		
<i>disclusa</i> (Foerster, 1868)	*B *S		
<i>erythrogaster</i> (Foerster, 1868)	B M		
<i>falcator</i> (Foerster, 1775)	B M		
<i>foersteri</i> (Roman, 1942)	B M *S		
<i>genalis</i> (Thomson, 1887)	B		
<i>habermehli</i> (Kriechbaumer, 1898)	M		
<i>heterocera</i> (Foerster, 1868)	B *S		
<i>humilis</i> (Foerster, 1868)	*M		
<i>inermis</i> (Foerster, 1868)	*B *M		
<i>infesta</i> (Foerster, 1868)	B M		

<i>insignita</i> (Foerster, 1868)	*M*S		
<i>lapponica</i> (Holmgren, 1860)	*S		
<i>leptogaster</i> (Holmgren, 1860)	B S		
<i>libertatis</i> (Teunissen, 1947)	*B		
<i>limnobia</i> (Thomson, 1887)	*B*M		
<i>myrtilla</i> (Desvignes, 1865)	*B		
<i>nidulator</i> (Fabricius, 1804)	B M		
<i>obliterata</i> (Holmgren, 1872)	B S		
<i>opaca</i> (Thomson, 1887)	B		
<i>oxyacanthae</i> (Boie, 1855)	B M S		
<i>perditor</i> (Foerster, 1868)	B S		
<i>peregrina</i> (Wollaston, 1858)	B M S		
<i>petiolator</i> (Fabricius, 1804)	*S		
<i>pineticola</i> (Holmgren, 1872)	S		
<i>polita</i> (Foerster, 1868)	B M*S		
<i>splendens</i> (Thomson, 1887)			
<i>pugillator</i> (Linnaeus, 1758)	*B M		
<i>pulchripes</i> (Holmgren, 1860)	*B		
<i>pulmentariae</i> (Hinz, 1963)	B S		
<i>recta</i> (Thomson, 1887)	*B*M		
<i>remota</i> (Foerster, 1868)	B M		
<i>rugifer</i> (Foerster, 1868)	*S		
<i>rugulosa</i> (Foerster, 1868)	*B		
<i>signator</i> (Brauns, 1895)	M		
<i>sobolicida</i> (Foerster, 1868)	*B		
<i>spinipes</i> (Thomson, 1887)	B S		
<i>stragifer</i> (Foerster, 1868)	*B *S		
<i>stygia</i> (Foerster, 1868)	*B*M		
<i>subcincta</i> (Foerster, 1868)	B M S		
<i>subimpressa</i> (Foerster, 1868)	B M S		
<i>tenuis</i> (Foerster, 1868)	B M S		
<i>terebrator</i> (Foerster, 1868)	*B*M		
		<i>ucrainica</i> (Hinz, 1972)	S
		<i>victor</i> (Thunberg, 1822)	*B M
		<i>vigilator</i> (Foerster, 1868)	*B
		<i>xenocampta</i> (Foerster, 1868)	*B M*S
		<i>zonella</i> (Foerster, 1868)	M
		<b>Pseudocymodusa</b> Habermahl, 1922	
		<i>alternans</i> (Gravenhorst, 1829)	*B*M
		<b>Meloboris</b> Holmgren, 1859	
		<i>Nepiera</i> Foerster, 1868	
		<i>collector</i> (Thunberg, 1822)	B M
		<i>concinna</i> (Holmgren, 1860)	
		<i>grisescens</i> (Gravenhorst, 1829)	*S
		<i>litoralis</i> (Holmgren, 1860)	*B
		<b>Cymodusa</b> Holmgren, 1859	
		<i>antennator</i> (Holmgren, 1855)	B M
		<i>cruentata</i> (Gravenhorst, 1829)	*M
		<i>flavipes</i> Brischke, 1880	*M
		<i>leucocera</i> Holmgren, 1859	*B M
		<i>petulans</i> Holmgren, 1859	*M
		<b>Dolopron</b> Foerster, 1868	
		<i>pedellus</i> (Holmgren, 1860)	B M
		<b>Phobocampe</b> Foerster, 1868	
		<i>bicingulata</i> (Gravenhorst, 1829)	S
		<i>confusa</i> (Thomson, 1887)	B M
		<i>crassiuscula</i> (Gravenhorst, 1829)	B M S
		<i>disparis</i> Gupta, 1983	B M S
		<i>unicincta</i> (Gravenhorst, 1829)	*B M*S
		<b>Tranosema</b> Foerster, 1868	
		<i>arenicola</i> Thomson, 1887	B M
		<i>rostralis</i> (Brischke, 1880)	
		<i>exoleta</i> (Thomson, 1887)	*B
		<i>nigridens</i> (Thomson, 1887)	*S
		<b>Macrus</b> Gravenhorst, 1829	

<i>parvulus</i> (Gravenhorst, 1829)	*S	<i>tenuipes</i> (Thomson, 1887)	S
<b>Enythus</b> Cameron, 1905		<i>trochanterata</i> (Thomson, 1887)	M S
<i>Dioces</i> Foerster, 1868		<i>truncata</i> (Thomson, 1887)	S
<i>apostatus</i> (Gravenhorst, 1829)	*B	<i>varians</i> (Brischke, 1880)	M
<i>exareolatus</i> (Ratzeburg, 1852)		<b>Hyposoter</b> Foerster, 1868	
<b>Tranosemella</b> Horstmann, 1977		<i>carbonarius</i> (Ratzeburg, 1844)	B M S
<i>praerogator</i> Linnaeus, 1758	*B *S	<i>coxalis</i> (Brischke, 1880)	M
<i>interrupta</i> (Holmgren, 1860)		<i>didymator</i> (Thunberg, 1822)	*B *S
<i>lacticrus</i> (Thomson, 1887)		<i>ruficinctus</i> (Gravenhorst, 1829)	
<b>Diadegma</b> Foerster, 1868		<i>ebeninus</i> (Gravenhorst, 1829)	B M S
<i>Angitia</i> Holmgren, 1859		<i>vulgaris</i> (Tschech, 1871)	
<i>agilis</i> (Brischke, 1880)	B	<i>neglectus</i> (Holmgren, 1858)	M
<i>apostata</i> (Gravenhorst, 1829)	S	<i>varicoxa</i> (Thomson, 1887)	
<i>armillata</i> (Gravenhorst, 1829)	B M S	<i>notatus</i> (Gravenhorst, 1829)	*M
<i>tibialis</i> (Gravenhorst, 1829)		<i>orbator</i> (Gravenhorst, 1829)	*B*M
<i>chrysostictos</i> (Gmelin in Linnaeus, 1790)	B M S	<i>pectinatus</i> (Thomson, 1887)	B
<i>crataegellae</i> (Thomson, 1887)	*B	<i>tenuicosta</i> (Thomson, 1887)	*B
<i>eucerophaga</i> Horstmann, 1969	B M S	<i>virginalis</i> (Gravenhorst, 1829)	B
<i>fenestralis</i> (Holmgren, 1860)	B M S	<b>Olesicampe</b> Foerster, 1868	
<i>holopyga</i> (Thomson, 1887)	*S	<i>Limneria</i> Holmgren, 1859	
<i>incompleta</i> Horstmann, 1973	S	<i>Holocremnus</i> Foerster, 1868	
<i>insectator</i> (Schrank, 1781)	S	<i>auctor</i> (Gravenhorst, 1829)	B
<i>laticrus</i> (Thomson, 1887)	S	<i>buccata</i> (Thomson, 1887)	*B*M
<i>latungula</i> (Thomson, 1887)	B S	<i>clandestina</i> (Holmgren, 1860)	B
<i>major</i> (Szépligeti, 1916)	S	<i>frutetorum</i> (Thomson, 1887)	S
<i>melia</i> (Thomson, 1887)	M	<i>fulcrans</i> (Thomson, 1887)	*B*M
<i>monospila</i> (Thomson, 1887)	S	<i>fulviventris</i> (Gmelin in Linnaeus, 1790)	M
<i>nana</i> (Gravenhorst, 1829)	*S	<i>genuicincta</i> (Hedwig, 1932)	M
<i>neocerophaga</i> Horstmann, 1969	*B*M*S	<i>macellator</i> (Thunberg, 1822)	M S
<i>parvicauda</i> (Thomson, 1887)	S	<i>melanogaster</i> (Thomson, 1887)	*S
<i>pusio</i> (Holmgren, 1860)	*B	<i>monticola</i> (Hedwig, 1938)	B M
<i>specularis</i> (Thomson, 1887)	*B		

<i>patellana</i> (Thomson, 1887)	*M		<b>Temelucha</b> Foerster, 1868
<i>ratzeburgi</i> (Tschek, 1871)	B		<i>albipennis</i> (Zetterstedt, 1838)
<i>signata</i> (Brischke, 1880)	*B		<i>annulata</i> (Szépligeti, 1900)
<i>sordidella</i> (Holmgren, 1858)	B M		M
<i>sternella</i> (Thomson, 1887)	M		<i>arenosa</i> (Szépligeti, 1900)
<i>vexata</i> (Holmgren, 1860)	M		B M S
<b>Lathrostizus</b> Foerster, 1868			<i>buoliania</i> (Curtis, 1854)
<i>lugens</i> (Gravenhorst, 1829)	*B		B M S
<i>macrostoma</i> (Thomson, 1887)		*S	<i>caudata</i> (Szépligeti, 1900)
<i>sternocerus</i> (Thomson, 1887)	*B		<i>confluens</i> (Gravenhorst, 1829)
<b>Lemophagus</b> Townes, 1965			B M S
<i>curtus</i> Townes, 1965	*B		<i>decorata</i> (Gravenhorst, 1829)
CREMASTINAE			S
<b>Dimophora</b> Foerster, 1868			<i>genalis</i> (Szépligeti, 1900)
<i>evanialis</i> (Gravenhorst, 1829)	B M		<i>interruptor</i> (Gravenhorst, 1829)
<i>robusta</i> Brischke, 1880	B M		M
<b>Pristomerus</b> Curtis, 1836			<i>lucida</i> (Szépligeti, 1900)
<i>armatus</i> Lucas, 1849	B M S		B M
<i>genalis</i> Sawoniewicz, 1978		*S	<i>mohelnensis</i> Šedivý, 1970
<i>horribilis</i> Narolsky, 1987	*M		<i>ophthalmica</i> (Holmgren, 1858)
<i>rivalis</i> Narolsky, 1987	*B		B M S
<i>vulnerator</i> (Panzer, 1799)	B M S		<i>signata</i> (Holmgren, 1858)
<b>Cremastus</b> Gravenhorst, 1829			B M
<i>bellicosus</i> Gravenhorst, 1829	B M S		<i>schoenobius</i> (Thomson, 1890)
<i>crassicornis</i> Thomson, 1890			M S
<i>geminus</i> Gravenhorst, 1829	B M S		<i>variipes</i> (Szépligeti, 1900)
<i>infirmus</i> Gravenhorst, 1829			B S
<i>kratochvili</i> Šedivý, 1970	B M		TERSILOCHINAE
<i>puberulus</i> Szépligeti, 1900	M		
<i>pungens</i> Gravenhorst, 1829			<b>Phradis</b> Foerster, 1868
<i>renominatus</i> Šedivý, 1970	S		<i>Isurgus</i> Foerster, 1868
<i>spectator</i> Gravenhorst, 1829	B M S		<i>brevicornis</i> Horstmann, 1971

<i>rufiventris</i> Horstmann, 1971	B	<i>flavipes</i> (Szépligeti, 1899)	B M
<b>Diaparsis</b> Foerster, 1868		<i>erythrostromus</i> (Gravenhorst, 1829)	M
<i>Diaparsis</i> s. str.		<b>Euporizon</b> Horstmann, 1971	
<i>carinifer</i> (Thomson, 1889)	B	<i>brevicornis</i> Horstmann, 1981	B M
<i>cingulata</i> Horstmann, 1971	M	<i>clavicornis</i> Horstmann, 1971	M
<i>jucunda</i> (Holmgren, 1860)	M	<i>corsicator</i> (Aubert, 1960)	M
<i>multiplicator</i> Aubert, 1969	M	<i>exilis</i> (Holmgren, 1860)	B M
<i>truncata</i> (Gravenhorst, 1829)	M	<i>gilvipes</i> (Gravenhorst, 1829)	B S
<b>Nanodiaparsis</b> Horstmann, 1971		<i>montanus</i> Horstmann, 1971	B M
<i>aperta</i> (Thomson, 1884)	B M	<i>nigriventris</i> Horstmann, 1971	M
<i>frontellus</i> (Holmgren, 1860)	B M	<i>rufipes</i> (Holmgren, 1850)	B M
<b>Sathropterus</b> Foerster, 1868		<i>truncorum</i> (Holmgren, 1860)	S
<i>pumilus</i> (Holmgren, 1858)	M S	<i>xanthopodus</i> (Holmgren, 1860)	B
<b>Aneuclis</b> Foerster, 1868		<b>Rugodiaparsis</b> Horstmann, 1971	
<i>brevicauda</i> (Thomson, 1889)	B M	<i>crassipes</i> (Thomson, 1889)	M
<i>incidens</i> (Thomson, 1889)	B	<i>ruficornis</i> (Szépligeti, 1899)	B
<i>melanarius</i> (Holmgren, 1860)	B	<b>Microdiaparsis</b> Horstmann, 1971	
<b>Epistathmus</b> Foerster, 1868		<i>microcephalus</i> (Gravenhorst, 1829)	B M
<i>crassicornis</i> Horstmann, 1971	B M	<i>neoversutus</i> (Horstmann, 1971)	B M S
<b>Barycnemis</b> Foerster, 1868		<i>versutus</i> (Holmgren, 1860)	B
<i>Leptopygus</i> Foerster, 1868		<b>Allophrooides</b> Horstmann, 1971	
<i>Cratophion</i> Thomson, 1889		<i>boops</i> (Gravenhorst, 1829)	M
<i>Cryptophion</i> Thomson, 1889		<b>Tersilochus</b> Holmgren, 1858	
<i>agilis</i> (Holmgren, 1858)	B M	<i>Tersilochus</i> s. str.	
<i>angustipennis</i> (Holmgren, 1858)	M	<i>clypeatus</i> Horstmann, 1971	M
<i>claviventris</i> (Gravenhost, 1829)	B S	<i>fulvipes</i> (Gravenhorst, 1829)	B M
<i>gracillimus</i> (Thomson, 1889)	S	<i>heterocerus</i> Thomson, 1889	B M
<i>gravipes</i> (Gravenhorst, 1829)	B M S	<i>jocator</i> (Holmgren, 1858)	B M
<i>filicornis</i> (Thomson, 1889)	B M	<i>nitidipleuris</i> Horstmann, 1971	M
<i>harpurus</i> (Schrank, 1802)	B M S	<i>obliquus</i> (Thomson, 1889)	B S
<i>laeviceps</i> (Thomson, 1889)	M	<i>obscurator</i> Aubert, 1959	B M
<i>punctifrons</i> Horstmann, 1981	S		
<b>Probles</b> Foerster, 1868			
<b>Probles</b> s. str.			

<i>saltator</i> (Fabricius, 1781)	B	<b>Enicospilus</b> Stephens, 1835
<i>subdepressus</i> (Thomson, 1889)	B	<i>Allocamptus</i> Foerster, 1868
<i>thyridialis</i> Horstmann, 1971	B	<i>inflexus</i> (Ratzeburg, 1844) *B S
<i>triangularis</i> (Gravenhorst, 1807)	B M	<i>ramidulus ramidulus</i> (Linnaeus, 1758) B M S
<b>Pectinolochus</b> Aubert, 1960	M	<i>ramidulus merdarius</i> (Gravenhorst, 1829) B M S
<i>ensifer</i> (Brischke, 1880)	B	<i>undulatus</i> (Gravenhorst, 1829) *B*M
<i>striola</i> (Thomson, 1889)	B	<b>Stauropoctonus</b> Brauns, 1889
<i>terebrator</i> Horstmann, 1971	B	<i>bombycivorus</i> (Gravenhorst, 1829) *B
<b>Gonolochus</b> Foerster, 1868		<b>MESOCHORINAE</b>
<i>caudatus</i> (Holmgren, 1860)	B M S	<b>Cidaphus</b> Foerster, 1868
<i>fenestralis</i> (Thomson, 1889)	B M S	<i>Plesiophthalmus</i> Foerster, 1868
<i>stenocari</i> (Gregor, 1941)	M	<i>alarius</i> (Gravenhorst, 1829) S
OPHIONINAE		<i>potanini</i> Kokujev, 1906 *B
OPHIONINI		<i>thuringiacus</i> Brauns, 1889 *S
<b>Ophion</b> Fabricius, 1798		<b>Astiphromma</b> Foerster, 1868
<i>costatus</i> Ratzeburg, 1848	*B	<i>marginellum</i> Holmgren, 1858 M
<i>formosus</i> Bridgman, 1882	B M	<i>mandibulare</i> (Thomson, 1886) *M
<i>longigena</i> Thomson, 1888	*M	<i>nigriceps</i> Brischke, 1880 M
<i>luteus</i> (Linnaeus, 1758)	B M S	<i>plagiatum</i> (Thomson, 1886) M
<i>minutus</i> Kriechbaumer, 1879	B M S	<i>splenium</i> (Curtis, 1833) B M S
<i>mocsaryi</i> Brauns, 1889	B M S	<i>strenuum</i> (Holmgren, 1860)
<i>obscuratus</i> Fabricius, 1798	B M S	<i>scutellatum</i> (Gravenhorst, 1829) S
<i>obscurus</i> Fabricius, 1804		<b>Mesochorus</b> Gravenhorst, 1829
<i>parvulus</i> Kriechbaumer, 1879	*B *S	<i>anomalus</i> Holmgren, 1860 B S
<i>pteridis</i> Kriechbaumer, 1879	B	<i>anthracinus</i> Kriechbaumer, 1890 *B S
<i>scutellaris</i> Thomson, 1888	*B	<i>brevicollis</i> Thomson, 1886 S
<i>longicornis</i> Brauns, 1889		<i>brevipetiolatus</i> Ratzeburg, 1844 *M S
<i>slaviceki</i> Kriechbaumer, 1892	M	<i>curvulus</i> Thomson, 1886 B
<i>ventricosus</i> Gravenhorst, 1829	B M S	<i>discitergus</i> (Say, 1836) B
<i>impressus</i> (Thunberg, 1822)		<i>facialis</i> Bridgman, 1884
<b>Eremotylus</b> Foerster, 1868		<i>dispar</i> Ruschka, 1880 B
<i>marginatus</i> (Jurine, 1807)	B M*S	<i>formosus</i> Bridgman, 1882 *B
ENICOSPILINI		<i>fulgorans</i> Curtis, 1833 B M S

<i>fulvus</i> Thomson, 1886	B	S	<i>longicalcar longicalcar</i>
<i>gemellus</i> Holmgren, 1858	B		Thomson, 1887 B M
<i>giberius</i> (Thunberg, 1822)	*B		<i>longicornis</i> Thomson, 1887 B M S
			<i>talpa</i> (Haliday, 1838) M S
<i>thoracicus</i> Gravenhorst, 1829			
<i>globulator</i> (Thunberg, 1822)	B		<b>Trieces</b> Townes, 1946
			<i>facialis</i> (Thomson, 1887) S
<i>dimidiatus</i> Holmgren, 1858			<i>rufimitranae</i> Aeschlimann, 1973 S
<i>gracilentus</i> Brischke, 1880	B		<i>tricarinatus</i> (Holmgren, 1856) B M S
<i>longicauda</i> Thomson, 1886		*M	
<i>marginatus</i> Thomson, 1886	*B		<b>Metopius</b> Panzer, 1806
<i>nigriceps</i> Thomson, 1886		S	<b>Peltocarus</b> Thomson, 1887
<i>nigripes</i> Ratzeburg, 1852	B	S	<i>croceicornis</i> Thomson, 1887 B M
<i>orbitalis</i> Holmgren, 1860	B	S	<i>dentatus</i> (Fabricius, 1779) B *M
<i>pallidus</i> Brischke, 1880	*B		<i>migratorius</i> (Fabricius, 1804)
<i>pectoralis</i> Ratzeburg, 1844	B	S	<b>Metopius</b> s. str.
<i>petiolaris</i> Brischke, 1880	M		<i>anxius</i> Wesmael, 1849 B
<i>pictilis</i> Holmgren, 1860		*S	<i>brevispina</i> Thomson, 1887 B
<i>politus</i> Gravenhorst, 1829	*B		<i>connexarius</i> Wesmael, 1849 B *S
<i>ruficornis</i> Brischke, 1880	B		<b>Tylopius</b> Townes, 1959
<i>semirufus</i> Holmgren, 1860	M		<i>leiopygus</i> Foerster, 1850 *B
<i>sylvarum</i> Curtis, 1833	B M S		<i>pinatorius</i> Brullé, 1846 B M
<i>splendidulus</i> (Ratzeburg, 1844)	B		<b>Ceratopius</b> Clément in Schmiedeknecht, 1927
<i>temporalis</i> Thomson, 1886	*B	*S	<i>dissectorius</i> (Panzer, 1805—1806) B
<i>testaceus</i> Gravenhorst, 1829	B M		<i>fuscipennis</i> Wesmael, 1849 B M S
<i>velox</i> Holmgren, 1860	B	*S	<i>scrobiculatus</i> Hartig, 1838 *S
<i>vittator</i> (Zetterstedt, 1838)	B	S	<b>Triclistus</b> Foerster, 1868
<i>vitticollis</i> Holmgren, 1860	B M		<i>aethiops</i> (Gravenhorst, 1829) M
<b>Stictopisthus</b> Thomson, 1886			<i>albicinctus</i> Thomson, 1887 S
<i>complanatus</i> (Haliday, 1838)	B M		<i>areolatus</i> Thomson, 1887 S
METOPIINAE			<i>globulipes</i> (Desvignes, 1856) B M S
<b>Chorinaeus</b> Holmgren, 1856			<i>lativentris</i> Thomson, 1887 B M S
<i>cristator</i> (Gravenhorst, 1829)	B M S		<i>longicalcar</i> Thomson, 1887 *B
<i>flavipes</i> Bridgman, 1881	M S		<i>niger</i> (Bridgman, 1883) M
<i>funebris</i> (Gravenhorst, 1829)	B M		<i>pallipes</i> Holmgren, 1873 B M S
			<i>podagratus</i> (Gravenhorst, 1829) B M S
			<i>pubiventris</i> Thomson, 1887 S
			<i>pygmaeus</i> (Cresson, 1864) M S
			<i>squalidus</i> (Holmgren, 1856) M S

<i>spiracularis</i> Thomson,			<i>pictus</i> Holmgren, 1856	*B
1887	B		<i>prosopius</i> Gravenhorst,	
<i>yponomeutae</i> Aeschlimann,			1829	M S
1973	*B		<i>suborbitalis</i>	
<b>Colpotrochia</b> Holmgren, 1855			Schmiedeknecht, 1924	*M
<i>cincta</i> (Scopoli, 1763)	B M S		<i>tibialis</i> Holmgren, 1856	*M
<i>elegantulus</i> (Schrank, 1781)			<i>turgidus</i> Holmgren, 1856	B M S
<b>Spudaeus</b> Gistel, 1848			<b>Lapton</b> Klug, 1813	
<i>Tylocomnus</i> Holmgren, 1873			<i>femoralis</i> Nees, 1813	B M*S
<i>scaber</i> (Gravenhorst,				
1829)	B M		<b>ANOMALONINAE</b>	
<b>Drepanoctonus</b> Pfankuch,			<b>ANOMALONINI</b>	
1911		S		
<b>Hypsicera</b> Latreille			<b>Anomalon</b> Panzer, 1804	
in Cuvier, 1829			<i>Nototrachys</i> Marshall, 1872	
<i>Metacoelus</i> Foerster, 1868			<i>foliator</i> (Fabricius, 1798)	B M S
<i>curvator</i> (Fabricius, 1793)	B M S			
<i>mansuetor</i> (Gravenhorst, 1807)			<b>THERIONINI</b>	
<i>femoralis</i> (Geoffroy in				
Fourcroy, 1785)	*B M*S		<b>Heteropelma</b> Wesmael, 1849	
<i>flaviceps</i> (Ratzeburg,			<i>Schizoloma</i> Wesmael, 1849	
1852)	S		<i>amictum</i> (Fabricius, 1775)	B M S
<b>Exochus</b> Gravenhorst, 1829			<i>calcator</i> (Wesmael, 1849)	B M S
<i>albicinctus</i> Holmgren,			<b>Therion</b> Curtis, 1829—1831	
1873	*B		<i>Exochilum</i> Wesmael, 1849	
<i>australis</i> (Thomson,			<i>brevicornis</i> (Gravenhorst,	
1895)	*B M S		1829)	B M S
<i>carri</i> Schmiedeknecht,			<i>circumflexum</i> (Linnaeus,	
1924	M S		1758)	B M S
<i>consimilis</i> Holmgren, 1856	S		<i>giganteum</i> (Gravenhorst,	
<i>decoratus</i> Holmgren, 1873	B M S		1829)	B M
<i>erythronotus</i>			<b>Trichomma</b> Wesmael, 1849	
(Gravenhorst, 1820)	B M S		<i>enecator</i> (Rossi, 1790)	B M S
<i>flavomarginatus</i>			<i>fulvidens</i> (Wesmael, 1849)	B M
Holmgren, 1855	*B S		<i>intermedium</i> Krieger,	
<i>gravipes</i> <i>gravipes</i>			1904	B M
(Gravenhorst, 1820	B M S		<i>occisor</i> Habermehl, 1909	S
<i>gravis</i> Gravenhorst, 1829	*B		<b>Habronyx</b> Foerster, 1868	
<i>incidens</i> Thomson, 1887	*B M		<b>Camposcopus</b> Foerster, 1868	
<i>lentipes</i> Gravenhorst,			<i>Blaptocampus</i> Thomson, 1892	
1829	*B*M		<i>Labrorychus</i> Foerster, 1868	
<i>mitratus</i> Gravenhorst,			<i>canaliculatus</i> (Ratzeburg,	
1829	B M S		1844)	B M S
<i>nigripalpis</i> Thomson,			<i>nigricorne</i> (Wesmael, 1849)	
1887	*M S		<b>Habrocampulum</b> Gauld 1976	
<i>notatus</i> Holmgren, 1856	*B M		<i>biguttatus</i> (Gravenhorst,	

<i>bellicosus</i> (Wesmael, 1849)	B M S	<i>tenuicorne</i> (Gravenhorst, 1829)
<i>ruficornis</i> (Gravenhorst, 1829)	M*S	<i>polyxenae</i> (Szépligeti, 1899)
<b>Gravenhorstia</b> Boie, 1856		<i>varitarsum</i> (Wesmael, 1849)
<b>Gravenhorstia</b> s. str.		<b>Parania</b> Morley, 1913
<i>picta</i> Boie, 1856	B	<i>geniculata</i> (Holmgren, 1857)
<b>Erigorgus</b> Foerster, 1868		<b>Periphincter</b> Townes, 1961
<i>Paranomalon</i> Viereck, 1912		<i>brevicollis</i> (Wesmael, 1849)
<i>cerinops</i> (Gravenhorst, 1829)	B M S	<b>Atrometus</b> Foerster, 1868
<i>latro</i> (Schrank, 1781)	B S	<i>insignis</i> Foerster, 1878
<i>laucopus</i> (Szépligeti, 1905)	B M	ACAENITINAE COLEOCENTRINI
<i>melanobatus melanobatus</i> (Gravenhorst, 1829)	M	<b>Coleocentrus</b> Gravenhorst, 1829
<i>procerus</i> (Gravenhorst, 1829)	B M	<i>caligatus</i> (Gravenhorst, 1829)
<i>romani</i> (Héllen, 1926)	M	<i>croceicornis</i> (Gravenhorst, 1829)
<i>ruficornis</i> (Szépligeti, 1899)	M	<i>excitator</i> (Poda, 1761)
<i>villosus</i> (Gravenhorst, 1829)	B	<b>Mesoclistus</b> Foerster, 1868
<b>Barylypa</b> Foerster, 1868		<i>rufipes</i> (Gravenhorst, 1829)
<i>amabilis</i> Tosquinet, 1900	S	<b>Leptacoenites</b> Strobl, 1902
<i>humeralis</i> (Brauns, 1895)		<i>frauenfeldi</i> (Tschek, 1868)
<i>delictor</i> (Thunberg, 1822)	B	<b>Procinetus</b> Foerster, 1868
<i>insidiator</i> (Foerster, 1878)	S	<i>decimator</i> (Gravenhorst, 1829)
<i>carinata</i> (Brische, 1880)		ACAENITINI
<i>pallida</i> (Gravenhorst, 1829)	S	<b>Arotes</b> Gravenhorst, 1829
<i>rubricator</i> (Szépligeti, 1899)	B	<i>albicinctus albicinctus</i> Gravenhorst, 1820
<i>uniguttata</i> (Gravenhorst, 1829)	B M S	<b>Acaenitus</b> Latreille, 1809
<b>Agrypon</b> Foerster, 1860		<i>dubitator</i> (Panzer, 1800)
<i>Trichonotus</i> Cameron, 1905		<b>Phaenolobus</b> Foerster, 1868
<i>anomelas</i> (Gravenhorst, 1829)	B M	<i>fulvicornis</i> (Gravenhorst, 1829)
<i>anxius</i> (Wesmael, 1849)	*B S	<i>major</i> Szépligeti, 1914
<i>clandestinus</i> (Gravenhorst, 1829)	B M S	<i>saltans</i> (Gravenhorst, 1829)
<i>flaveolatum</i> (Gravenhorst, 1807)	B M S	
<i>gracilipes</i> (Curtis, 1839)	B M S	
<i>debilis</i> (Wesmael, 1849)		
<i>delarvatus</i> (Gravenhorst, 1829)	B M	
<i>flexorius</i> (Thunberg, 1822)	B M S	

<i>terebrator</i> (Scopoli, 1763)	B M S	<i>oligomera</i> Foerster, 1868	M
<i>arator</i> (Rossi, 1790)		<i>pseudoligomera</i> Gregor, 1941	M
OXYTORINAE		<i>refractaria</i> van Rossem, <i>subdola</i> Foerster, 1871	B B
Microleptinae			
Plestiscinae			
<b>Megastylus</b> Schioedte, 1838		<b>Helictes</b> Haliday, 1838	
<i>nigriventris</i> Foerster, 1871	M	<i>erythrostoma</i> (Gmelin, 1790)	*B*M
<b>Microleptes</b> Gravenhorst, 1829		<b>Proclitus</b> Haliday, 1838	
<i>Miomeris</i> Foerster, 1868		<i>Clepticus</i> Haliday, 1838	
<i>obenbergeri</i> Gregor, 1937	M	<i>grandis</i> Foerster, 1871	S
<i>splendidulus</i> Gravenhorst, 1829	B	<b>Catastenus</b> Foerster, 1868	
<b>Oxytorus</b> Foerster, 1868		<i>femoralis</i> Foerster, 1871	M
<i>Callidiotes</i> Foerster, 1868			
<i>armatus</i> Thomson, 1883	*B	ORTHOPELMATINAE	
<i>luridator</i> (Gravenhorst, 1820)	*M	<b>Orthopelma</b> Taschenberg, 1865	
<b>Hyperacmus</b> Holmgren, 1856		<i>mediator</i> (Thunberg, 1822)	B M S
<i>crassicornis</i> (Gravenhorst, 1829)	*M S	<i>luteolator</i> (Gravenhorst, 1829)	
<b>Cylloceria</b> Schioedte, 1838		COLLYRIINAE	
<i>caligata</i> (Gravenhorst, 1829)	B M S	<b>Collyria</b> Schioedte, 1839	
<i>marginator</i> Schioedte, 1839	B M	<i>coxator</i> (Villers, 1789)	B M S
<i>melancholica</i> (Gravenhorst, 1820)	B M	<i>trichophthalma</i> (Thomson, 1877)	B
<b>Plectiscidea</b> Viereck, 1914		ORTHOCENTRINAE	
<i>Plectiscus</i> Gravenhorst, 1829		<b>Orthocentrus</b> Gravenhorst, 1829	
<i>canaliculata</i> (Foerster, 1871)	*B	<i>asper</i> (Gravenhorst, 1829)	*B
<i>subcurvata</i> (Foerster, 1871)	*M	<i>corrugatus</i> Holmgren, 1856	B M
<i>subtilis</i> (Foerster, 1871)	*B*M	<i>frontator</i> (Zetterstedt, 1838)	*M
<i>terebrator</i> (Foerster, 1871)	*B	<i>fulvipes</i> Gravenhorst, 1829	B M
<b>Eusterinx</b> Foerster, 1868		<i>longicornis</i> Holmgren, 1856	*B
<i>Holomeristus</i> Foerster, 1868		<i>marginatus</i> Holmgren, 1856	M
<i>Dalaatorrea</i> Ashmead, 1902		<i>petiolaris</i> Thomson, 1897	*B
<i>alpigena</i> (Strobl, 1903)	*M	<i>sannio</i> Holmgren, 1856	B M
<i>basalis</i> Foerster, 1868	M	<i>spurius</i> Gravenhorst, 1829	B M
<i>divulgata</i> Foerster, 1871	M	<i>stigmaticus</i> Holmgren, 1856	*B
<i>fleischeri</i> Gregor, 1941	M	<i>protuberans</i> Holmgren, 1856	

<b>Stenomacrus</b> Foerster, 1868		
<i>binotatus</i> (Holmgren, 1856)	B M	
<i>exserens</i> (Thomson, 1897)	*B*M	
<i>molestus</i> (Holmgren, 1856)	*B	
<i>ochripes</i> (Holmgren, 1856)	*B	
<i>silvaticus</i> (Holmgren, 1856)	*B*M	
<i>ventralis</i> (Holmgren, 1856)	*M	
<b>Neurateles</b> Ratzeburg, 1848		
<i>papyraceus</i> Ratzeburg, 1848	B M S	
<i>compressus</i> (Thomson, 1897)		
<b>DIPLAZONTINAE</b>		
<b>Homotropus</b> Foerster, 1868		
<i>Homocidus</i> Morley, 1911		
<i>dimidiatus</i> (Schrank, 1802)	B M	
<i>elegans</i> (Gravenhorst, 1829)	M S	
<i>rufonotatus</i> (Holmgren, 1856)		
<i>fissorius</i> (Gravenhorst, 1829)	B M	
<i>megaspis</i> Thomson, 1890	M	
<i>neopulcher</i> Horstmann, 1968	M	
<i>nigritarsus nigritarsus</i> (Gravenhorst, 1829)	B M S	
<i>pallipes</i> (Gravenhorst, 1829)	*B	
<i>pictus</i> (Gravenhorst, 1829)	B M	
<i>signatus</i> (Gravenhorst, 1829)	M S	
<i>strigator</i> (Fabricius, 1793)	*B*M	
<i>tarsatorius</i> (Panzer, 1809)	B M S	
<i>pulchellus</i> (Desvignes, 1862)		
<i>tricolor</i> (Stelfox, 1941)	*B M	
<b>Enizemum</b> Foerster, 1868		
<i>nigricorne</i> (Thomson, 1890)	M	
<i>ornatum</i> (Gravenhorst, 1829)		B M
<b>Woldstedtius</b> Carlson, 1979		
<i>Syrphocontonus</i> Foerster, 1868		
<i>abdominator</i> (Bridgman, 1886)		B M
<i>citropectoralis</i> (Schmiedeknecht, 1926)		
<i>biguttatus</i> (Gravenhorst, 1829)		B M S
<i>flavolineatus</i> <i>flavolineatus</i> Gravenhorst, 1829		B M
<b>Campocraspedon</b> Uchida, 1957		
<i>alpinus</i> (Holmgren, 1856)		*S
<i>caudatus</i> (Thomson, 1890)		*M
<b>Phthorima</b> Foerster, 1868		
<i>compressa</i> (Desvignes, 1856)		M S
<b>Syrphophilus</b> Dasch, 1964		
<i>bizonarius</i> (Gravenhorst, 1829)		B M S
<i>cingulatus</i> (Holmgren, 1856)		
<i>tricinctarius tricinctarius</i> (Thunberg, 1822)		B M S
<i>cinctus</i> (Gravenhorst, 1829)		
<i>lateralis</i> (Gravenhorst, 1829)		
<b>Tymmophorus</b>		
Schmiedeknecht, 1913		
<i>graculus</i> Gravenhorst, 1829		M
<i>obscuripes</i> (Holmgren, 1856)		
<i>rufocinctus</i> Desvignes, 1862)		
<i>rufiventris</i> (Gravenhorst, 1829)		B M
<i>suspiciosus</i> (Brischke, 1871)		
<b>Diplazon</b> , Nees, 1818		
<i>annulatus</i> (Gravenhorst, 1829)		M
<i>deletus</i> (Thomson, 1890)		M
<i>laetatorius</i> (Fabricius, 1781)		B M S
<i>pectoratorius</i> (Gravenhorst, 1829)		B M S
<i>tetragonus</i> (Thunberg, 1822)		B M S
<i>tricinctus</i> (Gravenhorst, 1829)		

<i>tibiatorius tibiatorius</i> (Thunberg, 1822)	B M S	<i>debilis</i> (Gravenhorst, 1829)	M
<i>albosignatus</i> (Gravenhorst, 1829)		<i>moravicus</i> (Gregor, 1939)	M
<i>varicoxa</i> (Thomson, 1890)	*B*M	<i>nigricollis</i> (Wesmael, 1844)	M
<i>Bioblapsis</i> Foerster, 1868		<i>rufipes</i> (Wesmael, 1848)	
<i>polita</i> (Vollenhoven, 1878)	*M	<i>rufescens</i> (Pic, 1914)	*B
<i>flavipes</i> (Holmgren, 1856)		<i>thoracicus</i> (Gravenhorst, 1829)	B M S
<i>Promethes</i> Foerster, 1868		<i>tineidarum</i> (Girault, 1871)	B M
<i>Promethus</i> Thomson, 1890		<i>truncator</i> (Fabricius, 1798)	B M S
<i>nigriventris</i> (Thomson, 1890)	*B	<i>Heterischnus</i> s. str.	
<i>melanaspis</i> (Thomson, 1890)	*S	<i>pullex</i> (Müller, 1776)	B M
<i>sulcator sulcator</i> (Gravenhorst, 1829)	B M S	<i>Misetus</i> Wesmael, 1844	
<i>Sussaba</i> Cameron, 1909		<i>oculatus</i> Wesmael, 1844	*B
<i>cognata cognata</i> (Holmgren, 1856)	B	<i>Hemichneumon</i> Wesmael, 1857	
<i>cognata albicoxa</i> (Thomson, 1890)	B*M	<i>elongatus</i> (Ratzeburg, 1852)	B M S
<i>dorsalis dorsalis</i> (Holmgren 1856)	B	<i>Trachyarus</i> Thomson, 1891	
<i>elongata elongata</i> Provancher, 1874)	B M	<i>corvinus</i> Thomson, 1891	S
<i>erigator</i> (Fabricius, 1793)	B M S	<i>Diaschiaspis</i> Foerster, 1868	
<i>festivus</i> (Fabricius, 1793)		<i>campoplegoides</i> Holmgren, 1889	M
<i>pulchella</i> (Holmgren, 1856)	B M	<i>Oronotus</i> Wesmael, 1844	
<i>punctiventris</i> (Thomson, 1890)	M	<i>binotatus</i> (Gravenhorst, 1829)	B S
ICHNEUMONINAE		<i>Dicaelotus</i> Wesmael, 1844	
ALOMYINI		<i>Deloglyptus</i> Foerster, 1868	
<i>Alomya</i> Panzer, 1806		<i>kriechbaumeri</i> (Holmgren, 1890)	M
<i>debelator</i> (Fabricius, 1775)	B M S	<i>parvulus</i> (Gravenhorst, 1829)	M
<i>semiflava</i> Stephens, 1835	B	<i>pictus</i> (Schmiedeknecht, 1908)	*B*M
PHAEOGENINII		<i>pumilus</i> (Gravenhorst, 1829)	M S
<i>Heterischnus</i> Wesmael, 1859		<i>morosus</i> (Wesmael, 1855)	
RHEXIDERMUS Foerster, 1868		<i>punctiventris</i> (Thomson, 1891)	*B
<i>Ischnopsidea</i> Viereck, 1914		<i>pusillator</i> (Gravenhorst, 1822)	*B
<i>anomalus</i> (Holmgren, 1857)	B	<i>pusillus</i> Holmgren, 1890	M
		<i>resplendens</i> Holmgren, 1890	M

<i>rufilimbatus</i> Gravenhorst, 1820	M		<i>collaris</i> (Gravenhorst, 1829)	*B
<b>Cinxaelotus</b> Holmgren, 1889			<i>pimplarius</i> Wesmael, 1848	M
<i>erythrogaster</i> Holmgren, 1889	*B*M		<i>subtilicornis</i> (Gravenhorst, 1829)	B M
<b>Colpognathus</b> Wesmael, 1844			<i>troglodytes</i> (Gravenhorst, 1829)	B M
<i>armatus</i> Thomson, 1891	*B		<i>ustulatus</i> Holmgren, 1889	*B
<i>celerator</i> (Gravenhorst, 1807)	B M S		<b>Aethecerus</b> Wesmael, 1844	
<i>divisus</i> Thomson, 1891	*B		<i>discolor</i> Wesmael, 1844	M
<b>Centeterus</b> Wesmael, 1844			<i>dispar</i> Wesmael, 1844	M
<i>confector</i> (Gravenhorst, 1829)	B M S		<i>graniger</i> Thomson, 1891	M
<i>major</i> Wesmael, 1844	B M S		<i>foveolatus</i> Gregor, 1940	M S
<i>nigricornis</i> Thomson, 1891	*S		<i>moravicus</i> Gregor, 1940	M
<i>opprimator</i> (Gravenhorst, 1820)	B M		<i>nitidus</i> Wesmael, 1844	M S
<b>Eparces</b> Foerster, 1868			<i>pallicoxa</i> Thomson, 1891	B M
<i>grandiceps</i> Thomson, 1891	M		<i>placidus</i> Wesmael, 1844	B M
<b>Herpestomus</b> Wesmael, 1844			<b>Epitomus</b> Foerster, 1868	
<i>arridens</i> Gravenhorst, 1829	M S		<i>parvus</i> Thomson, 1891	B M
<i>xanthops</i> (Gravenhorst, 1829)			<b>Dirophanes</b> Foerster, 1868	
<i>brunnicornis</i> (Gravenhorst, 1829)	B M		<i>fulvitarsis</i> (Wesmael, 1844)	B M S
<i>flavoclypeatus</i> Strobl, 1900	S		<b>Phaeogenes</b> Wesmael, 1844	
<i>nasutus</i> Wesmael, 1844	M		<i>Proscus</i> Holmgren, 1889	
<i>phaeocerus</i> (Wesmael, 1844)	M		<i>amoenus</i> Wesmael, 1844	B
<b>Eriplatys</b> Foerster, 1868			<i>bisignatus</i> Holmgren, 1890	B M S
<i>ardeicollis</i> (Wesmael, 1844)	B M S		<i>callopus</i> Wesmael, 1844	*B
<i>kratochvili</i> Gregor, 1943	M		<i>coryphaeus</i> Wesmael, 1844	*B
<b>Stenodontus</b> Berthoumieu, 1896			<i>curator</i> (Thunberg, 1822)	*M
<i>marginellus</i> Gravenhorst, 1829	*S		<i>eques</i> Wesmael, 1844	M
<b>Notosemus</b> Foerster, 1868			<i>flavidens</i> Wesmael, 1844	S
<i>bohemani</i> (Wesmael, 1855)	*B		<i>fuscicornis</i> Wesmael, 1844	B M
<b>Micrope</b> Thomson, 1891			<i>infimus</i> Wesmael, 1844	*B*M S
<i>macilenta</i> (Wesmael, 1844)	*B		<i>invisor</i> (Thunberg, 1822)	B M S
<b>Diadromus</b> Wesmael, 1844			<i>impiger</i> Wesmael, 1844	B
			<i>ischiomelinus</i> (Gravenhorst, 1829)	B M S
			<i>lascivus</i> Wesmael, 1855	B M S
			<i>maculicornis</i> (Stephens, 1835)	B M S
			<i>melanogonos</i> (Gmelin in Linnaeus, 1790)	M S
			<i>montanus</i> Thomson, 1891	*M
			<i>muricifer</i> Holmgren, 1890	M
			<i>mysticus</i> Wesmael, 1855	M
			<i>nigridens</i> Wesmael, 1844	M

<i>ophthalmicus</i> Wesmael,			
1844	B M		
<i>osculator</i> (Thunberg, 1822)	B M		
<i>planifrons</i> Wesmael, 1844	B M		
<i>semivulpinus</i> (Gravenhorst, 1829)	M		
<i>rufator</i> (Stephens, 1835)			
<i>scutellaris</i> Wesmael, 1844	M		
<i>socialis</i> (Ratzeburg, 1852)	B M		
<i>spiniger</i> (Gravenhorst, 1829)	*M		
<i>stipator</i> Wesmael, 1855	M		
<i>tegularis</i> Thomson, 1891	*B		
PLATYLABINI			
<b>Apaeleticus</b> Wesmael, 1844			
<i>inimicus</i> (Gravenhorst, 1820)	M		
<b>Ectopius</b> Wesmael, 1859			
<i>rubellus</i> (Gmelin in Linnaeus, 1790)	B M		
<b>Linyxus</b> Cameron, 1903			
<i>exhortator exhortator</i> (Fabricius, 1787)	M		
<b>Cyclolabus</b> Heinrich, 1935			
<i>nigricollis</i> (Wesmael, 1844)	*M		
<i>pactor</i> (Wesmael, 1844)	M		
<b>Hypomecus</b> Wesmael, 1844			
<i>quadriannulatus</i> <i>quadriannulatus</i> Gravenhorst, 1829)	*B M		
<b>Asthenolabus</b> Heinrich, 1951			
<i>Stenolabus</i> Heinrich, 1936			
<i>daemon</i> (Wesmael, 1844)	*M		
<i>stali</i> (Holmgren, 1871)	M		
<b>Rhyssolabus</b> Berthoumieu, 1896			
<i>bassicus</i> (Tischbein, 1868)	*B*M		
<b>Pristiceros</b> Gravenhorst, 1829			
<i>serrarius</i> Gravenhorst, 1829	*B		
Platylabus Wesmael, 1844			
<i>decipiens</i> Wesmael, 1848	B M S		
<i>gigas</i> Kriechbaumer, 1886	*M		
<i>histrio</i> Wesmael, 1855	M		
<i>iridipennis</i> (Gravenhorst, 1829)	B M		
<i>opaculus</i> Thomson, 1888	*M		
<i>pedatorius</i> (Fabricius, 1793)	B M		
<i>pumilio</i> Holmgren, 1871	S		
<i>rufus</i> Wesmael, 1844	B M		
<i>suborbitalis</i>			
Kriechbaumer, 1894	M		
<i>sulci</i> Gregor, 1940	B M		
<i>transversus</i> Bridgman, 1889	*M		
<i>lativentris</i> Thomson, 1894			
<i>vibratorius</i> (Thunberg, 1822)	M		
Dentilabus Heinrich, 1975			
<i>variegatus</i> (Wesmael, 1844)	*M		
Pyramidophorus Tischbein, 1882			
<i>flavoguttatus</i> Tischbein, 1882	B M		
Eurylabini			
Eurylabus Wesmael, 1844			
<i>larvatus</i> (Christ, 1791)	*M		
<i>tristis</i> (Gravenhorst, 1829)	*B		
Probolus Wesmael, 1844			
<i>alticola</i> (Gravenhorst, 1820)	B M		
<i>concinnus</i> Wesmael, 1853	B M S		
<i>slavicekti</i> (Kriechbaumer, 1893)			
<i>culpatorius</i> (Linnaeus, 1758)	*M		
Goedartini			
Goedartia Boie, 1841			
<i>alboguttata</i> (Gravenhorst, 1829)	B M S		

<b>LISTRODROMINI</b>			
<b>Listrodromus</b> Wesmael, 1844			
<i>nycthemerus</i> (Gravenhorst, 1820)	*B M S		
<b>Neotypus</b> Foerster, 1868			
<i>lapidator lapidator</i> (Fabricius, 1793)	B M S		
<i>nobilitor nobilitator</i> (Gravenhorst, 1807)	B M S		
<b>Anisobas</b> Wesmael, 1844			
<i>cingulatorius</i> (Gravenhorst, 1820)	B M		
<i>hostilis</i> (Gravenhorst, 1820)	B M S		
<i>platystylus</i> Thomson, 1888	*M*S		
<b>ACANTHOJOPPINI</b>			
<b>Pseudoplatylabus</b> Smits van Burgst, 1920			
<i>uniguttatus</i> (Gravenhorst, 1829)	B M S		
<b>ICHNEUMONINAE</b>			
Joppini			
<b>Cratichneumon</b> Thomson, 1893			
<i>cordiger</i> (Kriechbaumer, 1882)	*S		
<i>coruscator</i> (Linnaeus, 1758)	*M		
<i>culex</i> (Müller, 1776)	B M S		
<i>fabricator</i> (Fabricius, 1793)	B M		
<i>foersteri</i> (Wesmael, 1848)	*B		
<i>fugitivus</i> (Gravenhorst, 1829)	*M S		
<i>infidus</i> (Wesmael, 1848)	*M		
<i>hostylus</i> (Thomson, 1887)			
<i>luteiventris</i> (Gravenhorst, 1820)	B M		
<i>melanocastanus</i> (Gravenhorst, 1820)	*B		
<i>rubro-ater</i> (Ratzeburg, 1852)			
<i>polystictus</i> (Kriechbaumer, 1887)	M S		
<i>rufifrons</i> (Gravenhorst, 1829)	M		
<i>semirufus</i> (Gravenhorst, 1820)		*M	
<i>sicarius</i> (Gravenhorst, 1829)		B M	
<i>varipes</i> (Gravenhorst, 1829)		M	
<i>versator</i> (Thunberg, 1822)		B M	
<i>viator</i> (Scopoli, 1763)		B M S	
<i>nigritarius</i> (Gravenhorst, 1820)			
<b>Eupalamus</b> Wesmael, 1844			
<i>lacteator</i> (Gravenhorst, 1829)		*B	
<i>oscillator</i> Wesmael, 1844		*M	
<i>wesmaeli</i> Thomson, 1886		*B M	
<b>Anisopygus</b> Kriechbaumer, 1888			
<i>pseudonymus</i> Wesmael, 1888		*B	
<b>Platylabops</b> Heinrich, 1950			
<i>corniculus</i> Wesmael, 1855		M	
<b>Aoplus</b> Tischbein, 1874			
<i>altercator</i> (Wesmael, 1855)		*B	
<i>castaneus</i> (Gravenhorst, 1820)		B M	
<i>defraudator</i> (Wesmael, 1844)		B M	
<i>ochropis</i> (Gmelin in Linnaeus, 1790)		*M	
<i>ratzeburgii</i> (Hartig, 1838)		*S	
<i>ruficeps ruficeps</i> (Gravenhorst, 1829)		B M	
<b>Stenolabus</b> Heinrich, 1938			
<i>pictus</i> (Gravenhorst, 1829)		B M S	
<b>Homotrechus</b> Foerster, 1868			
<i>locutor</i> (Thunberg, 1822)		B M	
<i>varipes</i> (Gravenhorst, 1829)		M	
<b>Crypteffigies</b> Heinrich, 1961			
<i>lanius</i> (Gravenhorst, 1829)		M	

<b>Melanichneumon</b> Thomson, 1893		<b>Vulgichneumon</b> Heinrich, 1962	
<i>albibictus</i> (Gravenhorst, 1820)	*B M	<i>deceptor</i> (Gravenhorst, 1820)	M
<i>leucocheilus</i> (Wesmael, 1844)	*B M	<i>vestigator</i> (Wesmael, 1893)	
<b>Virgichneumon</b> Heinrich, 1977		<i>saturatorius saturatorius</i> (Linnaeus, 1758)	B
<i>albilineatus</i> (Gravenhorst, 1820)	B	<i>suavis</i> (Gravenhorst, 1829)	M
<i>leucomelas</i> (Gmelin in Linnaeus, 1790)		<i>lepidus</i> (Gravenhorst, 1829)	
<i>albosignatus</i> (Gravenhorst, 1829)	*B	<b>Sympasis</b> Townes in Townes, Townes et Momoi, 1965	
<i>callicerus</i> (Gravenhorst, 1820)	*S	<i>alboguttatus</i> (Gravenhorst, 1820)	B
<i>digrammus</i> (Gravenhorst, 1820)	B	<i>eburnifrons</i> (Wesmael, 1857)	*M
<i>faunus</i> (Gravenhorst, 1829)	S	<i>lineator</i> (Fabricius, 1781)	B M S
<i>monostagon</i> (Gravenhorst, 1820)	B M	<i>rimulosus</i> (Thomson, 1893)	*M
<i>tergenus</i> (Gravenhorst, 1820)	M	<i>rufinus</i> (Gravenhorst, 1820)	*B
<b>Barichneumon</b> Thomson, 1893		<i>scutellator</i> (Gravenhorst, 1829)	M
<i>bilunulatus</i> (Gravenhorst, 1829)	B M S	<b>Orgichneumon</b> Heinrich, 1961	
<i>gibbosus</i> Brischke, 1878	*B	<i>calcatorius</i> (Thunberg, 1822)	*B *M
<i>chionomus</i> (Wesmael, 1844)	*S	<b>Paracoelichneumon</b> Heinrich, 1978	
<i>citrinops</i> (Wesmael, 1857)	M	<i>rubens</i> (Fonscolombe, 1847)	M *S
<i>derogator</i> (Wesmael, 1844)	M S	<b>Lymantrichneumon</b> Heinrich, 1968	
<i>perversus</i> (Kriechbaumer, 1893)	B M S	<i>disparis</i> (Poda, 1761)	B M S
<i>scriptorius</i> (Thunberg, 1822)	B M S	<b>Ichneumon</b> Linnaeus, 1756	
<b>Stenobarichneumon</b> Heinrich, 1962		<i>Pterocormus</i> Foerster, 1850	
<i>basiglyptus</i> (Kriechbaumer, 1890)	M S	<i>albicollis</i> Wesmael, 1857	*M
<i>citator</i> (Thunberg, 1822)	M S	<i>albiger</i> Wesmael, 1844	M
<b>Rictichneumon</b> Heinrich, 1961		<i>tempestivus</i> Holmgren, 1864	
<i>pachymerus</i> (Hartig, 1838)	*B	<i>albiornatus</i> Tischbein, 1877	B M
		<i>variegatorius</i> Holmgren, 1864	
		<i>analis</i> Gravenhorst, 1829	B M
		<i>bucculentus</i> Wesmael, 1844	B M

<i>caedator</i> Gravenhost, 1829	B M	<i>mordax</i> Kriechbaumer, 1875	*M*S
<i>caloscelis</i> Wesmael, 1844	*B M	<i>paragravipes</i> Rasnitsyn, 1981	*M
<i>cessator</i> Müller, 1776	B*M	<i>gravipes</i> Wesmael, 1855	
<i>computatorius</i> Müller, 1776	B M S	<i>primatorius primatorius</i>	
<i>confusor</i> Gravenhorst, 1820	B M	<i>Foerster</i> , 1771	B M
<i>crassifemur</i> Thomson, 1886	M	<i>quadrialbatus</i>	
<i>croceipes</i> Wesmael, 1848	B M	Gravenhorst, 1820	M S
<i>didymus</i> Gravenhorst, 1829	*B	<i>quaesitorius</i> Linnaeus, 1761	*M
<i>emancipatus</i> Wesmael, 1844	M*S	<i>rarus</i> Tischbein, 1873	*M
<i>eumerus</i> Wesmael, 1857	*B M	<i>rogenhoferi rogenhoferi</i>	
<i>extensorius</i> Linnaeus, 1758	B M	Kriechbaumer, 1888	M
<i>formosus</i> Gravenhorst, 1829	M	<i>sarcitorius sarcitorius</i>	
<i>obessor</i> Wesmael, 1844		Linnaeus, 1758	B M S
<i>gracilentus</i> Wesmael, 1844	B M	<i>septentrionalis</i>	
<i>gracilicornis</i>		Holmgren, 1864	*M
Gravenhorst, 1829	B M	<i>stramentarius</i>	
<i>haglundi</i> Holmgren, 1864	*B	Gravenhorst, 1820	*B*M
<i>illuminatorius</i>		<i>subquadratus</i> Thomson, 1887	*S
Gravenhorst, 1829	*M*S	<i>suspiciosus</i> Wesmael, 1844	B M
<i>zonalis</i> Gravenhorst, 1829		<i>terminatorius</i>	
<i>inops</i> Holmgren, 1880	*M	Gravenhorst, 1820	B M
<i>insidiosus</i> Wesmael, 1844	*M S	<i>trispilus</i> Thomson, 1888	*M
<i>inquinatus</i> Wesmael, 1844	*B M	<i>tuberculipes</i> Wesmael, 1848	*B M S
<i>lanquidus</i> Wesmael, 1844	B M	<i>validicornis</i> Holmgren, 1864	B M S
<i>latrator</i> Fabricius, 1781	*B*M	<i>vulneratorius</i>	
<i>ligatorius</i> Thunberg, 1822	B M S	Zetterstedt, 1838	*B*M
<i>longeareolatus</i> Thomson, 1886	B M	<i>xanthorius</i> Foerster, 1771	*B
<i>lugens</i> Gravenhorst, 1829	B M	<b>Thyrateles</b> Perkins, 1953	
<i>macrocerus</i> Thomson, 1886	*B	<i>camelinus</i> (Wesmael, 1844)	B M S
<i>melanotis</i> Holmgren, 1864	*M	<i>haereticus</i> (Wesmael, 1854)	B*M*S*
<i>memorator</i> Wesmael, 1844	*B M	<b>Stenichneumon</b> Thomson, 1893	
<i>möllerii</i> Holmgren, 1884	*B	<i>culpator culpator</i>	
<i>molitorius</i> Holmgren, 1861	B M	(Schrank, 1802)	B M S
		<i>militarius militarius</i>	
		(Thunberg, 1822)	B M
		<b>Patrocloides</b> Heinrich, 1961	
		<i>sputator</i> (Fabricius, 1793)	B M S

<b>Patroclus</b> Cresson 1873		
<i>Pseudoamblyteles</i> Heinrich, 1926		
<i>homocerus homocerus</i> (Wesmael, 1854)	*B	
<b>Chasmias</b> Wesmael, 1844		
<i>motatorius</i> (Fabricius, 1775)	B M	
<i>paludator</i> (Desvignes, 1854)	M	
<b>Exephanes</b> Wesmael, 1844		
<i>ischioxanthus</i> (Gravenhorst, 1829)	M S	
<i>occupator</i> (Gravenhorst, 1829)	*B *S	
<i>variegator</i> Tischbein, 1881	B M	
<b>Diphyus</b> Kriechbaumer, 1890		
<i>amatorius</i> (Müller, 1776)	B M S	
<i>luctatorius</i> (Linnaeus, 1758)	*B	
<i>mercatorius mercatorius</i> (Fabricius, 1793)	B M S	
<i>monitorius</i> (Panzer, 1801)	B M S	
<i>montivagans</i> (Berthoumieu, 1896)	B M S	
<i>palliatorius</i> (Gravenhorst, 1829)	B M S	
<i>pseudomercator</i> Heinrich, 1978	B M S	
<i>quadripunctarius</i> (Müller, 1776)	B M S	
<i>raptorius</i> (Linnaeus, 1758)	B M S	
<i>septemguttatus</i> (Gravenhorst, 1829)	*M	
<i>tricolor</i> Kriechbaumer, 1890	B M	
<i>trifasciatus</i> (Gravenhorst, 1829)	*B *S	
<b>Limerodops</b> Heinrich, 1949		
<i>elongatus</i> (Brischke, 1878)	*B	
<i>subsericans</i> (Gravenhorst, 1820)	B M S	
<i>uniguttatus</i> (Gravenhorst, 1829)	B	
<b>Amblyteles</b> Wesmael, 1844		
<i>armatorius</i> Foerster, 1771	B M	
<i>diversorius</i> (Stephens, 1835)		
<b>Achaius</b> Cameron, 1903		
<i>oratorius oratorius</i> (Fabricius, 1793)	B M	
<b>Hepiopelmus</b> Wesmael, 1844		
<i>melanogaster</i> (Gmelin in Linnaeus, 1790)	B M	
<i>variegatorius variegatorius</i> (Panzer, 1800)	M	
<b>Ctenichneumon</b> Thomson, 1894		
<i>castigator</i> (Fabricius, 1793)	B M S	
<i>devylderi</i> (Holmgren, 1871)	B M	
<i>divisorius</i> (Gravenhorst, 1820)	*B	
<i>funereus</i> (Geoffroy in Fourcroy, 1785)	*M	
<i>inspector</i> (Wesmael, 1844)	M	
<i>panzeri</i> (Wesmael, 1844)	*B*M	
<i>repentinus</i> (Gravenhorst, 1820)	*M*S	
<b>Obtusodonta</b> Heinrich, 1962		
<i>equitatoria</i> Panzer, 1786	*B	
<b>Spilichneumon</b> Thomson, 1894		
<i>ammonius</i> (Gravenhorst, 1820)	*B	
<i>occisor</i> (Fabricius, 1793)	B M	
<i>occisorius</i> (Fabricius, 1804)		
<i>johansoni</i> (Holmgren, 1871)	*B M	
<b>Eutanyacra</b> Cameron, 1903		
<i>crispatorius</i> (Linnaeus, 1758)	*B	
<i>glaucatorius glaucatorius</i> (Fabricius, 1793)	B M S	
<i>pictus</i> (Schrank, 1776)	*M S	
<i>vadatorius</i> (Illiger, 1807)		
<b>Triptognathus</b> Berthoumieu, 1904		
<i>atrides</i> (Gravenhorst, 1820)	B M S	

<b>Triptognathops</b> Heinrich, 1978			
<i>bicolorus</i> (Kriechbaumer, 1882)	*B		
<b>Limerodes</b> Wesmael, 1844			
<i>arctiventris</i> (Boie, 1841)	B M		
<b>Tricholabus</b> Thomson, 1894			
<i>strigatorius strigatorius</i> (Gravenhorst, 1829)	B M		
<b>Hoplismenus</b> Gravenhorst, 1829			
<i>Zanthonjoppa</i> Cameron, 1901			
<i>albifrons</i> Gravenhorst, 1829	B M		
<i>armatorius</i> (Fabricius, 1784)			
<i>bidentatus</i> (Gmelin in Linnaeus, 1790)	*M*S		
<i>luteus</i> Gravenhorst, 1829	B M		
PROTICHNEUMONINI			
<b>Coelichneumon</b> Thomson, 1893			
<i>buguttulus</i> (Kriechbaumer, 1875)	M		
<i>bilineatus</i> (Gmelin in Linnaeus, 1790)	*S		
<i>castaneiventris</i> (Gravenhorst, 1829)	*M		
<i>centummaculatus</i> (Christ, 1791)	*S		
<i>comitator</i> (Linnaeus, 1758)	B M		
<i>ferreus</i> (Gravenhorst, 1829)			
<i>cyaniventris</i> (Wesmael, 1859)	*B		
<i>delitorius delitorius</i> (Linnaeus, 1758)	B M S		
<i>desinistratorius</i> (Thunberg, 1822)	M S		
<i>subguttatus</i> (Gravenhorst, 1829)			
<i>falsificus</i> (Wesmael, 1844)	S		
<i>haemorrhoidalis</i> (Gravenhorst, 1820)	M S		
<i>leucocerus</i> (Gravenhorst, 1820)	M		
<i>nigricornis</i> (Wesmael, 1844)	B M		
<i>nobilis</i> (Wesmael, 1857)	*M		
<i>opulentus</i> (Taschenberg, 1871)	M		
<i>serenus</i> (Gravenhorst, 1820)	*B*M		
<i>sugillatorius</i> (Linnaeus, 1758)	*B*M		
<b>Protichneumon</b> Thomson, 1893			
<i>coqueberti</i> (Wesmael, 1848)	B M		
<i>fusorius</i> (Linnaeus, 1758)	B M		
<i>pisorius</i> (Linnaeus, 1758)	*B		
<b>Amblyjoppa</b> Cameron, 1902			
<i>fuscipennis</i> (Wesmael, 1844)	B M S		
<i>proteus proteus</i> (Christ, 1791)	B M S		
<i>laminatorius</i> (Fabricius, 1798)			
<b>Catadelphus</b> Wesmael, 1854			
<i>arrogator</i> (Fabricius, 1781)	B M		
TROGINI			
<b>Trogs</b> Panzer, 1806			
<i>lapidator lapidator</i> (Fabricius, 1787)	B M S		
<i>lapidator coerulator</i> (Fabricius, 1804)	B		
<b>Psilomastax</b> Tischbein, 1868			
<i>pyramidalis</i> Tischbein, 1868	*B		
<i>pictus</i> (Kriechbaumer, 1882)			
<b>Callajoppa</b> Cameron, 1903			
<i>exaltatoria exaltatoria</i> (Panzer, 1804)	B M S		
<i>cirrogaster cirrogaster</i> (Schrank, 1781)	B M S		

STEPHANOIDEA

JOSEF SEDIVÝ

A single species is known to occur in the warm part of Czechoslovakia. The first findings in Czechoslovakia have been published by Kolubajiv (1938). The species was reared from oak branches infested with larvae of cerambycid beetles.

Reference: Kolubajiv, S., 1938: Výskyt vzácných hymenopter v Československu. *Časopis Čs. Spol. Ent.* XXXV; 21—28.

STEPHANOIDEA  
STEPHANIDAE

**Stephanus** Panzer, 1801

*coronatus* Panzer, 1801      B M S  
*serrator* (Fabricius, 1798)

## CHALCIDOIDEA

VLADIMÍR KALINA

The superfamily Chalcidoidea is probably one of the most difficult groups of parasitic Hymenoptera. For a considerable part of the family, comprehensive works and keys are still lacking. Some of the families (e. g., Trichogrammatidae, Mymaridae) have not been sufficiently studied as to their occurrence in Czechoslovakia.

Investigations into this superfamily in Czechoslovakia began only after 1945. Until that time, only a quite insignificant number of species was reported from the territory of Czechoslovakia. After 1945, the greatest credit for the knowledge of the chalcidoid fauna of the territory is due to A. Hoffer (who published exclusively on the family Encyrtidae) and Z. Bouček (who published numerous papers on most chalcid families). S. Kolubajiv treated the chalcids fairly systematically from the point of view of forestry entomology. At present, only the present author studies this group in Czechoslovakia.

The collections of A. Hoffer and Z. Bouček are deposited in the National Museum in Prague, the author's collection in the Institute of Applied Ecology and Ecotechnology at Kostelec nad Černými lesy.

Most chalcidoid species are solitary or gregarious parasitoids of other insects or, exceptionally, some other arthropods. They infest all developmental stages of their hosts. Some species are hyperparasitic, some are polyembryonic. The hosts of many species are still unknown. Phytophagous species are less common; they are mainly represented in the families Eurytomidae and Torymidae. Some of the entomophagous species are utilised in the biological control of insect pests.

In Czechoslovakia, the least thoroughly studied regions include northern and western Bohemia, northern Moravia and northern Slovakia. Discoveries of further members of the chalcidoid fauna of Czechoslovakia may be expected, above all, in the families Mymaridae, Trichogrammatidae, Eurytomidae, Pteromalidae, Eulophidae and Encyrtidae.

In compiling this check-list, the classification has been used as stated in the papers by Bouček et Askew 1968, Bouček et Graham 1978, Domenichini 1966, Graham 1969, 1987, Hayat 1983, Hansson 1985, Jasnoch 1976, Noyes et Hayat 1984 and Tryapitzin 1973a, b.

References: Bouček Z. et Askew A. A., 1968: Palearctic Eulophidae (excl. Tetrastichinae). Index of entomophagous insects. Le Francois, Paris, 254 pp. — Bouček Z. et Graham M. W. R. de V., 1978: Chalcidoidea, in Fitton et al.: Hymenoptera, A check list of British insects. London, 1—159. — Domenichini G., 1966: Palearctic Tetrastichinae. Index of entomofagous insects. Le Francois, Paris, 101 pp. — Graham M. W. R.

de V., 1969: The Pteromalidae of North-Western Europe (Hymenoptera: Chalcidodea). *Bull. Br. Mus. nat. Hist., Ent. Supplement* 16: 908 pp; 1987: A reclassification of the European Tetrastichinae (Hymenoptera: Eulophidae), with a revision of certain genera. *Bull. Br. Mus. nat. Hist., Ent.* 55 (1): 1—392. — Hayat M., 1983: The genera of Aphelinidae (Hymenoptera) of the world. *Syst. Ent.*, 8: 63—102. — Hansson C., 1985: Taxonomy and biology of the Palearctic species of Chrysocharis Foerster 1856 (Hymenoptera: Eulophidae). *Ent. scand. Supplement* 26: 1—130. — Jasnoch V. A., 1976: Classification of parasitic Hymenoptera of the family Aphelinidae (Chalcidoidea). *Ent. Obozr.*, 55 (1): 159—168. — Noyes J. S. et Hayat M., 1984: A review of the genera of Indo-Pacific Encyrtidae (Hymenoptera: Chalcidoidea). *Bull. Br. Mus. nat. Hist. Ent.*, 48 (3): 131—395. — Tryapitzin V. A., 1973a: Classification of parasitic Hymenoptera of the family Encyrtidae (Chalcidoidea). Part I. Review of systems of classifications, the subfamily Tetracneminae Howard, 1892. *Ent. Obozr.*, 52 (1): 163—175; Classification of parasitic Hymenoptera of the family Encyrtidae (Chalcidoidea). Part II. The subfamily Encyrtinae Walker, 1837. *Ent. Obozr.*, 52 (2): 416—429.

My sincere thanks to all those who helped in preparing this list, in particular to Dr. Z. Bouček, Dr. J. S. Noyes (London) and Dr. V. A. Tryapitzin (Leningrad) for their critical review of the manuscript, corrections and additional data.

CHALCIDOIDEA			
LEUCOSPIDAE			
<b>Leucospis</b> Fabricius, 1775			
<i>biquetina</i> Jurine, 1807	B M S		
<i>dorsigera</i> Fabricius, 1775	B M S		
<i>gigas</i> Fabricius, 1793	S		
<i>intermedia</i> Illiger, 1807	B S		
CHALCIDIDAE			
DIRHININAE			
<b>Dirhinus</b> Dalman, 1820			
<i>hesperidum</i> (Rossi, 1790)	S		
CHALCIDINAE			
<b>Chalcis</b> Fabricius, 1789			
<i>myrifex</i> (Sulzer, 1776)	B M S		
<i>sispes</i> (Linnaeus, 1761)	B M S		
<b>Spilochalcis</b> Thomson, 1875			
<i>xanthostigma</i> (Dalman, 1820)	B S		
<b>Brachymeria</b> Westwood, 1832			
<i>coloradensis</i> (Cresson, 1872)	M S		
<i>femorata</i> (Panzer, 1801)	B M S		
<i>inermis</i> (Fonscolombe, 1840)	M S		
intermedia (Nees, 1834)	B M S		
<i>minuta</i> (Linnaeus, 1767)	B M S		
<i>moerens</i> (Ruschka, 1922)	S		
<i>podagraca</i> (Fabricius, 1787)	S		
<i>rugulosa</i> (Foerster, 1859)	B M S		
<i>secundaria</i> (Ruschka, 1922)	S		
<i>vitripennis</i> (Foerster, 1859)	B M S		
<i>walkeri</i> Dalla Torre, 1898	B M S		
HALTICHELLINAE			
<b>Belaspidia</b> Masi, 1916			
<i>obscura</i> Masi, 1916	B M S		
<b>Kriechbaumerella</b>			
Dalla Torre, 1897			
<i>hofferi</i> (Bouček, 1852)	S		
<b>Haltichella</b> Spinola, 1811			
<i>rufipes</i> (Olivier, 1790)	B M S		
<b>Neochalcis</b> Kirby, 1883			
<i>fertoni</i> (Kieffer, 1899)	B M S		
<i>osmicida</i> (Saunders, 1873)	M S		
<b>Euchalcis</b> Dufour, 1861			
<i>minor</i> Hoffer, 1951	S		
<b>Hockeria</b> Walker, 1834			
<i>bifasciata</i> Walker, 1834	M S		

<i>hockerioides</i> (Bouček, 1952)	S	<i>calamagrostis</i> (von Schlechtendal, 1891)	B M
<i>hofferi</i> Bouček, 1952	M S	<i>cylindrica</i> (von Schlechtendal, 1891)	B S
<i>inopinata</i> Bouček, 1952	M S	<i>cereipes</i> (Erdös, 1955)	B M S
<i>magna</i> Bouček, 1952	B M S	<i>eximia</i> (Giraud, 1863)	B M S
<i>singularis</i> Bouček, 1952	M	<i>fulvicollis</i> (Walker, 1832)	B M S
<i>susterai</i> Bouček, 1952	B M	<i>giraudi</i> (von Schlechtendal, 1891)	B
<i>unicolor</i> Walker, 1834	B M S	<i>hyalipennis</i> (Walker, 1832)	B M
<b>Hybothorax</b> Ratzeburg, 1844		<i>linearis</i> (Walker, 1832)	B M S
<i>graffi</i> Ratzeburg, 1844	B M S	<i>longicornis</i> (Walker, 1832)	B M
<b>Neohybothorax</b>		<i>phleicola</i> (Hedicke, 1920)	M
<i>Nikolskaja</i> , 1960		<i>phragmitis</i> (Erdös, 1952)	B M
<i>hetera</i> (Walker, 1834)	M S	<i>poae</i> (von Schlechtendal, 1891)	B M
<b>Lasiochalcidia</b> Masi, 1927		<i>scheppigi</i> (von Schlechtendal, 1891)	B M
<i>dargelasi</i> (Latreille, 1806)	B M S	<i>schlechtendali</i> (Hedicke, 1920)	B M
<i>guineensis</i> (Steffan, 1951)	M S	<b>Pseudotetramesa</b> <i>Kalina</i> , 1970	
<b>Proconura</b> Girault, 1915		<i>doksenensis</i> Kalina, 1970	B
<i>nigripes</i> (Walker, 1834)	B S	<b>Sycophila</b> Walker, 1871	
<b>Invreia</b> Masi, 1927		<i>concinna</i> (Boheman, 1836)	B
<i>benoisti</i> Steffan, 1948	B M S	<i>flavicollis</i> (Walker, 1834)	B M S
<i>ligistica</i> Masi, 1934	M S	<i>mayri</i> (Erdös, 1959)	B M S
<i>mirabilis</i> Bouček, 1951	S	<i>scorzonerae</i> (Mayr, 1905)	M S
<i>rufitarsis</i> (Illiger, 1807)	S	<i>stagnalis</i> (Erdös, 1947)	B S
<i>subaenea</i> Masi, 1927	S	<i>variegata</i> (Curtis, 1831)	S
<i>subarmata</i> (Foerster, 1856)	B M S	<b>Eurytoma</b> Illiger, 1807	
EURYTOMIDAE		<i>aciculata</i> Ratzeburg, 1848	B M
RILEYINAE		<i>appendigaster</i> (Swederus, 1795)	B
<b>Archirileya</b> Silvestri, 1920		<i>arctica</i> Thomson, 1876	B M S
<i>inopinata</i> Silvestri, 1920	B M S	<i>atra</i> (Walker, 1832)	B
EURYTOMINAE		<i>brunniventris</i>	
<b>Tetramesa</b> Walker, 1848		Ratzeburg, 1852	B M
<i>aciculata</i>			
(von Schlechtendal, 1891)	B M S		
<i>airae</i>			
(von Schlechtendal, 1891)	M		
<i>angustipennis</i> (Walker, 1832)	B		
<i>brevicollis</i> (Walker, 1836)	M		
<i>brevicornis</i> (Walker, 1832)	B M		

<i>collaris</i> Walker, 1932	B M S	<i>brevicaudis</i> Ratzeburg, 1852	B M
<i>coluteae</i> Bouček, 1954	B	<i>dorsalis</i> (Fabricius, 1798)	B M S
<i>crassinervis</i> Thomson, 1876	B	<i>puctus</i> (Foerster, 1841)	B M S
<i>flavimana</i> Boheman, 1836	B M	<i>rosae</i> Bouček, 1971	M
<i>globiceps</i> Bouček, 1954	B S	<i>spermotrophus</i> Wachtl, 1893	B M S
<i>goidanichi</i> Bouček, 1970	S	<i>stigmatizans</i> (Fabricius, 1798)	M S
<i>gyoerfii</i> Erdős, 1957	S	<i>strobilobius</i> Ratzeburg, 1848	B M
<i>maura</i> Boheman, 1836	B S	<i>suspectus</i> Borries, 1895	B M S
<i>morio</i> Boheman, 1836	B M S	MONODONTOMERINAE	
<i>nodularis</i> Boheman, 1836	B M S		
<i>robusta</i> Mayr, 1878	B M S	<b>Podagrion</b> Spinola, 1811	
<i>rosae</i> Nees, 1834	B M S	<i>pachymerum</i> (Walker, 1833)	M S
<i>rubicola</i> Girault, 1866	B S	<i>splendens</i> Spinola, 1811	S
<i>rufipes</i> Walker, 1832	B	<b>Exopristus</b> Ruschka, 1923	
<i>salicis</i> Walker, 1834	B M	<i>trigonomerus</i> (Masi, 1916)	M S
<i>serratulae</i> (Fabricius, 1798)	B	<b>Cryptopristus</b> Foerster, 1856	
<i>spinipes</i> Kalina, 1970	M S	<i>caliginosus</i> (Walker, 1833)	B M S
<i>squamea</i> Walker, 1834	S	<b>Idarnotorymus</b> Masi, 1916	
<i>stenostigma</i> Thomson, 1876	B M	<i>pulcher</i> Masi, 1916	S
<i>strigifrons</i> Thomson, 1876	B	<b>Eridontomerus</b> Crawford, 1907	
<i>tibialis</i> Boheman, 1836	B M S	<i>arabonicus</i> Erdős, 1954	B M
<i>tilicola</i> Hedquist, 1966	?	<i>biroi</i> Ruschka, 1923	M S
<i>verticillata</i> (Fabricius, 1798)	B M S	<i>isosomatis</i> (Rilley, 1881)	B M S
<b>Ipideurytoma</b> Bouček et Novicky, 1954		<i>laticornis</i> Foérster, 1859)	B S
<i>spessivtsevi</i> Bouček et Novicky, 1954	B M	Monodotomerus Westwood, 1833	
<b>Nikanoria</b> Nikolskaja, 1955		<i>aereus</i> (Walker, 1834)	B M S
<i>metallica</i> (Erdős, 1956)	M	<i>dentipes</i> (Dalman, 1820)	B M S
<b>Systole</b> Walker, 1832		<i>minor</i> (Ratzeburg, 1848)	B M S
<i>albipennis</i> Walker, 1832	B	<i>obscurus</i> Westwood, 1833	B M S
<i>bipunctata</i> Erdős, 1952	S	<i>obsoletus</i> (Fabricius, 1798)	B
<b>Pseudosystole</b> Kalina, 1969		<i>rugulosus</i> (Thomson, 1876)	B
<i>hofferi</i> Kalina, 1969	B	<i>vicicellae</i> (Walker, 1847)	B M S
TORYMIDAE		Glyphomerus Foerster, 1856	
MEGASTIGMINAE			
<b>Megastigmus</b> Dalman, 1820			
<i>aculeatus</i> (Swederus, 1795)	B M S		
<i>bipunctatus</i> (Swederus, 1795)	S		

<i>europaeus</i> (Erdös, 1957)	M	<i>auratus</i> (Geoffroy in Fourcroy, 1785)	B M S
<i>stigma</i> (Fabricius, 1793)	B M S	<i>azureus</i> Boheman, 1834	B M
<i>tibialis</i> Foerster, 1859	B M S	<i>baudysi</i> Bouček, 1954	B M S
<b>Microdontomerus</b> Grawford, 1907		<i>bedeguaris</i> (Linnaeus, 1758)	B M S
<i>coturnatus</i> (Masi, 1921)	M S	<i>bohemani</i> Thomson, 1876	B S
<b>Liodontomerus</b> Gahan, 1914		<i>campanulae</i> (Cameron, 1880)	B
<i>balasi</i> Szelényi, 1957	S	<i>caudatus</i> Boheman, 1834	B M S
<i>mayri</i> (Wachtl, 1883)	B M	<i>chloromerus</i> (Walker, 1834)	B M S
<i>pannonicus</i> (Ruschka, 1923)	S	<i>chrysoccephalus</i> (Boheman, 1834)	B
<i>papaveris</i> Foerster, 1856]	B M	<i>cingulatus</i> Nees, 1834	B M
<i>splendidus</i> Szelényi, 1959	M S	<i>confinis</i> (Walker, 1833)	B
<i>terebrator</i> (Masi, 1916)	S	<i>corni</i> Mayr, 1874	M
<i>variegatus</i> Szelényi, 1959	M	<i>cultiventris</i> Ratzeburg, 1844	B M S
<b>Pseuderimerus</b> Gahan, 1919		<i>cupratus</i> Boheman, 1834	B
<i>luteus</i> Bouček, 1954	B M	<i>cyaneus</i> Boheman, 1834	B M S
<b>Didactyliocerus</b> Masi, 1916		<i>cynipedis</i> Boheman, 1834	B S
<i>dispar</i> Masi, 1916	S	<i>druparum</i> Boheman, 1834	S
<b>Dimeromicrus</b> Ruschka, 1923		<i>erucarum</i> (Schrank, 1871)	B M S
<i>kiesenwetteri</i> (Mayr, 1874)	B M S	<i>fastuosus</i> Boheman, 1834	B M
<b>Pseudotorymus</b> Masi, 1922		<i>formosus</i> (Walker, 1833)	B
<i>apionis</i> (Mayr, 1874)	B M S	<i>fuscicornis</i> (Walker, 1833)	B
<i>brassicae</i> (Ruschka, 1923)	B M S	<i>fuscipes</i> Boheman, 1834	B
<i>leguminum</i> Ruschka, 1923	B	<i>galii</i> Boheman, 1834	M
<i>medicaginis</i> (Mayr, 1974)	B	<i>glechomae</i> Mayr, 1874	B M
<i>militaris</i> (Boheman, 1834)	B M	<i>hederae</i> (Walker, 1833)	B S
<i>salicis</i> Ruschka, 1923	M	<i>heyperi</i> Wachtl, 1883	B M
<i>sapphyrinus</i> (Fonscolombe, 1832)	B M S	<i>igniceps</i> Mayr, 1874	B M
TORYMINAE		<i>incertus</i> Foerster, 1841	S
<b>Diomorus</b> Walker, 1834		<i>insolitus</i> Walker, 1874	B S
<i>armatus</i> Boheman, 1834	B M	<i>juniperi</i> (Linnaeus, 1758)	B M
<i>calcaratus</i> (Nees, 1834)	B M S	<i>millefolii</i> Ruschka, 1931	B M
<i>cupreus</i> (Spinola, 1808)	B M S	<i>nigricornis</i> Boheman, 1834	B M S
<b>Torymus</b> Dalman, 1820		<i>nigritarsus</i> (Walker, 1833)	B
<i>acrophilae</i> Ruschka, 1921	M	<i>nitidulus</i> (Walker, 1833)	B
<i>affinis</i> (Fonscolombe, 1832)	B	<i>nobilis</i> Boheman, 1834	B
<i>artcticus</i> (Thomson, 1876)	M	<i>notatus</i> (Walker, 1833)	B
<i>arundinis</i> (Walker, 1833)	B M S	<i>pulchellus</i> Thomson, 1876	B
		<i>purpurascens</i> (Boheman, 1834)	B
		<i>quercinus</i> Boheman, 1834	B
		<i>roboris</i> (Walker, 1833)	B
		<i>rubi</i> (Schrank, 1781)	B M S

<i>ruschkai</i> (Hoffmeyer, 1930)	M	<i>chrysonotus</i> Foerster, 1859	B M S
<i>saliciperdae</i> Ruschka, 1921	B M	<i>eximius</i> Masi, 1932	B S
<i>scaposus</i> Thomson, 1876	B	<i>intermedius</i> Bouček, 1956	S
<i>schizothecae</i> Ruschka, 1921	B	<i>laevifrons</i> Dalman, 1822	B M S
<i>socius</i> Mayr, 1874	B	<i>maceki</i> Bouček, 1956	B
<i>spilopterus</i> Boheman, 1834	B	<i>masculinus</i> Bouček, 1956	B M S
<i>tanaceticola</i> Ruschka, 1921	M	<i>micans</i> Dalman, 1820	S
<i>thymi</i> Ruschka, 1921	B	<i>neglectus</i> Bouček, 1956	B M S
<i>tipulariarum</i> Zetterstedt, 1838	B M	<i>nitens</i> Walker, 1834	B M S
<i>ulmariae</i> Ruschka, 1921	B M	<i>polypori</i> Bouček, 1971	B S
<i>varians</i> (Walker, 1833)	B M S	<i>ruficornis</i> (Fabricius, 1793)	B M S
<i>ventralis</i> (Fonscolombe, 1832)	B M S	<i>tristis</i> Mayr, 1905	B M S
ORMYRIDAE			
<b>Ormyrus</b> Westwood, 1832		PTEROMALIDAE	
<i>diffinis</i> (Fonscolombe, 1832)	B M S	CLEONYMINAE	
<i>gratiosus</i> (Foerster, 1860)	B M S	<b>Cleonus</b> Latreille, 1809	
<i>nitidulus</i> (Fabricius, 1804)	B M S	<i>brevis</i> Bouček, 1972	S
<i>orientalis</i> Walker, 1871	B M S	<i>laticornis</i> Walker, 1837	B S
<i>papaveris</i> (Perris, 1840)	S	<b>Heydenia</b> Foerster, 1856	
<i>punctiger</i> Westwood, 1832	B M S	<i>pretiosa</i> Foerster, 1856	B M S
<i>speculifer</i> Erdös, 1946	M	<b>Pannoniella</b> (Erdös, 1960)	
<i>wachtli</i> Mayr, 1904	B M S	<i>sexramosa</i> (Erdös, 1946)	B M S
EUCHARITIDAE			
<b>Eucharis</b> Latreille, 1802		<b>Notanisus</b> Walker, 1837	
<i>adscendens</i> (Fabricius, 1787)	B M S	<i>versicolor</i> Walker, 1837	S
<i>dobrogica</i> Andriescu, 1968	S	MACROMESINAE	
<i>hyalinipennis</i> Hoffer et Novicky, 1954	B	<b>Macromesus</b> Walker, 1848	
<b>Stilbula</b> Spinola, 1811		<i>amphiretus</i> Walker, 1848	B S
<i>cyniformis</i> (Rossi, 1792)	M S	CEINAE	
PERILAMPIDAE			
<b>Perilampus</b> Latreille, 1809		<b>Cea</b> Walker, 1837	
<i>aureoviridis</i> Walker, 1833	B S	<i>pulicaris</i> Walker, 1837	B M
<i>aeneus</i> (Rossi, 1790)	B M S	<b>Spalangiopelta</b> Masi, 1922	
<i>auratus</i> (Panzer, 1798)	M S	<i>alata</i> Bouček, 1952	B
<i>cephalotes</i> Bouček, 1956	S	<i>dudichi</i> Erdös, 1956	S
		SPALANGIINAE	
		<b>Spalangia</b> Latreille, 1805	
		<i>cameroni</i> Perkins, 1910	B M
		<i>crassicornis</i> Bouček, 1963	B
		<i>endius</i> Walker, 1839	B
		<i>erythromera</i> Foerster, 1850	B M S
		<i>fuscipes</i> Nees, 1834	B M S
		<i>nigra</i> Latreille, 1805	B M
		<i>nigripes</i> Curtis, 1839	B S

<i>nigroaenea</i> Curtis, 1839	B	M	<i>nidicola</i> Ferrière, 1934	B
<i>rugulosa</i> Foerster, 1850	B		CHRY SOLAMPINAE	
<i>slovaca</i> Bouček, 1963		S	<b>Chrysolampus</b> Spinola, 1811	
<i>subpunctata</i> Foerster, 1850	B	S	<i>dentatus</i> (Bouček, 1956)	B
CEROCEPHALINAE			<i>punctatus</i> (Foerster, 1859)	B M S
<b>Cerocephala</b> Westwood, 1832			<i>rufitarsis</i> (Foerster, 1859)	B M S
<i>cornigera</i> Westwood, 1832	B	S	<i>splendidus</i> (Spinola, 1808)	B S
<i>rufa</i> (Walker, 1833)	B		<i>thenae</i> (Walker, 1848)	B M S
<b>Theocolax</b> Westwood, 1832			<b>Chrysomala</b> Foerster, 1859	
<i>formiciformis</i> Westwood,			<i>roseri</i> Foerster, 1859	B M
1832		B	<b>Brachylatus</b> Hoffer	
DIPARINAE			et Novicky, 1954	
<b>Dipara</b> Walker, 1833			<i>viridis</i> Hoffer	
<i>petiolata</i> Walker, 1833		B M S	et Novicky, 1954	M
NETOMOCERA			PANSTENONINAE	
<b>Netomocera</b> Bouček, 1954			<b>Panstenon</b> Walker, 1846	
<i>setifera</i> Bouček, 1954		B M S	<i>agylla</i> (Walker, 1850)	B
NEODIPARINAE			<i>oxylus</i> Walker, 1834	B M S
<b>Neodipara</b> Erdös, 1955			MISCOGASTERINAE	
<i>masneri</i> Bouček, 1961		B	<b>Trigonoderus</b> Westwood,	
EUNOTINAE			1832	
<b>Eunotus</b> Walker, 1834			<i>cyanescens</i> (Foerster,	
<i>acutus</i> Kurdjumov, 1912	B	M	1841)	B M S
<i>areolatus</i> (Ratzeburg,	B	M	<i>filatus</i> Walker, 1836	M S
1852)			<i>princeps</i> Westwood, 1832	B M S
<i>cretaceus</i> Walker, 1834	B	M S	<i>pulcher</i> Walker, 1836	B M S
<i>hofferi</i> Bouček, 1972	B	M S	<b>Plutothrix</b> Foerster, 1856	
<i>nigriclavus</i> (Foerster,	B	M	<i>cisae</i> Hedqvist, 1967	B M S
1856)			<i>scenicus</i> (Walker, 1836)	B M S
<i>obscurus</i> Masi, 1931		S	<i>trifasciatus</i> (Thomson,	
<i>parvulus</i> Masi, 1931	B	S	1878)	B M S
Scutellista Motschulsky,			<b>Jansonielia</b> Kerrich, 1957	
1859			<i>ambigua</i> Graham, 1969	B
<i>cyanea</i> Motschulsky, 1859	M		<i>caudata</i> Kerrich, 1957	B
<i>obscura</i> (Foerster, 1878)	M	S	<b>Platygerrhus</b> Thomson, 1878	
ASAPHINAE			<i>affinis</i> (Walker, 1836)	B M
<b>Asaphes</b> Walker, 1834			<i>dolosus</i> (Walker, 1836)	B
<i>suspensus</i> (Nees, 1834)	B	M S	<i>ductilis</i> (Walker, 1836)	B M S
<i>vulgaris</i> Walker, 1834	B	M S	<i>longigena</i> Graham, 1969	B S
<b>Hyperimerus</b> Girault, 1917			<i>maculatus</i> Erdös, 1957	B
<i>pusillus</i> (Walker, 1833)	B		<i>unicolor</i> Graham, 1969	B M
Bairamlia Waterston, 1929			Gastracanthus Westwood,	
			1833	

<b>pulcherrimus</b> Westwood, 1833	B M S	<b>polita</b> (Walker, 1834)	B M
<b>Erdoesia</b> Bouček, 1957		<b>Halticopterina</b> Erdös, 1946	
<b>tessellata</b> Bouček, 1957	B M	<i>moczari</i> Erdös, 1954	S
<b>Sphegigaster</b> Spinola, 1811		<i>triannulata</i> Erdös, 1946	B S
<i>aculeata</i> (Walker, 1833)	B	<b>Thinodytes</b> Graham, 1956	
<i>intersita</i> Graham, 1969	B	<i>cyzius</i> (Walker, 1839)	B M S
<i>nigricornis</i> (Nees, 1834)	B M	<b>Tricyclomischus</b> Graham, 1956	
<i>pallicornis</i> (Spinola, 1808)	B M S	<i>celticus</i> Graham, 1956	B
<i>stepicola</i> Bouček, 1965	B M S	<b>Rhincocoelia</b> Graham, 1956	
<i>truncata</i> Thomson, 1878	B M S	<i>constans</i> (Walker, 1836)	B M
<b>Syntomopus</b> Walker, 1833		<i>coretas</i> (Walker, 1848)	B
<i>incisus</i> Thomson, 1878	B M S	<i>grahami</i> Bouček, 1970	B
<i>incurvus</i> Walker, 1833	B M	<b>Merismus</b> Walker, 1833	
<i>oviceps</i> Thomson, 1878	B M S	<i>megapterus</i> Walker, 1833	B M
<i>thoracicicus</i> Walker, 1833	B S	<i>nitidus</i> (Walker, 1833)	B
<b>Notoglyptus</b> Masi, 1917		<i>rufipes</i> Walker, 1833	B M
<i>niger</i> Masi, 1917	B S	<b>Callimerismus</b> Graham, 1956	
<b>Cryptoprymna</b> Foerster, 1856		<i>suecicus</i> Graham, 1969	B
<i>atra</i> (Walker, 1833)	B S	<b>Ksenoplata</b> Bouček, 1965	
<b>Novitzkyanus</b> Bouček, 1961		<i>quadrata</i> Bouček, 1965	B S
<i>cryptogaster</i> Bouček, 1961	S	<b>Susterai</b> Bouček, 1972	
<b>Cyrtogaster</b> Walker, 1833		<i>acerina</i> Bouček, 1972	B
<i>vulgaris</i> Walker, 1833	B M S	<b>Nodisoplata</b> Graham, 1969	
<b>Polycystus</b> Westwood, 1839		<i>diffinis</i> (Walker, 1874)	B
<i>clavicornis</i> (Walker, 1833)	B M S	<b>Seladerma</b> Walker, 1834	
<b>Toxeumia</b> Walker, 1833		<i>berani</i> (Delucchi, 1953)	B
<i>acilius</i> (Walker, 1848)	B	<i>coeruleovirens</i>	
<i>funcicorne</i> Walker, 1833	B M	(Foerster, 1861)	M
<i>substruncatum</i> Graham, 1959	M	<i>diffine</i> (Walker, 1833)	B
<b>Halticoptera</b> Spinola, 1811		<i>geniculatum</i> (Zetterstedt, 1838)	B M S
<i>aenea</i> (Walker, 1833)	B	<i>laetum</i> Walker, 1834	B
<i>brevicornis</i> Thomson, 1876	B S	<i>sabbas</i> (Walker, 1848)	B
<i>circulus</i> (Walker, 1833)	B M S	<i>saurus</i> Walker, 1844	B
<i>collaris</i> (Walker, 1836)	B	<i>simplex</i> (Thomson, 1876)	B
<i>crius</i> Walker, 1839	B M	<i>tarsale</i> (Walker, 1833)	B
<i>flavicornis</i> (Spinola, 1808)	B M S	<b>Thektogaster</b> Delucchi, 1955	
<i>hippeus</i> (Walker, 1839)	B M	<i>chrysia</i> (Foerster, 1861)	M
<i>laevigata</i> Thomson, 1876	B M	<b>Glyphognathus</b> Graham, 1956	
<i>patellana</i> (Dalman, 1818)	B	<i>flameus</i> (Delucchi, 1953)	B
		<i>umbelliferae</i> Graham, 1956	B
		<b>Sphaeripalpus</b> Foerster, 1841	

<i>fuscipes</i> (Walker, 1833)	B	<i>parvula</i> Thomson, 1876	B
<i>laevis</i> (Delucchi, 1953)	B	<b>Bugacia</b> Erdős, 1946	
<i>laevigatus</i> (Delucchi, 1953)	B	<i>arenaria</i> Erdős, 1946	B S
<i>microstolus</i> (Graham, 1969)	B	<i>submontana</i> Bouček, 1955	B
<i>viridis</i> Foerster, 1841	B	<b>Melancistrus</b> Graham, 1969	
<b>Stictomischus</b> Thomson, 1876		<i>mucronatus</i> (Thomson, 1876)	B
<i>obscurus</i> (Walker, 1833)	B	<b>Gastrancistrus</b> Westwood, 1833	
<i>tumidus</i> (Walker, 1833)	B	<i>acutus</i> Walker, 1834	B
<b>Miscogaster</b> Walker, 1833		<i>citripes</i> (Thomson, 1876)	B
<i>elegans</i> Walker, 1833	B	<i>clavatus</i> (Thomson, 1876)	B
<i>maculata</i> Walker, 1833	B M	<i>consors</i> Graham, 1969	B
<i>rufipes</i> Walker, 1833	B M	<i>coxalis</i> (Thomson, 1876)	B
<b>Lamprotatus</b> Westwood, 1833		<i>dispar</i> Graham, 1969	B
<i>annularis</i> (Walker, 1833)	B	<i>fuscicornis</i> Walker, 1834	B S
<i>crassipes</i> Thomson, 1876	B	<i>glabellus</i> (Thomson, 1876)	B
<i>cupreus</i> Delucchi, 1953	M	<i>hamillus</i> Walker, 1848	B
<i>piceinervis</i> Thomson, 1876	B	<i>pusztensis</i> (Erdős, 1946)	B
<i>pschorni</i> Delucchi, 1953	B	<i>puncticollis</i> (Thomson, 1876)	B
<i>simillimus</i> Delucchi, 1953	B	<i>salicis</i> (Nees, 1834)	B M S
<i>splendens</i> Westwood, 1833	B M	<i>vagans</i> Westwood, 1833	B
<b>Skleroceras</b> Delucchi, 1953		<i>vulgaris</i> Walker, 1834	B
<i>clavigerum</i> (Thomson, 1878)	B S	<b>Ecrizotes</b> Foerster, 1861	
<i>socium</i> (Zetterstedt, 1838)	S	<i>filicornis</i> (Thomson, 1876)	B
<b>Micradelus</b> Walker, 1834		<i>longicornis</i> (Walker, 1848)	B
<i>acutus</i> Graham, 1969	B M	<i>monticola</i> (Foerster, 1861)	B
<i>rotundus</i> Walker, 1834	B M S	<b>Spathopus</b> Ashmead, 1904	
<b>Termolampa</b> Bouček, 1961		<i>hofferi</i> Bouček, 1964	S
<i>pinicola</i> Bouček, 1961	B S	<b>Stenophrus</b> Foerster, 1841	
<b>Ormocerus</b> Walker, 1834		<i>compresus</i> Foerster, 1841	B
<i>vernalis</i> Walker, 1834	B M S	<b>Pirene</b> Haliday, 1833	
<b>Ammeia</b> Delucchi, 1962		<i>bouceki</i> Graham, 1969	B
<i>pulchella</i> Delucchi, 1962	M S	<i>chalybea</i> Haliday, 1833	B M
<b>Semiotelus</b> Westwood, 1839		<i>conjugens</i> Graham, 1969	B
<i>diversus</i> (Walker, 1834)	B M S	<i>penetrans</i> (Kirby, 1800)	B
<i>mundus</i> (Walker, 1834)	B M S	<i>varicornis</i> Haliday, 1833	B S
<b>Systasis</b> Walker, 1834		PTEROMALINAE	
<i>annulipes</i> (Walker, 1834)	B M S	<b>Dinotiscus</b> Ghesquière, 1946	
<i>encyrtoides</i> Walker, 1834	B M S	<i>aponius</i> (Walker, 1848)	B S
<i>longula</i> Bouček, 1955	M	<i>colon</i> (Linnaeus, 1758)	B S
		<i>eupterus</i> (Walker, 1836)	B M S

<b>Rhopalicus</b> Foerster, 1856			
<i>guttatus</i> (Ratzeburg, 1844)	M		
<i>quadratus</i> (Ratzeburg, 1844)	B S		
<i>tutela</i> (Walker, 1836)	B M S		
<b>Acrocormus</b> Foerster, 1856			
<i>semifasciatus</i> Thomson, 1878	B S		
<b>Cheiropachus</b> Westwood, 1828			
<i>quadrum</i> (Fabricius, 1787)	B M S		
<b>Metacolus</b> Foerster, 1856			
<i>azureus</i> (Ratzeburg, 1844)	B		
<i>unifasciatus</i> Foerster, 1856	B S		
<b>Raphitellus</b> Walker, 1834			
<i>maculatus</i> Walker, 1834	B M S		
<i>ladenbergi</i> (Ratzeburg, 1844)	B		
<b>Panelus</b> Foerster, 1856			
<i>flavipes</i> (Foerster, 1841)	B		
<b>Roptrocerus</b> Ratzeburg, 1848			
<i>brevicornis</i> Thomson, 1878	B M S		
<i>mirus</i> (Walker, 1834)	B S		
<i>xylophagorum</i> (Ratzeburg, 1844)	B M S		
<b>Xiphydriophagus</b> Ferrière, 1952			
<i>meyerinckii</i> (Ratzeburg, 1848)	M		
<b>Habritys</b> Thomson, 1878			
<i>brevicornis</i> (Ratzeburg, 1844)	B M S		
<b>Perniphora</b> Ruschka, 1923			
<i>robusta</i> Ruschka, 1923	B M S		
<b>Caenocrepis</b> Thomson, 1878			
<i>arenicola</i> Thomson, 1878	B M S		
<b>Oxysuchus</b> Delucchi, 1956			
<i>pilosulus</i> (Thomson, 1878)	B		
<b>Ischyroptyx</b> Delucchi, 1956			
<i>ligusticus</i> (Masi, 1921)	S		
<b>Anisopteromalus</b> Ruschka, 1912			
		<i>calandrae</i> (Howard, 1881)	B M S
		<b>Dinarmus</b> Thomson, 1878	
		<i>acutus</i> Thomson, 1878	B M S
		<b>Dinarmoides</b> Masi, 1924	
		<i>spilopterus</i> Masi, 1924	M S
		<b>Stenoselma</b> Delucchi, 1956	
		<i>nigrum</i> Delucchi, 1956	B M S
		<b>Norbanus</b> Walker, 1843	
		<i>meridionalis</i> (Masi, 1922)	S
		<i>scabriculus</i> (Nees, 1834)	B M S
		<b>Picroscytoides</b> Masi, 1922	
		<i>albiventris</i> Bouček, 1970	M S
		<i>cerasiops</i> Masi, 1922	B S
		<b>Merisus</b> Walker, 1835	
		<i>splendidus</i> Walker, 1835	B M S
		<b>Homoporus</b> Thomson, 1878	
		<i>aphareetus</i> (Walker, 1839)	B
		<i>arestor</i> (Walker, 1848)	B M S
		<i>auratus</i> Erdős, 1953	B M
		<i>biroi</i> Szélényi, 1957	M S
		<i>budensis</i> Erdős, 1953	S
		<i>brunneiventris</i> Szélényi, 1956	B
		<i>cephalotes</i> Szélényi, 1957	S
		<i>chalcidiphagus</i> (Walsh & Riley, 1936)	B M S
		<i>cupreus</i> Erdős, 1953	B S
		<i>destructor</i> (Say, 1817)	B S
		<i>febriculosus</i> (Girault, 1917)	B
		<i>fulviventris</i> (Walker, 1835)	B M S
		<i>gibbiscuta</i> (Thomson, 1878)	B
		<i>gusztavi</i> Bouček, 1974	M
		<i>laeviusculus</i> Erdős, 1953	B S
		<i>luniger</i> (Nees, 1834)	B M S
		<i>pulchripes</i> Erdős, 1953	B M S
		<i>sashegyensis</i> Erdős, 1953	B M S
		<i>semiluteus</i> (Walker, 1872)	B M S
		<i>subniger</i> (Walker, 1835)	B M S
		<i>titanes</i> Szélényi, 1957	B S
		<b>Callitula</b> Spinola, 1811	
		<i>angioneurae</i> Bouček, 1970	M
		<i>bicolor</i> Spinola, 1811	B M S
		<i>elongata</i> (Thomson, 1878)	B

<i>jerrierei</i> Bouček, 1964	B M S	<i>bifoveolatus</i> Foerster, 1861	M S
<i>pyrrhogaster</i> (Walker, 1833)	B M S	<i>capreae</i> (Linnaeus, 1761) B	S
<b>Psilocera</b> Walker, 1833	B M S	<i>cardui</i> (Erdös, 1953)	S
<i>atra</i> (Walker, 1834)	B M S	<i>chrysos</i> Walker, 1836	S
<i>crassispina</i> (Thomson, 1878)	B M S	<i>cioni</i> (Thomson, 1878)	B
<i>obscura</i> (Walker, 1833)	B M S	<i>cionobius</i> (Erdös, 1953)	B
<b>Catolaccus</b> Thomson, 1878	B M S	<i>conopidarum</i> (Bouček, 1961)	B
<i>ater</i> (Ratzeburg, 1852)	B M S	<i>crassicornis</i> Zetterstedt, 1838	B
<i>crassiceps</i> (Masi, 1911)	S	<i>crassinervis</i> (Thomson, 1878)	B
<b>Erythromalus</b> Graham, 1956		<i>decipiens</i> (Graham, 1969)	B
<i>nubilipennis</i> (Walker, 1835)	B S	<i>dispar</i> (Curtis, 1827)	B S
<i>rufiventris</i> (Walker, 1835)	B S	<i>dolichurus</i> (Thomson, 1878)	B
<b>Psychophagus</b> Mayr, 1904		<i>elevatus</i> (Walker, 1834)	B
<i>omnivorus</i> (Walker, 1835)	B M S	<i>grandis</i> Walker, 1835	B
<b>Stenetra</b> Masi, 1931		<i>hieracii</i> (Thomson, 1878)	B
<i>ligustica</i> Masi, 1931	B	<i>isarchus</i> Walker, 1839	S
<b>Eurydinota</b> Foerster, 1878		<i>musaeus</i> Walker, 1844	B
<i>leptomera</i> Foerster, 1878	S	<i>pallidicola</i> Bouček, 1972	B
<b>Capellia</b> Delucchi, 1958		<i>parietinae</i> (Graham, 1969)	B
<i>cecidomyidae</i> (Ratzeburg, 1844)	B	<i>platyphilus</i> Walker, 1874	B M
<i>orneus</i> (Walker, 1839)	B M S	<i>puparum</i> (Linnaeus, 1758)	B M S
<b>Mokrzeckia</b> Mokrzecki, 1933		<i>semotus</i> (Walker, 1834)	B M S
<i>pini</i> (Hartig, 1838)	B S	<i>sequester</i> Walker, 1835	B S
<b>Spilomalus</b> Graham, 1956		<i>tereus</i> Walker, 1839	B
<i>quadrinota</i> (Walker, 1835)	B S	<i>venustus</i> Walker, 1835	B S
<b>Spintherus</b> Thomson, 1878		<i>vibulenus</i> (Walker, 1839)	M
<i>dubius</i> (Nees, 1841)	B M S	<b>Tricolas</b> Bouček, 1967	
<b>Sceptrothelys</b> Graham, 1956		<i>xylocleptis</i> Bouček, 1967	B
<i>deione</i> (Walker, 1839)	B S	<b>Cecidostiba</b> Thomson, 1878	
<i>gradiclava</i> (Walker, 1835)	B	<i>adana</i> Askew, 1961	S
<b>Pteromalus</b> Swederus, 1795		<i>geganius</i> (Walker, 1848)	S
<i>albibipennis</i> (Walker, 1835)	B M S	<i>hilaris</i> (Walker, 1836)	B M S
<i>altus</i> (Walker, 1834)	B M	<i>semifascia</i> (Walker, 1835)	B M S
<i>aureolus</i> (Thomson, 1878)	B S	<b>Hobbya</b> Delucchi, 1957	
<i>bedeguaris</i> (Thomson, 1878)	B M S	<i>stenonota</i> (Ratzeburg, 1848)	B
<i>berylli</i> Walker, 1835	B M S	<b>Caenacis</b> Foerster, 1856	
		<i>inflexa</i> (Walker, 1848)	B
		<i>lauta</i> (Walker, 1835)	B M S
		<b>Dinotoides</b> Bouček, 1957	
		<i>tenebricus</i> (Walker, 1834)	B

<b>Ablaxia</b> Delucchi, 1957			<i>favorinus</i> (Walker, 1839)	B
<i>anaxenor</i> (Walker, 1845)	M S		<i>fervida</i> Graham, 1956	B
<i>crasicornis</i> (Thomson, 1878)	B		<i>gracilis</i> (Walker, 1834)	B M S
<i>parviclava</i> (Thomson, 1878)	B S		<i>illudens</i> (Walker, 1836)	B
<i>squamifera</i> (Thomson, 1878)	M		<i>laticeps</i> (Walker, 1848)	B M
<b>Aggelma</b> Delucchi, 1956			<i>liparae</i> (Giraud, 1863)	B M
<i>abdominalis</i> Delucchi, 1956	M		<i>micans</i> (Olivier, 1813)	B
<i>violacea</i> (Zetterstedt, 1838)	B		<i>oxygyne</i> (Walker, 1835)	B
<b>Apelioma</b> Delucchi, 1956			<b>Chlorocytus</b> Graham, 1956	
<i>pteromalimum</i> (Thomson, 1878)	B S		<i>breviscapus</i> Graham, 1965	B
<b>Synedrus</b> Graham, 1956			<i>deschampsiae</i> Graham, 1965	
<i>transiens</i> (Walker, 1835)	B		<i>diversus</i> (Walker, 1836)	B S
<b>Holcaeus</b> Thomson, 1878			<i>formosus</i> (Walker, 1835)	B
<i>calligetus</i> (Walker, 1839)	B		<i>harmolitae</i> Bouček, 1957	B M
<i>compressus</i> (Walker, 1836)	B M S		<i>laogore</i> (Walker, 1836)	B
<i>gorgasus</i> (Walker, 1839)	B M		<i>longicauda</i> (Thomson, 1878)	B
<i>stenogaster</i> (Walker, 1836)	S		<i>phalaridis</i> Graham, 1965	B
<i>stylatus</i> Graham, 1969	B		<i>pilosus</i> Graham, 1965	B
<i>varro</i> (Walker, 1839)	B		<i>pulchripes</i> (Walker, 1834)	B
<b>Cricellius</b> Thomson, 1878			<i>spicatus</i> (Walker, 1835)	B
<i>decipiens</i> Thomson, 1878	B		<b>Isocyrthus</b> Walker, 1833	
<i>gracilentus</i> (Bouček, 1954)	B M S		<i>laetus</i> Walker, 1833	B M
<i>gracilis</i> (Walker, 1836)	B		<b>Psilonotus</b> Walker, 1834	
<b>Trychnosoma</b> Graham, 1957			<i>achaeus</i> Walker, 1848	B
<i>punctipleura</i> (Thomson, 1878)	B S		<i>adamas</i> Walker, 1834	B
<b>Lonchetron</b> Graham, 1956			<b>Anogmus</b> Foerster, 1856	
<i>fennicum</i> Graham, 1956	B		<i>hohenheimensis</i> (Ratzeburg, 1844)	B M S
<b>Kaleva</b> Graham, 1957			<i>piceae</i> (Ruschka, 1922)	B
<i>corynocera</i> Graham, 1957	B		<i>strobilarum</i> (Thomson, 1878)	B M
<b>Eulonchetron</b> Graham, 1966			<i>vala</i> (Walker, 1839)	B M S
<i>scalptrum</i> (Askew, 1962)	B		<b>Karpinskiella</b> Bouček, 1955	
<b>Stenomalina</b> Ghesquière, 1946			<i>pityophthori</i> Bouček, 1955	B
<i>continua</i> (Walker, 1836)	B		<b>Eumacepolus</b> Graham, 1957	
<i>dives</i> (Walker, 1835)	B		<i>grahami</i> von Rosen, 1960	B S
<i>epistena</i> (Walker, 1835)	B M S		<b>Mesopolobus</b> Westwood, 1833	
			<i>aequus</i> (Walker, 1834)	B M S
			<i>agropyricola</i> von Rosen, 1960	B
			<i>albitarsus</i> (Walker, 1834)	B S
			<i>amaenus</i> (Walker, 1834)	B M

<i>anogmoides</i> Graham, 1969	B	S	<i>crassiceps</i> Bouček, 1961	S
<i>aspilus</i> (Walker, 1835)	B		<b>Rohatina</b> Bouček, 1954	
<i>diffinis</i> (Walker, 1834)	B	S	<i>inermis</i> Bouček, 1954	B M
<i>dubius</i> (Walker, 1834)	B M		<i>monstrosa</i> Bouček, 1954	B S
<i>fasciiventris</i> Westwood, 1833	B M		<b>Peridesmia</b> Foerster, 1856	
<i>fuscipes</i> (Walker, 1834)	B		<i>congrua</i> (Walker, 1836)	B S
<i>incultus</i> (Walker, 1834)	B	S	<i>discus</i> (Walker, 1835)	B S
<i>laticornis</i> (Walker, 1834)	B M		<i>montana</i> Bouček, 1972	B S
<i>maculicornis</i> (Giraud, 1863)	B		<b>Spaniopus</b> Walker, 1833	
<i>mediterraneus</i> (Mayr, 1903)		S	<i>amoenus</i> Foerster, 1856	B S
<i>morys</i> (Walker, 1848)	B M S		<i>dissimilis</i> Walker, 1833	B
<i>nobilis</i> (Walker, 1834)	B M S		<i>fulvicornis</i> Bouček, 1972	B S
<i>subfumatus</i> (Ratzeburg, 1852)	B M S		<i>peisonis</i> (Erdős, 1957)	B M S
<i>squamifer</i> (Thomson, 1878)	B	S	<i>varicornis</i> Bouček, 1972	B S
<i>teliformis</i> (Walker, 1834)	B M S		<b>Trichomalus</b> Thomson, 1878	
<i>tibialis</i> (Westwood, 1833)	B M S		<i>acuminatus</i> Delucchi et Graham, 1956	M
<i>typographi</i> (Ruschka, 1924)	B M S		<i>annulatus</i> (Foerster, 1841)	B M S
<i>zetterstedti</i> (Dalla Torre, 1898)		M	<i>campestris</i> (Walker, 1834)	B M S
<b>Meraporus</b> Walker, 1834			<i>flagellaris</i> Graham, 1969	B
<i>graminicola</i> Walker, 1834	B M S		<i>fulvipes</i> (Walker, 1836)	B S
<b>Pegopus</b> Foerster, 1856			<i>gynetus</i> (Walker, 1835)	B S
<i>inornatus</i> (Walker, 1834)	B		<i>helvipes</i> (Walker, 1834)	B
<i>leptomerus</i> Graham, 1969	B		<i>inops</i> (Walker, 1835)	B
<b>Leptomerapus</b> Graham, 1957			<i>inscitus</i> (Walker, 1835)	B
<i>nicaee</i> (Walker, 1839)	B		<i>lepidus</i> (Foerster, 1841)	B
<b>Lampoterma</b> Graham, 1956			<i>lonchaeae</i> Bouček, 1959	B S
<i>viride</i> (Thomson, 1878)	B		<i>nanus</i> (Walker, 1836)	B M S
<b>Stinoplus</b> Thomson, 1878			<i>perfectus</i> (Walker, 1835)	B M S
<i>militaris</i> Thomson, 1878	B		<i>posticus</i> (Walker, 1834)	B
<i>pervasus</i> (Walker, 1836)	B		<i>repandus</i> (Walker, 1835)	B
<b>Pseudocatolaccus</b> Masi, 1908			<i>robustus</i> (Walker, 1835)	B S
<i>nitescens</i> (Walker, 1834)	B M S		<i>rufinus</i> (Walker, 1835)	B S
<b>Apsilocera</b> Bouček, 1955			<i>rugosus</i> Delucchi et Graham, 1956	B
<i>verticilata</i> Bouček, 1955		S	<i>statutus</i> (Foerster, 1841)	B M
<b>Heteroprymna</b> Graham, 1956			<b>Atrichomalus</b> Graham, 1956	
<i>longicornis</i> (Walker, 1835)	B		<i>triannelatus</i> Graham, 1956	B S
<b>Ghelcia</b> Bouček, 1961			<b>Platypieromalus</b> Bouček, 1955	
			<i>pragensis</i> Bouček, 1955	B S
			<b>Trichomalopsis</b> Crawford, 1913	
			<i>acuminatus</i> (Graham, 1969)	B
			<i>albopilosa</i> (Graham, 1969)	S
			<i>arzoneae</i> (Bouček, 1970)	B

<i>exiqua</i> (Walker, 1834)	B		<b>Tritneptis</b> Girault, 1908	
<i>genalis</i> (Graham, 1969)	B		<i>klugii</i> (Ratzeburg, 1844)	B
<i>hemiptera</i> (Walker, 1836)	B		<i>lophyrorum</i> (Ruschka, 1915)	B M
<i>microptera</i> (Lindeman, 1887)	B		<b>Dibrachys</b> Foerster, 1913	
<i>peregrina</i> (Graham, 1969)	B M S		<i>affinis</i> Masi, 1907	B
<i>submarginata</i> (Thomson, 1878)	B S		<i>boarmiae</i> (Walker, 1863)	B M S
<b>Gyrinophagus</b> Ruschka, 1914			<i>braconidis</i> (Ferrière et Faure, 1925)	S
<i>aper</i> (Walker, 1839)	B		<i>cavus</i> (Walker, 1835)	B M S
<b>Nasonia</b> Ashmead, 1904			<i>hians</i> Bouček, 1965	B
<i>vitripennis</i> (Walker, 1836)	B M S		<i>maculipennis</i> Szelényi, 1957	S
<b>Urolepis</b> Walker, 1846			<b>Dibrachoides</b> Kurdjumov, 1913	
<i>maritima</i> (Walker, 1834)	B		<i>dynastes</i> (Foerster, 1841)	B M S
<b>Rakosina</b> Bouček, 1955			<b>Schizonotus</b> Ratzeburg, 1852	
<i>deplanata</i> Bouček, 1955	S		<i>latus</i> (Walker, 1833)	S
<b>Halomalus</b> Erdös, 1953			<i>sieboldi</i> (Ratzeburg, 1852)	B S
<i>crucifer</i> Erdös, 1953	S		<b>Coelopisthia</b> Foerster, 1856	
<b>Pezilepsis</b> Delucchi, 1955			<i>extenta</i> (Walker, 1835)	B M S
<i>dentifera</i> (Thomson, 1878)	B		<b>Conomorium</b> Masi, 1924	
<b>Diglochis</b> Foerster, 1856			<i>patulum</i> (Walker, 1835)	B M S
<i>sylvicola</i> (Walker, 1835)	B M S		<b>Stichocrepis</b> Foerster, 1878	
<b>Tomicobia</b> Ashmead, 1899			<i>armata</i> Foerster, 1878	B M S
<i>promulus</i> (Walker, 1840)	B M S		<b>Muscidifurax</b> Girault et Saunders, 1910	
<i>seitneri</i> (Ruschka, 1924)	B S		<i>raptor</i> Girault et Saunders, 1910	B M S
<b>Dirhicnus</b> Thomson, 1878			<b>Dimachus</b> Thomson, 1878	
<i>pirus</i> (Walker, 1839)	B M S		<i>discolor</i> (Walker, 1836)	B
<b>Arthrolytus</b> Thomson, 1878			<b>Lariophagus</b> Crawford, 1909	
<i>discoideus</i> (Nees, 1834)	B		<i>distinguendus</i> (Foerster, 1841)	B M S
<i>glandium</i> Bouček, 1967	S		<i>fimbriatus</i> Bouček, 1965	S
<i>maculipennis</i> (Walker, 1836)	B		<b>Hemitrichus</b> Thomson, 1878	
<i>ocellus</i> (Walker, 1834)	B		<i>oxygaster</i> Bouček, 1965	S
<i>slovacus</i> Graham, 1969	M S		<i>seniculus</i> (Nees, 1834)	B M S
<b>Erdoesina</b> Graham, 1957			<b>Veltrusia</b> Bouček, 1972	
<i>albomaculata</i> (Ratzeburg, 1852)	B S		<i>rara</i> Bouček, 1972	B
<b>Cyclogastrella</b> Bukowski, 1838			<b>Strejcekia</b> Bouček, 1972	
<i>clypealis</i> Bouček, 1965	B M S		<i>brevior</i> Bouček, 1972	B
<i>deplanata</i> (Nees, 1834)	B M S			
<b>Platneptis</b> Bouček, 1961				
<i>laeta</i> (Walker, 1848)	B			

<i>elegans</i> Bouček, 1972	B	<i>viridis</i> Masi, 1926	M S
<b>Vrestovia</b> Bouček, 1961		<b>Eusandalum</b> Ratzeburg, 1852	
<i>fidenas</i> (Walker, 1848)	B S	<i>coronatum</i> (Thomson, 1876)	B M
<b>Metastenus</b> Walker, 1834		<i>dezorti</i> Bouček, 1967	M
<i>concinnus</i> Walker, 1834	B M S	<i>inerme</i> (Ratzeburg, 1848)	B M S
<b>Pachyneuron</b> Walker, 1833		<i>merceti</i> (Bolivar y Pieltain, 1926)	B S
<i>aphidis</i> (Bouché, 1834)	B M S	<i>walkeri</i> Curtis, 1836	B
<i>gibbiscuta</i> Thomson, 1878	B		
<i>grande</i> Thomson, 1878	B S		
<i>groenlandicum</i> (Holmgren, 1872)	B M S		
<i>leucopiscida</i> Mani, 1939	B S		
<i>muscarium</i> (Linnaeus, 1758)	B M S		
<i>planiscuta</i> Thomson, 1878	B		
<i>solitarium</i> (Hartig, 1838)	B		
<i>vitudurens</i> Delucchi, 1955	B		
<b>Euneura</b> Walker, 1844			
<i>augaurus</i> Walker, 1844	B M S		
<i>lachni</i> Ashmead, 1887	B		
<b>Coruna</b> Walker, 1833			
<i>clavata</i> Walker, 1833	B M S		
<b>Pachycrepoideus</b> Ashmead, 1904			
<i>vindemiae</i> (Rondani, 1875)	B M S		
CRATOMINAE			
<b>Cratomus</b> Dalman, 1820			
<i>megacephalus</i> (Fabricius, 1793)	B		
COLOTRECHNINAE			
<b>Colotrechus</b> Thomson, 1878			
<i>subcoeruleus</i> Thomson, 1878	B M S		
<i>viridis</i> (Masi, 1921)	B S		
EUPELMIDAE			
CALOSOTINAE			
<b>Calosota</b> Curtis, 1836			
<i>acron</i> (Walker, 1848)	B	<i>Anastatus</i> Motschulsky, 1859	
<i>agrili</i> Nikolskaja, 1952	S	<i>bifasciatus</i> (Geoffroy in Fourcroy, 1785)	B M S
<i>aestivalis</i> Curtis, 1836	B M S	<i>giraudi</i> (Ruschka, 1921)	B M S
<i>lixobia</i> Erdös, 1946	S	<i>japonicus</i> Ashmead, 1904	S
<i>obscura</i> Ruschka, 1921	B M S	<i>oscari</i> (Ruthe, 1859)	B M S
<i>vernalis</i> Curtis, 1836	B M	<i>ruficaudus</i> Ferrière, 1954	S
		<b>Metapelma</b> Westwood, 1835	

<i>nobile</i> (Foerster, 1860)	M S	<i>securicornis</i> Domenichini, 1953	B M
<b>Merostenus</b> Walker, 1837		<i>singularis</i> Hoffer, 1953	M
<i>excavatus</i> (Dalman, 1820)	B M S		
ENCYRTIDAE			
TETRACNEMINAE			
<b>Charitopus</b> Foerster, 1856		<b>Paraenasioidea</b> Hoffer, 1953	
<i>fulviventris</i> Foerster, 1860	B M S	<i>marani</i> Hoffer, 1953	S
<i>obscurus</i> (Erdös, 1946)	B M S	<b>Leptomastix</b> Foerster, 1856	
<b>Moraviella</b> Hoffer, 1954		<i>epona</i> (Walker, 1844)	M
<i>inexpectata</i> Hoffer, 1954	M	<i>histrio</i> Mayr, 1876	B M S
<b>Ericydnus</b> Walker, 1837		<b>Rhopus</b> Foerster, 1856	
<i>longicornis</i> (Dalman, 1820)	M	<i>budensis</i> Erdös, 1957	S
<i>sipylus</i> (Walker, 1837)	M	<i>flavidus</i> (Mercet, 1921)	M
<i>strigosus</i> (Nees, 1834)	B M	<i>piso</i> (Walker, 1838)	M
<i>ventralis</i> (Dalman, 1820)	M	<b>Doliphoceras</b> Mercet, 1921	
<b>Lyka</b> Mercet, 1921		<i>belibus</i> (Walker, 1837)	B M
<i>submetallica</i> Mercet, 1921	B M S	<b>Platyrhopus</b> Erdös, 1955	
<b>Dinocarsis</b> Foerster, 1856		<i>meridionalis</i> Ferrière, 1955	S
<i>hemiptera</i> (Dalman, 1820)	B M S	<b>Asitus</b> Erdös, 1955	
<i>hofferi</i> Graham, 1966	B M	<i>phragmitis</i> (Ferrière, 1955)	S
<b>Dinocarsiella</b> Mercet, 1921		<b>Leptomastidea</b> Mercet, 1916	
<i>alpina</i> (Girault, 1917)	B M S	<i>bifasciata</i> (Mayr, 1876)	M S
<b>Dicarnosis</b> Mercet, 1921		<b>Callipteroma</b> Motschulsky, 1863	
<i>helena</i> Hoffer, 1953	B M S	<i>sexguttata</i> Motschulsky, 1863	S
<b>Dusmetia</b> Mercet, 1921		<b>Anusia</b> Foerster, 1856	
<i>cardinalis</i> Hoffer, 1969	M S	<i>fulvescens</i> Westwood, 1833	S
<i>ceballosi</i> Mercet, 1921	B M S	<i>nasicornis</i> Foerster, 1860	M
<i>pulex</i> (Ruschka, 1922)	B M S	<b>Paratetracnemoidea</b> Girault, 1915	
<b>Aquaencyrtus</b> Hoffer, 1953		<i>malenottii</i> (Mercet, 1918)	S
<i>boemicus</i> Hoffer, 1953	B	<b>Metaphaenodiscus</b> Mercet, 1921	
<b>Xanthoectroma</b> Mercet, 1925		<i>nemoralis</i> Mercet, 1921	M S
<i>aquilinum</i> Mercet, 1925	B M S	<b>Monodiscodes</b> Hoffer, 1954	
<b>Anagyrus</b> Howard, 1896		<i>intermedius</i> (Mayr, 1875)	M S
<i>bohemani</i> (Westwood, 1837)	S	<b>Tetraenemus</b> Westwood, 1837	
<i>bouceki</i> Hoffer, 1953	B M	<i>bifasciatellus</i> Mercet, 1919	S
<i>diversicornis</i> Mercet, 1921	M	<i>diversicornis</i> Westwood, 1837	B M S
<i>minium</i> (Mercet, 1921)	M		
<i>novickyi</i> Hoffer, 1953	S		
<i>schoenherri</i> Westwood, 1837	B M S		

<i>heydeni</i> (Mayr, 1876)	M	<i>matritensis</i> (Mercet, 1921)	B M S
<i>simillimus</i> (Hoffer, 1953)	B	<i>micropterus</i> (Mercet, 1921)	M S
<b>Mohelniella</b> Hoffer, 1964		<i>nikolskajae</i> Erdős, 1955	B M S
<i>silhavyi</i> Hoffer, 1964	M S	<i>problematicus</i> Hoffer, 1977	B M S
<b>Paramasia</b> Hoffer, 1953		<i>rogenhoferi</i> (Mayr, 1875)	B M S
<i>slovaca</i> Hoffer, 1953	M S	<i>sylvius</i> (Dalman, 1820)	B M S
<b>Tricladia</b> Mercet, 1918		<i>subcupratus</i> (Dalman, 1820)	B M
<i>humilis</i> Mercet, 1918	M	<i>tessellatus</i> (Dalman, 1820)	B M S
<b>Paracopidosoma</b> Hoffer, 1957		<b>Aschitus</b> Mercet, 1921	
<i>parallelum</i> Hoffer, 1957	B	<i>annulatus</i> Erdős, 1957	S
<b>Tetracnemoidea</b> Howard, 1898		<i>carpathicus</i> (Hoffer, 1958)	M S
<i>piceae</i> (Erdős, 1946)	B M S	<i>jalus</i> (Walker, 1837)	B M S
<b>Mohelenencyrtus</b> Hoffer, 1969		<i>javorinensis</i> (Hoffer, 1958)	M
<i>acuminatus</i> Hoffer, 1969	M S	<i>madyes</i> (Walker, 1837)	B M S
<b>Anomalicornia</b> Mercet, 1921		<i>mariae</i> (Hoffer, 1953)	B
<i>tenuicornis</i> Mercet, 1921	S	<i>semihyalinus</i> (Hoffer, 1953)	M S
ENCYRTINAE			
<b>Aphycooides</b> Mercet, 1921		<b>Trichomasthus</b> Thomson, 1876	
<i>clavellatus</i> (Dalman, 1820)	B M S	<i>coeruleus</i> Mercet, 1923	B M S
<i>niger</i> (Hoffer, 1957)	B	<i>bavarici</i> Hoffer, 1965	B M
<b>Plesiomicroterys</b> Ishii, 1928		<i>calonotus</i> (Mercet, 1921)	B M
<i>speciosus</i> Hoffer, 1968	B S	<i>dissimilis</i> (Tshumakova, 1961)	B S
<b>Microterys</b> Thomson, 1876		<b>Amicroterys</b> Myartseva, 1983	
<i>aeneiventris</i> (Walker, 1937)	B M S	<i>tetricus</i> (Hoffer, 1958)	S
<i>barbarus</i> (Dalman, 1820)	M	<b>Hoplopsis</b> DeStefani, 1891	
<i>brachypterus</i> (Mercet, 1921)	B S	<i>minuta</i> (Fabricius, 1793)	S
<i>carpathicus</i> Hoffer, 1977	S	<b>Subprionomitus</b> Mercet, 1921	
<i>chalcostomus</i> (Dalman, 1820)	B M S	<i>festucae</i> (Mayr, 1876)	M
<i>contractus</i> (Hoffer, 1953)	B M S	<i>paludosus</i> Hoffer, 1969	B
<i>dichrous</i> (Mercet, 1921)	M S	<b>Brachyencyrtus</b> Hoffer, 1959	
<i>duplicatus</i> (Nees, 1834)	B M S	<i>araneoides</i> (Hoffer, 1957)	B
<i>ferrugineus</i> (Nees, 1834)	B M S	<b>Platencyrtus</b> Ferrière, 1955	
<i>feudatarius</i> (Mercet, 1921)	B M S	<i>parkeri</i> Ferrière, 1955	B S
<i>fuscipennis</i> (Dalman, 1820)	B M S	<b>Syrphophagus</b> Ashmead, 1900	
<i>herbaceus</i> Sugonjaev, 1962	B M S	<i>aeruginosus</i> (Dalman, 1820)	B M S
<i>hortulanus</i> Erdős, 1956	S	<i>annelicornis</i> (Hoffer, 1965)	B
<i>lunatus</i> (Dalman, 1820)	B M S	<i>annulipes</i> (Thomson, 1876)	B S

<i>aphidivorus</i> (Mayr, 1876)	B	M	S
<i>ariantes</i> (Walker, 1837)	B	M	S
<i>arundinicola</i> (Hoffer, 1965)	B	M	S
<i>danuvicus</i> (Erdös, 1957)	B		
<i>fabulosus</i> (Hoffer, 1965)	B		
<i>fuscipes</i> (Dalman, 1820)	B		
<i>herbidus</i> (Dalman, 1820)	B	M	S
<i>mamitus</i> (Walker, 1837)	B		
<i>quadrii</i> (Alam, 1961)		S	
<i>quercicola</i> Hoffer, 1970	B	M	
<i>rugulosus</i> Hoffer, 1963	B		
<i>subviridis</i> Hoffer, 1970		S	
<i>staryi</i> (Hoffer, 1970)	B		
<i>taeniatus</i> (Foerster, 1861)	B	M	
<b>Ooencyrtus</b> Ashmead, 1900			
<i>albopilosus</i> Hoffer, 1963	B	M	S
<i>azureus</i> (Mercet, 1921)	B		
<i>fulvipes</i> Hoffer, 1963	B		
<i>salicinus</i> (Erdös, 1957)		M	
<i>telenomicida</i> (Vasiljev, 1904)	B	M	S
<b>Cerchysius</b> Westwood, 1833			
<i>gigas</i> Erdös, 1955	B	S	
<i>subplanus</i> (Dalman, 1820)	B	M	S
<b>Helegonatopus</b> Perkins, 1906			
<i>citripes</i> (Erdös, 1957)		M	S
<i>dimorphus</i> (Hoffer, 1954)	B	M	
<i>formosus</i> (Mercet, 1919)		S	
<i>nikolskajae</i> Hoffer, 1965	B	S	
<b>Blastothrix</b> Mayr, 1876			
<i>klapaleki</i> Hoffer, 1963	B		
<i>longipennis</i> Howard, 1881	B	M	S
<i>confusa</i> Erdös, 1959			
<i>matesovae</i> Sugonjaev, 1964		?	
<i>nikolskajae</i> Sugonjaev, 1964		?	
<i>pragensis</i> Hoffer, 1963	B		
<i>trjapitzini</i> Sugonjaev, 1976		?	
<b>Metablastothrix</b> Sugonjaev, 1964			
<i>truncatipennis</i> (Ferrière, 1955)	B	S	
<b>Aphytus</b> Mayr, 1875			
<i>apicalis</i> (Dalman, 1820)	B		
<i>atratulus</i> Hoffer, 1954		M	
<i>fulvohirtus</i> Hoffer, 1954		M	
<i>moravicus</i> (Hoffer, 1953)		M	
<i>nigrociliatus</i> Hoffer, 1954		M	S
<i>sumavicus</i> Hoffer, 1954	B		
<b>Pseudococcobius</b> Timberlake, 1916			
<i>obenbergeri</i> (Novicky, 1928)		B	M
<b>Pseudaphytes</b> Clausen, 1915			
<i>austriacus</i> Mercet, 1925	B	M	
<b>Aphytulus</b> Hoffer, 1954			
<i>perparvus</i> Hoffer, 1954	B	M	S
<i>zavadili</i> Hoffer, 1954		M	S
<b>Pseudectroma</b> Girault, 1915			
<i>nocturna</i> (Hoffer, 1955)		M	
<b>Stemmatosteres</b> Timberlake, 1918			
<i>bohemicus</i> Hoffer, 1954	B		S
<b>Xenaphytes</b> Trjapitzin, 1982			
<i>flavovarius</i> (Mercet, 1921)	B	M	S
<b>Metaphytes</b> Mercet, 1921			
<i>asterolecanii</i> (Mercet, 1923)			?
<i>callunae</i> (Alam, 1957)			?
<i>dispar</i> (Mercet, 1925)		S	
<i>erythraeus</i> Hoffer, 1954		S	
<i>flavus</i> (Howard, 1881)		M	
<i>fulvifrons</i> (Walker, 1838)		?	
<i>hanstediensis</i> Bakkendorf, 1965		B	
<i>hubai</i> Hoffer, 1969		S	
<i>nitens</i> (Kurdjumov, 1912)		?	
<i>notatus</i> Hoffer, 1954	B		
<i>piceus</i> Hoffer, 1954	B	M	S
<i>punctipes</i> (Dalman, 1820)		M	
<i>roubali</i> Hoffer, 1954	B		
<i>stagnarum</i> Hoffer, 1954	B		
<i>unicolor</i> Hoffer, 1954		M	
<i>zebratus</i> (Mercet, 1917)		M	
<b>Trechnites</b> Thomson, 1876			
<i>alni</i> Erdös, 1957		B	

<i>flavipes</i> (Mercet, 1921)	B M	<i>talitzkii</i> Trjapitzin, 1978	S
<i>fuscitarsis</i> Thomson, 1876	B M S	<i>tertia</i> (Hoffer, 1954)	B M
<i>psyllae</i> Ruschka, 1923	B S	<b>Baeocharis</b> Mayr, 1876	
<b>Sectiliclava</b> Hoffer, 1957		<i>pascuorum</i> Mayr, 1876	M
<i>cleone</i> (Walker, 1844)	B	<b>Mayridia</b> Mercet, 1921	
<b>Prionomitus</b> Mayr, 1876		<i>borkovicensis</i> (Hoffer, 1957)	B
<i>mitratus</i> (Dalman, 1820)	B M	<i>egidiopolitana</i> Erdös, 1957	M
<i>perbellus</i> (Erdös, 1957)	S	<i>subfuscipennis</i> Erdös et Novicky, 1957	M S
<i>tiliaris</i> (Dalman, 1820)	B M	<i>tatrica</i> (Hoffer, 1957)	S
<i>vicinus</i> Hoffer, 1963	B M		
<b>Psyllaephagus</b> Ashmead, 1900		<b>Ectroma</b> Westwood, 1833	
<i>abbreviatus</i> (Hoffer, 1963)	B M S	<i>aeneiventre</i> (Walker, 1837)	B M S
<i>belanensis</i> (Hoffer, 1963)	S	<i>albiblavatum</i> (Hoffer, 1957)	B M S
<i>smaragdinus</i> (Hoffer, 1963)	S	<i>arenarium</i> (Erdös, 1955)	B M S
<i>taborita</i> (Hoffer, 1963)	B	<i>fulvescens</i> Westwood, 1833	M
<i>vendicus</i> Erdös, 1961	S	<i>reinhardi</i> (Mayr, 1875)	B M S
<b>Cercobelus</b> Walker, 1840		<b>Ixodiphagus</b> Howard, 1907	
<i>jugaetus</i> (Walker, 1840)	B M	<i>hookeri</i> (Howard, 1908)	B S
<b>Discodes</b> Foerster, 1856		<b>Thomsonisca</b> Ghesquière, 1946	
<i>aeneus</i> (Dalman, 1820)	B M S	<i>amathus</i> (Walker, 1838)	B
<i>anthores</i> (Walker, 1848)	M	<b>Habrolepis</b> Foerster, 1856	
<i>cercopiformis</i> (Walker, 1875)	M S	<i>dalmanni</i> (Westwood, 1837)	B M S
<i>monachus</i> (Mercet, 1921)	B M	<i>pascuorum</i> Mercet, 1921	S
<i>rubcovi</i> Sugonjaev, 1971	S	<b>Epitetracnemus</b> Girault, 1915	
<b>Choreia</b> Westwood, 1833		<i>zetterstedtii</i> (Westwood, 1837)	B S
<i>inepta</i> (Dalman, 1820)	B M S	<b>Adelencyrtus</b> Asmead, 1900	
<i>maculata</i> (Hoffer, 1954)	S	<i>aulacaspidis</i> (Brèthes, 1914)	B
<i>stepicola</i> (Hoffer, 1953)	S	<i>arenarius</i> Erdös, 1957	S
<b>Semen</b> Hoffer, 1954		<b>Comperiella</b> Howard, 1906	
<i>apterum</i> Hoffer, 1954	M S	<i>bifasciata</i> Howard, 1906	B
<b>Mira</b> Schellenberg, 1803		<b>Cerapterocerus</b> Westwood, 1833	
<i>mucora</i> Schellenberg, 1803	M	<i>mirabilis</i> Westwood, 1833	B M S
<b>Echthroplexiella</b> Mercet, 1921		<i>pilicornis</i> Thomson, 1875	B M S
<i>aeneiventris</i> Mercet, 1921	S	<b>Eusemion</b> Dahlbom, 1857	
<i>flava</i> Mercet, 1921	B	<i>cornigerum</i> Walker, 1838	B M S
<i>metatibialis</i> Erdös, 1957	S		
<i>moravica</i> Hoffer, 1953	B M S		
<i>niveicornis</i> Hoffer, 1963	S		
<i>obscura</i> (Hoffer, 1954)	M		
<i>orientalis</i> Hoffer, 1953	M S		
<i>similis</i> Hoffer, 1953	B M S		

<b>Cheiloneurus</b> Westwood, 1833			
<i>boldyrevi</i> Trjapitzin et Agekian, 1978		?	
<i>claviger</i> Thomson, 1876	B	M	S
<i>formosus</i> Boheman, 1852	B	M	S
<i>fulvescens</i> Hoffer, 1957	B	M	
<i>kollari</i> Mayr, 1875		M	S
<i>longicornis</i> Ruschka, 1923	B	M	S
<i>paralia</i> (Walker, 1837)	B	M	
<i>quercus</i> Mayr, 1875		M	
<i>submuticus</i> Thomson, 1876	B	M	S
<i>victor</i> Hoffer, 1957	B	M	S
<b>Zaomma</b> Ashmead, 1900			
<i>eriococci</i> (Ferrière, 1955)	B	M	
<i>lambinus</i> (Walker, 1838)	B	M	S
<i>trichomasthooides</i> (Hoffer, 1965)	B		
<b>Achalcerinys</b> Girault, 1915			
<i>lindus</i> Mercet, 1921		M	S
<b>Mahencyrtus</b> Masi, 1917			
<i>comara</i> (Walker, 1837)	B	M	S
<b>Tyndarichus</b> Howard, 1910			
<i>melanacis</i> (Dalman, 1820)	B	M	S
<i>scaurus</i> (Walker, 1838)	B	M	S
<i>pulcherrimus</i> Hoffer, 1969	B		
<b>Leiocyrtus</b> Erdös et Novicky, 1955			
<i>clavatus</i> Erdös et Novicky, 1955	B	M	
<b>Boucekiella</b> Hoffer, 1954			
<i>depressa</i> Hoffer, 1954		M	S
<b>Prochiloneurus</b> Silvestri, 1915			
<i>bolivari</i> Mercet, 1921	B	M	S
<b>Mayrencyrtus</b> Hincks, 1944			
<i>imandes</i> (Walker, 1837)	B		
<i>maculatus</i> (Hoffer, 1957)	B		
<b>Trjapitzinellus</b> Viggiani, 1967			
<i>nigricornis</i> Hoffer, 1975	B		
<i>obscurus</i> (Mercet, 1921)		M	
<i>semidaliphagus</i> Viggiani, 1967	B	M	
<b>Cerchysiella</b> Girault, 1914			
<i>centennalis</i> (Erdös, 1956)			S
<i>planiscutellum</i> (Mercet, 1921)			M
<b>Coenocercus</b> Thomson, 1876			
<i>puncticollis</i> Thomson, 1876		B	M
<b>Copidosoma</b> Ratzeburg, 1844			
<i>agrotis</i> (Fonscolombe, 1832)		B	M
<i>albipes</i> (Westwood, 1837)	B		
<i>ancharus</i> (Walker, 1837)	B	M	S
<i>arenarium</i> Erdös, 1961			S
<i>aretas</i> (Walker, 1838)	B		S
<i>auricolle</i> (Thomson, 1875)	B	M	S
<i>bohemicum</i> Hoffer, 1969	B		
<i>bolivari</i> Mercet, 1921	B	M	S
<i>boreale</i> Hoffer, 1970	B		
<i>boucheanum</i> Ratzeburg, 1844		B	
<i>breviclava</i> Hoffer, 1970		M	S
<i>budense</i> Erdös, 1955	B		
<i>cervius</i> (Walker, 1848)	B	M	S
<i>chalconotum</i> (Dalman, 1820)	B		S
<i>clavigerum</i> (Mercet, 1921)	B		
<i>coleophorae</i> Mayr, 1875	B	M	S
<i>cultriforme</i> Mayr, 1875	B		S
<i>cyaneum</i> Hoffer, 1970			M
<i>dius</i> (Walker, 1837)	B	M	S
<i>extraneum</i> (Hoffer, 1970)	B	M	
<i>filocrine</i> (Dalman, 1820)	B		
<i>flagellare</i> (Dalman, 1820)	B	M	S
<i>flavomaculatum</i> (Ratzeburg, 1848)	B		S
<i>fulvipes</i> (Erdös, 1960)	B	M	S
<i>fuscisquama</i> (Thomson, 1875)	B		S
<i>genale</i> (Thomson, 1875)	B	M	S
<i>geniculatum</i> (Dalman, 1820)	B	M	S
<i>giganteum</i> Hoffer, 1957	B		
<i>globiceps</i> Erdös, 1955			S
<i>hartmanni</i> (Mayr, 1875)	B	M	
<i>incertum</i> Mercet, 1921			S
<i>intermedium</i> (Mercet, 1921)	B	M	S
<i>iracundum</i> Erdös, 1957	B	M	S

<i>kriechbaumeri</i> (Mayr, 1875)	B	<i>hybridus</i> Hoffer, 1963	S
<i>mohelense</i> Hoffer, 1970	M	<i>nigricornis</i> Mercet 1921	?
<i>moldavicum</i> (Hoffer, 1957)	B	<i>platynaspidis</i> Hoffer, 1963	B M S
<i>monochroum</i> Hoffer, 1970	M	<i>singularis</i> Hoffer, 1963	B
<i>nanellae</i> Silvestri, 1923	M S	<b>Homalotyloidea</b> Mercet, 1921	
<i>nanum</i> (Hoffer, 1970)	S	<i>crassa</i> (Hoffer, 1957)	M S
<i>peticus</i> (Walker, 1846)	B M S	<i>erginus</i> (Walker, 1837)	B
<i>phalaenarum</i> (Thomson, 1875)	B M S	<i>nigrociliata</i> (Hoffer, 1954)	M S
<i>pragense</i> Novicky, 1925	B M S	<i>nowickyi</i> Hoffer, 1957	B M S
<i>prasimum</i> Hoffer, 1970	S	<i>snoflaki</i> (Hoffer, 1954)	B M
<i>ratzeburgi</i> Mercet, 1921	M S	<b>Isodromus</b> Howard, 1886	
<i>salicinum</i> (Erdös, 1956)	M	<i>flaviscutum</i> Hoffer et Trjapitzin, 1978	?
<i>sashegyense</i> Erdös, 1957	B S	<i>limosus</i> Hoffer, 1969	B
<i>scutellare</i> (Hoffer, 1970)	B	<i>puncticeps</i> (Howard, 1855)	B M S
<i>serricorne</i> (Dalman, 1820)	B	<i>vinulus</i> (Dalman, 1820)	B M S
<i>sosares</i> (Walker, 1837)	B	<b>Eupcecilopoda</b> Novicky et Hoffer, 1953	
<i>thebe</i> (Walker, 1838)	B	<i>perpunctata</i> (Masi, 1942)	M S
<i>tibiale</i> Hoffer, 1970	B	<b>Encyrtus</b> Latreille, 1809	
<i>triangulare</i> (Thomson, 1875)	B M S	<i>albitarsis</i> Zetterstedt, 1838	B M S
<i>truncatellum</i> (Dalman, 1820)	B M S	<i>infidus</i> (Rossi, 1790)	B M S
<i>viridiaeneum</i> Hoffer, 1970	B	<i>lecaniorum</i> (Mayr, 1876)	B
<b>Paralitomastix</b> Mercet, 1921		<i>swederi</i> Dalman, 1820	B M S
<i>subalbicornis</i> (Nees, 1834)	B M S	<b>Eugahania</b> Mercet, 1921	
<i>varicornis</i> Hoffer, 1960	B	<i>fumipennis</i> (Ratzeburg, 1852)	S
<b>Copidosomopsis</b> Girault, 1915		<b>Prionomastix</b> Mayr, 1876	
<i>bohemica</i> (Hoffer, 1960)	B S	<i>morio</i> (Dalman, 1820)	S
<b>Parablastothrix</b> Mercet, 1917		<b>Pseudorhopus</b> Timberlake, 1926	
<i>metatibialis</i> Erdös, 1955	B	<i>testaceus</i> (Ratzeburg, 1848)	B
<i>montana</i> Erdös, 1955	B S	<b>Psilophrys</b> Mayr, 1876	
<i>vespertina</i> Mercet, 1917	?	<i>tenuicornis</i> Graham, 1969	B
<b>Calometopia</b> Mercet, 1921		<b>Quadrenencyrtus</b> Hoffer, 1953	
<i>kodensis</i> Hoffer, 1957	B	<i>paradoxus</i> Hoffer, 1953	B M S
<b>Coelopencyrtus</b> Timberlake, 1919		<b>Arrhenophagus</b> Aurivillius, 1888	
<i>arenarius</i> (Erdös, 1957)	B S	<i>chionaspidis</i> Aurivillius, 1888	B S
<b>Homalotylus</b> Mayr, 1876			
<i>ephiphium</i> Ruschka, 1923	?		
<i>eytelweinii</i> (Ratzeburg, 1844)	B		
<i>flaminius</i> (Dalman, 1820)	B M S		

<b>Anthemus</b> Howard, 1896					
<i>funicularis</i> (Bakkendorf, 1939)	B	S			
<b>Encyrtinae</b> — incertae sedis					
<b>Admireencyrtus</b> Hoffer, 1960					
<i>insolitus</i> Hoffer, 1960	S				
<b>Agromyzaphagus</b> Gahan, 1912					
<i>detrimentosus</i> Gahan, 1912	B				
<b>Allocerchysius</b> Hoffer, 1963					
<i>biroi</i> (Erdös, 1955)	B M				
<b>Parablatticida</b> Girault, 1915					
<i>montana</i> (Mercet, 1921)	B				
<b>Ceballosia</b> Mercet, 1921					
<i>dusmeti</i> Mercet, 1921	B				
<b>Eucoccidophagus</b> Hoffer, 1963					
<i>breviventris</i> (Kurdjumov, 1912)	B				
<i>semiluniger</i> (Hoffer, 1959)	B				
<b>Ginsiana</b> Erdös					
<i>et Novicky, 1955</i>					
<i>carpetana</i> (Mercet, 1921)	B M				
<i>obscura</i> Erdös					
<i>et Novicky, 1955</i>	?				
<i>praepannonica</i> Erdös, 1957	B M S				
<b>Globulencyrtus</b> Hoffer, 1975					
<i>politum</i> (Hoffer, 1957)	S				
<b>Lamennaisia</b> Girault, 1922					
<i>ambiguus</i> (Nees, 1834)	M				
<b>Negenaspidius</b> Trjapitzin, 1982					
<i>nobilis</i> (Nees, 1834)	M				
<b>Parasauleia</b> Hoffer, 1958					
<i>trjapitzini</i> Hoffer, 1958	B M				
<b>SIGNIPHORIDAE</b>					
<b>Clytina</b> Erdös, 1957					
<i>giraudi</i> Erdös, 1957	B				
<b>Chartocerus</b> Motschulsky, 1859					
<i>subaeneus</i> (Foerster, 1878)	B M S				
<b>Thysanus</b> Haliday					
<i>in Walker, 1840</i>					
<i>ater</i> Haliday in Walker, 1840	B M				
<b>APHELINIDAE</b>					
<b>Coccophagus</b> Westwood, 1833					
<i>lycimnia</i> (Walker, 1839)	B M S				
<i>obscurus</i> Westwood, 1833	B S				
<i>piceae</i> Erdös, 1956	B M S				
<i>pulchellus</i> Westwood, 1833	B				
<i>semicircularis</i> (Foerster, 1841)	B M S				
<i>silvestrii</i> Compere, 1931	B				
<b>Coccobius</b> Ratzeburg, 1852					
<i>annulicornis</i> Ratzeburg, 1852	B M S				
<b>Archenomus</b> Howard, 1898					
<i>bicolor</i> Howard, 1898	B S				
<i>longiclava</i> Girault, 1915	B M S				
<i>opacus</i> (Erdös, 1953)	B M S				
<b>Coccophagooides</b> Girault, 1915					
<i>moeris</i> (Walker, 1839)	B				
<b>Encarsia</b> Foerster, 1878					
<i>citrina</i> (Craw, 1891)	S				
<i>formosa</i> Gahan, 1924					
introd.	B M S				
<i>gigas</i> (Tschumakova, 1957)	B S				
<i>leucaspidis</i> (Mercet, 1912)	B M				
<i>lutea</i> (Masi, 1910)	B S				
<i>longicornis</i> Mercet, 1928	B M				
<i>perniciosi</i> (Tower, 1913)					
introd.	B M S				
<i>tricolor</i> Foerster, 1878	B M				
<b>Hispaniella</b> Mercet, 1911					
<i>lauri</i> Mercet, 1911	S				
<b>Pteroptrix</b> Westwood, 1833					
<i>dimidiata</i> Westwood, 1833	S				
<i>macropedicellata</i> (Maláč, 1947)	B M S				
<b>Aphelinus</b> Dalman, 1820					
<i>abdominalis</i> (Dalman, 1820)	B M S				
<i>annulipes</i> (Walker, 1839)	B M S				

<i>argiope</i> (Walker, 1839)	B M S	<i>arcuatus</i> Ferrière, 1947	M S
<i>asychis</i> (Walker, 1839)	B M S	<i>flabellatus</i> (Fonscolombe, 1832)	B S
<i>chaonia</i> (Walker, 1841)	B M S	<i>platyedrae</i> Ferrière, 1935	B M S
<i>confusus</i> Kalina, 1976	B M S	<i>rufiventris</i> Ferrière, 1947	B M S
<i>daucicola</i> Kurdjumov, 1913	B M S	<i>vibicellae</i> Ferrière, 1947	B M S
<i>flavipes</i> (Foerster, 1841)	B M S	<i>viridiceps</i> Thomson, 1878	B M S
<i>flaviventris</i> Kurdjumov, 1913	B M	<b>Euryischia</b> Riley, 1889	
<i>fulvus</i> Jasnoch, 1963	B	<i>inopinata</i> Masi, 1907	S
<i>humilis</i> Mercet, 1928	B M S	TETRACAMPIDAE	
<i>mali</i> (Haldeman, 1851) introd.	B M S	PLATYNOCHEILINAE	
<i>subflavescens</i> (Westwood, 1837)	B M S	<b>Platynocheilus</b> Westwood, 1837	
<i>tetrataenion</i> (Erdös et Novicky, 1953)	B S	<i>cuprifrons</i> (Nees, 1834)	B M S
<i>thomsoni</i> Graham, 1976	B M S	<i>gravenhorsti</i> (Ratzeburg, 1852)	B
<i>varipes</i> (Foerster, 1841)	B M S	TETRACAMPINAE	
<b>Aphytis</b> Howard, 1900		<b>Dipriocampe</b> Bouček, 1958	
<i>aonidiae</i> (Mercet, 1911)	B S	<i>diprioni</i> (Ferrière, 1935)	B
<i>hispanicus</i> (Mercet, 1912)	B	<i>elongata</i> (Erdös, 1951)	B M S
<i>moldavicus</i> Jasnoch, 1966	B	<b>Tetracampe</b> Foerster, 1841	
<i>mytilaspidis</i> (Le Baron, 1870)	B M S	<i>impressa</i> Foerster, 1841	B M S
<i>proclia</i> (Walker, 1839)	B M S	<b>Foersterella</b> Dalla Torre, 1897	
<b>Centrodora</b> Foerster, 1878		<i>erdeosi</i> Bouček, 1958	B M S
<i>acridiphaga</i> (Otten, 1941)	B M	<i>flavipes</i> (Foerster, 1841)	B M S
<i>amoena</i> Foerster, 1878	B M S	<b>Epiclerus</b> Haliday, 1843	
<i>dorsati</i> Mercet, 1930	B S	<i>nomocerus</i> (Masi, 1934)	S
<i>hexatricha</i> Erdös et Novicky, 1953	S	<i>panyas</i> (Walker, 1839)	B
<i>livens</i> (Walker, 1851)	B M	<i>temenus</i> (Walker, 1839)	B M S
<i>locustarum</i> (Giraud, 1864)	S	EULOPHIDAE	
<i>ochrura</i> Erdös et Novicky, 1953	B M S	EULOPHINAE	
<b>Eretmocerus</b> Haldeman, 1850		<b>Euplectrus</b> Westwood, 1832	
<i>mundus</i> Mercet, 1931	B M	<i>bicolor</i> (Swederus, 1795)	B M S
<b>Marietta</b> Motschulsky, 1863		<i>flavipes</i> (Fonscolombe, 1832)	S
<i>picta</i> (André, 1877)	B M S	<i>liparidis</i> Ferrière, 1941	S
<b>Azotus</b> Howard, 1898		<b>Trichoplectrus</b> Erdös, 1951	
<i>atomon</i> (Walker, 1847)	B M S	<i>bouceki</i> (Erdös, 1960)	S
<i>celsus</i> (Walker, 1839)	B	<b>Stenomesius</b> Westwood, 1833	
ELASMIDAE		<i>rufescens</i> (Rossi, 1790)	B M S
<b>Elasmus</b> Westwood, 1833		<b>Elachertus</b> Spinola, 1811	
<i>albipennis</i> Thomson, 1878	B M S		

<i>arenarius</i> Erdös, 1966	B		<i>pulcherrimus</i> (Mercet, 1916)	M S
<i>argissa</i> (Walker, 1839)	B M S		<i>salatis</i> Walker, 1838	B
<i>artaeus</i> (Walker, 1839)	B S		<i>singa</i> Walker, 1838	S
<i>charondas</i> (Walker, 1839)	B M S		<i>viticola</i> (Rondani, 1876)	B M S
<i>geniculatus</i> (Hartig, 1838)	M S		<i>staryi</i> Bouček, 1959	B
<i>inunctus</i> (Nees, 1834)	B M S		<i>variegatus</i> (Masi, 1907)	S
<i>isadas</i> (Walker, 1839)	B		<i>vittatus</i> Walker, 1838	B M S
<i>longipetiolatus</i> Bouček, 1971	B		<b>Diaulinopsis</b> Crawford, 1912	
<i>nigritulus</i> (Zetterstedt, 1838)	B M S		<i>arenaria</i> (Erdös, 1951)	B M S
<i>olivaceus</i> (Thomson, 1878)	B M S		<b>Sympiesis</b> Foerster, 1856	
<i>pilosiscuta</i> Bouček, 1968	B S		<i>acalle</i> (Walker, 1848)	B M S
<i>pulcher</i> (Erdös, 1961)	B M S		<i>angustipennis</i> (Erdös, 1954)	B M S
<b>Xanthellum</b> Erdös et Novicky, 1951			<i>asecta</i> Delucchi, 1962	S
<i>transsylvanicum</i> Erdös, 1951	B S		<i>capeki</i> Bouček, 1959	B S
<b>Rhienopelte</b> Foerster, 1878			<i>dolichogaster</i> Ashmead, 1898	B M S
<i>crassicornis</i> (Nees, 1834)	B M S		<i>euspilapterygis</i> (Erdös, 1958)	B
<b>Dermatopelte</b> Erdös et Novicky, 1951			<i>flavopicta</i> Bouček, 1959	B M S
<i>budensis</i> Erdös et Novicky, 1951	B S		<i>gordius</i> (Walker, 1839)	B M S
<b>Bryopezus</b> Erdös, 1951			<i>grahami</i> Erdös, 1966	B
<i>brevipennis</i> Erdös, 1951	B S		<i>gregori</i> Bouček, 1959	B M S
<b>Miotropis</b> Thomson, 1878			<i>gyoerfii</i> Erdös, 1954	S
<i>unipunctata</i> (Nees, 1834)	B M S		<i>notata</i> (Zetterstedt, 1838)	B M S
<b>Aulogymnus</b> Foerster, 1851			<i>sericeicornis</i> (Nees, 1834)	B M S
<i>arsames</i> (Walker, 1838)	B M		<i>viridula</i> (Thomson, 1878)	B M S
<i>gallarum</i> (Linnaeus, 1761)	B M		<i>xanthostoma</i> (Nees, 1834)	B M S
<i>obscuripes</i> (Mayr, 1877)	M S		<b>Dimmockia</b> Ashmead, 1904	
<i>skianeuros</i> (Ratzeburg, 1844)	B M		<i>brevicornis</i> (Erdös, 1954)	B S
<i>trilineatus</i> (Mayr, 1877)	M S		<b>Hemiptarsenus</b> Westwood, 1833	
<b>Cirrospilus</b> Westwood, 1832			<i>dropion</i> (Walker, 1839)	B M S
<i>argei</i> (Crawford, 1911)	B S		<i>fulvicollis</i> Westwood, 1833)	B S
<i>curvineurus</i> Askew, 1965	S		<i>unguicellus</i> (Zetterstedt, 1838)	B M S
<i>diallus</i> Walker, 1838	B S		<i>wailesellae</i> Novicky, 1929	B S
<i>elegantissimus</i> Westwood, 1832	B M		<i>waterhousei</i> Westwood, 1833	B M S
<i>elongatus</i> Bouček, 1959	B M		<i>zilahisebessi</i> Erdös, 1951	S
<i>lynCUS</i> Walker, 1838	B M S		<b>Cleolophus</b> Mercet, 1824	
<i>pictus</i> (Nees, 1834)	B M S		<i>autonomus</i> Mercet, 1924	M S
<i>pulcher</i> Masi, 1911	B M S		<b>Di cladocerus</b> Westwood, 1832	
			<i>breviramulus</i> Bouček, 1959	B M S

<i>euryalus</i> (Haliday, 1843)	B	<i>tidius</i> (Walker, 1839)	B M S
<i>westwoodii</i> Westwood, 1832	B M S	<b>Microlycus</b> Thomson, 1878	
<b>Ratzeburgiola</b> Erdös, 1958		<i>biroi</i> Erdös, 1951	B M S
<i>cristata</i> (Ratzeburg, 1848)	B	<i>erdoesi</i> Bouček, 1959	B M S
<i>incompleta</i> Bouček, 1971	S	<i>gyoerpii</i> (Erdös, 1954)	M
<b>Pnigalio</b> Schrank, 1802		<i>heterocerus</i> Thomson, 1878	B
<i>agraules</i> (Walker, 1839)	B	<i>virens</i> Erdös, 1951	B M S
<i>attis</i> (Walker, 1839)	B	<b>Colpocephalus</b> Lucchese, 1941	
<i>epilobii</i> Bouček, 1966	B	<i>florus</i> (Walker, 1839)	B S
<i>hirtulus</i> (Erdös, 1954)	B	<b>Diglyphus</b> Walker, 1848	
<i>longulus</i> (Zetterstedt, 1838)	B	<i>albiscapus</i> Erdös, 1951	B S
<i>monilicornis</i> (Zetterstedt, 1838)	B	<i>begini</i> (Ashmead, 1902)	B S
<i>nemati</i> (Westwood, 1838)	B M S	<i>chabrias</i> (Walker, 1838)	B S
<i>pectinicornis</i> (Linnaeus, 1758)	B M S	<i>crassinervis</i> Erdös, 1958	B S
<i>phragmitis</i> (Erdös, 1954)	B	<i>isaea</i> (Walker, 1838)	B M S
<i>soemius</i> (Walker, 1839)	B M S	<i>minoeus</i> (Walker, 1838)	B S
<i>tricuspidis</i> (Erdös, 1954)	B	<i>poppaea</i> Walker, 1848	B M
<i>tridentatus</i> (Thomson, 1878)	B	<i>pusztensis</i> (Erdös et Novicky, 1951)	B S
<i>xerophilus</i> (Erdös, 1954)	S	<i>sabulosus</i> Erdös, 1951	B S
<b>Dahlbominus</b> Hincks, 1945		TETRASTICHINAE	
<i>fuscipennis</i> (Zetterstedt, 1838)	B M S	<b>Eutetrastichus</b> Kostjukov, 1977	
<b>Eulophus</b> Müller, 1764		<i>adalia</i> (Walker, 1839)	B
<i>abdominalis</i> Nees, 1834	B M S	<i>amethystinus</i> (Ratzeburg, 1848)	B
<i>aeneicoxa</i> (Thomson, 1878)	B M S	<i>cirsii</i> (Kurdjumov, 1913)	B S
<i>cyanescens</i> Bouček, 1959	B	<i>crassicornis</i> (Erdös, 1954)	B
<i>larvarum</i> (Linnaeus, 1758)	B M S	<i>daira</i> (Walker, 1839)	B
<i>pectinicornis</i> Nees, 1834	B	<i>diaphanthus</i> (Walker, 1839)	B S
<i>smerinthicida</i> Bouček, 1959	B	<i>evonymellae</i> (Bouché, 1834)	B
<i>slovacus</i> Bouček, 1959	S	<i>galactopus</i> (Ratzeburg, 1844)	B
<i>thespius</i> Walker, 1839	B	<i>oophagus</i> (Otten, 1942)	B
<b>Necremnus</b> Thomson, 1878		<i>principiae</i> (Domenichini, 1965)	B S
<i>artynes</i> (Walker, 1839)	B M	<i>turionum</i> (Hartig, 1838)	M
<i>capitatus</i> Bouček, 1959	B	<b>Tetrastichus</b> Haliday, 1844	
<i>cosconius</i> (Walker, 1839)	S	<i>epilachnae</i> (Giard, 1896)	S
<i>folia</i> (Walker, 1839)	B M S	<i>halidayi</i> (Graham, 1961)	B
<i>hungaricus</i> (Erdös, 1951)	S	<i>hylocomarum</i> (Bouché, 1834)	B S
<i>leucarthos</i> (Nees, 1834)	B M S	<i>julii</i> (Walker, 1839)	B
<i>metalarus</i> (Walker, 1839)	B		

<b>legionarius</b> Giraud, 1863	B	S	<b>Minotetrastichus</b> Kostjukov, 1977
<i>lyridice</i> (Walker, 1839)		M	
<i>miser</i> (Nees, 1834)	B		
<i>murcia</i> (Walker, 1839)	B	M	
<i>sempronius</i> Erdős, 1954	B	S	
<i>telon</i> (Graham, 1961)	B		
<i>temporalis</i> Graham, 1961	B		
<i>trichops</i> Thomson, 1878	B		
<i>ulmi</i> Erdős, 1954	B		
<b>Holcotetrastichus</b> Graham, 1987			
<i>rhosaces</i> (Walker, 1839)	B	M	
<b>Oomyzus</b> Rondani, 1870			
<i>cocinellae</i> (Kurdjumov, 1912)	B	S	
<i>fumatus</i> (Erdős, 1954)	B		
<i>gallerucae</i> (Fonscolombe, 1832)	B		
<i>galerucivorus</i> (Hedquist, 1959)	B	M	
<i>incertus</i> (Ratzeburg, 1844)	B		
<b>Sphaenolepis</b> Nees, 1834			
<i>pygmaea</i> Nees, 1834	B	M	
<b>Tamarixia</b> Mercet, 1924			
<i>actis</i> (Walker, 1839)	B		
<i>monesus</i> (Walker, 1839)	B	M	
<i>pubescens</i> (Nees, 1834)	B	S	
<i>upis</i> (Walker, 1839)	B	M	
<b>Cecidotetrastichus</b> Kostjukov, 1977			
<i>brevinervis</i> (Zetterstedt, 1838)	B		
<i>citrinellus</i> (Graham, 1961)	B		
<i>lasiocera</i> (Graham, 1961)	B		
<i>podicularis</i> (Graham, 1961)	B		
<i>sajoi</i> (Szelényi, 1941)	B	S	
<i>vacuna</i> (Walker, 1839)	B	S	
<b>Mischotetrastichus</b> Graham, 1987			
<i>petiolatus</i> (Erdős, 1961)	B		
<b>Apotetrastichus</b> Graham, 1987			
<i>postmarginalis</i> (Bouček, 1971)	B		
			<b>Minotetrastichus</b> Kostjukov, 1977
			<i>ecus</i> (Walker, 1838) B M S
			<i>loxotoma</i> (Graham, 1961) S
			<i>platanellus</i> (Mercet, 1922) M S
			<b>Xenaprostocetus</b> Graham, 1987
			<i>pungens</i> Graham, 1987 M
			<b>Aprostocetus</b> Westwood, 1833
			<i>abydenus</i> (Walker, 1848) B
			<i>acron</i> Graham, 1987 B
			<i>aethiops</i> (Zetterstedt, 1838) B S
			<i>agevilleae</i> Graham, 1987 S
			<i>agrus</i> (Walker, 1839) B S
			<i>amenon</i> (Walker, 1839) B
			<i>apama</i> (Walker, 1839) B
			<i>apiculatus</i> Graham, 1987 S
			<i>arenarius</i> (Erdős, 1954) S
			<i>aristaeus</i> (Walker, 1839) B
			<i>arrabonicus</i> (Erdős, 1954) B S
			<i>aurantiacus</i> (Ratzeburg, 1848) B
			<i>boreus</i> (Delucchi, 1954) B
			<i>brachycerus</i> (Thomson, 1878) B
			<i>brevipennis</i> Graham, 1987 M S
			<i>bruzzonis</i> (Masi, 1930) B
			<i>calamarius</i> (Graham, 1961) B S
			<i>catius</i> (Walker, 1839) B
			<i>caudatus</i> (Westwood, 1833) B M S
			<i>cecidomyiarum</i> (Bouché, 1834) S
			<i>celtidis</i> (Erdős, 1954) B
			<i>cerricola</i> (Erdős, 1954) B
			<i>ciliatus</i> (Nees, 1834) B
			<i>citrinus</i> (Foerster, 1841) B
			<i>citripes</i> (Thomson, 1878) B
			<i>clavicornis</i> (Zetterstedt, 1838) B
			<i>collega</i> (Ratzeburg, 1844) B
			<i>crino</i> (Walker, 1838) B M S
			<i>dauci</i> Graham, 1987 B
			<i>deplanatus</i> (Walker, 1874) B
			<i>distichus</i> Graham, 1961 B
			<i>doksyensis</i> Graham, 1987 B
			<i>domenichini</i> (Erdős, 1969) S

<i>elongatus</i> (Foerster, 1841)	B	M	S	<i>pseudopodiellus</i> (Bakkendorf, 1953)	B
<i>emesa</i> (Walker, 1839)	B		S	<i>pygmaeus</i> (Zetterstedt, 1838)	B
<i>epilobii</i> Graham, 1987	B		S	<i>rhipheus</i> (Walker, 1839)	B
<i>escherichi</i> (Szelényi, 1941)	B		S	<i>rubi</i> Graham, 1987	B
<i>epilachnae</i> (Giard, 1896)			S	<i>rufus</i> Bakkendorf, 1953	B
<i>epicharmus</i> (Walker, 1839)	B	M	S	<i>salictorum</i> Graham, 1987	S
<i>eupatorii</i> Kurdjumov, 1913	B			<i>serratularum</i> Graham, 1987	B
<i>eurytomae</i> (Nees, 1834)	B			<i>setulosus</i> Graham, 1987	S
<i>eurytus</i> (Walker, 1838)			S	<i>stenus</i> Graham, 1987	B
<i>forsteri</i> (Walker, 1847)	B	M	S	<i>strobilanae</i> (Ratzeburg, 1844)	B
<i>fonscolombei</i> Graham, 1987	B			<i>subanellatus</i> Graham, 1961	B
<i>fulvipes</i> (Foerster, 1878)	B		S	<i>subcylidricus</i> Graham, 1987	B
<i>gaus</i> (Walker, 1839)	B			<i>subplanus</i> Graham, 1987	B
<i>gratus</i> (Giraud, 1863)	B			<i>suevius</i> (Walker, 1839)	B
<i>grylli</i> (Erdős, 1954)			S	<i>tenuiradialis</i> Graham, 1987	B
<i>leucone</i> (Walker, 1839)	B	M	S	<i>terebrans</i> Erdős, 1954	B
<i>longicauda</i> (Thomson, 1878)	B		S	<i>tiliaceae</i> Graham, 1987	B
<i>longiscapus</i> (Thomson, 1878)	B			<i>tompanus</i> (Erdős, 1954)	B
<i>luteus</i> (Ratzeburg, 1852)	B	M	S	<i>tymber</i> (Walker, 1839)	S
<i>lycidas</i> (Walker, 1839)	B	M	S	<i>venustus</i> (Gahan, 1914)	B
<i>lysippe</i> (Walker, 1839)	B			<i>viridinitens</i> Graham, 1987	B
<i>mandanis</i> (Walker, 1838)	B	M	S	<i>westwoodii</i> (Fonscolombe, 1840)	B
<i>menius</i> (Walker, 1839)	B		S	<i>xanthomelas</i> Graham, 1987	B
<i>mycerinus</i> (Walker, 1839)	B		S	<i>xanthopus</i> (Nees, 1834)	B
<i>neglectus</i> (Domenichini, 1956)	B			<i>zoilus</i> (Walker, 1839)	B
<i>novatus</i> (Walker, 1839)			S	<i>zosimus</i> (Walker, 1839)	B
<i>oreophilus</i> (Foerster, 1861)	B			<b>Neotrichoporooides</b> Girault, 1913	
<i>oreastes</i> Graham, 1987	B			<i>biogradensis</i> Graham, 1987	S
<i>orithyia</i> (Walker, 1839)	B	M	S	<i>cavigena</i> Graham, 1987	S
<i>ovivorax</i> Silvestri, 1920	B			<i>dubiosus</i> Graham, 1987	S
<i>pachyneuros</i> (Ratzeburg, 1844)	B			<i>erroneus</i> Graham, 1987	S
<i>pausiris</i> (Walker, 1839)	B			<i>gordensis</i> Graham, 1987	B
<i>percaudatus</i> (Silvestri, 1920)			S	<i>mediterraneus</i> Graham, 1986	S
<i>phineus</i> (Walker, 1839)	B	M	S	<i>viridimaculatus</i> (Fullaway, 1955)	S
<i>phragmitinus</i> (Erdős, 1954)	B			<b>Anaprostocetus</b> Graham, 1987	
<i>planiusculus</i> (Thomson, 1878)	B		S	<i>acuminatus</i> (Ratzeburg, 1848)	B
<i>polygoni</i> (Erdős, 1954)	B				
<i>prolidice</i> Graham, 1987	B				

<b>Sigmophora</b> Rondani, 1867					
<i>brevicornis</i> (Panzer, 1804)	B	M	S		
<b>Aceratoneuromyia</b> Girault, 1917					
<i>evanescens</i> (Ratzeburg, 1848)	B	S			
<b>Pronotalia</b> Gradwell, 1957					
<i>carlinarum</i> (Szelényi et Erdős, 1951)	B	S			
<i>hungarica</i> (Erdős, 1955)	B	M	S		
<b>Crataepus</b> Foerster, 1878					
<i>marbis</i> (Walker, 1839)	B	M	S		
<b>Kocourekia</b> Bouček, 1966					
<i>debilis</i> (Ratzeburg, 1852)	M				
<b>Melittobia</b> Westwood, 1847					
<i>acasta</i> (Walker, 1839)	B				
EUDERINAE					
<b>Euderus</b> Haliday, 1844					
<i>agrili</i> Bouček, 1963	B				
<i>albitarsis</i> (Zetterstedt, 1838)	B	M	S		
<i>arenarius</i> Erdős, 1951	S				
<i>brevicornis</i> Bouček, 1963	B	M	S		
<i>caudatus</i> Thomson, 1878	B				
<i>viridis</i> Thomson, 1878	B	S			
<b>Parasecodella</b> Girault, 1915					
<i>obscura</i> (Thomson, 1878)	B	S			
<b>Allocerastichus</b> Masi, 1923					
<i>doderi</i> Masi, 1923	B	S			
<b>Astichus</b> Foerster, 1856					
<i>arithmeticus</i> Foerster, 1851)	B				
ENTEDONTINAE					
<b>Entedon</b> Dalman, 1820					
<i>abdera</i> Walker, 1839	B	S			
<i>apionis</i> Erdős, 1951	B				
<i>astragali</i> Erdős, 1951	S				
<i>biroi</i> Erdős, 1944	B	S			
<i>costalis</i> Dalman, 1820	B	M	S		
<i>crassiscapus</i> Erdős, 1944	B	S			
<i>diotimus</i> Walker, 1839	B	S			
<i>ergias</i> Walker, 1839				B	M
<i>euphorion</i> Walker, 1839				B	S
<i>hercyna</i> Walker, 1839				B	S
<i>insignis</i> Erdős, 1944				B	M
<i>kerteszi</i> Erdős, 1944				B	
<i>lanceolatus</i> Erdős, 1944				B	S
<i>leucocnemis</i> Erdős, 1944				B	
<i>lixi</i> Erdős, 1951				B	M
<i>longiventrosus</i>				S	
				Dalla Torre, 1898	
				<i>loti</i> Erdős, 1944	
				<i>methion</i> Walker, 1839	
				<i>molybdaenus</i> Erdős, 1944	
				<i>nigrini</i> Bouček, 1974	
				<i>nubilatus</i> Erdős, 1944	
				<i>pallicrus</i> Erdős, 1944	
				<i>pannonicus</i> Erdős, 1944	
				<i>parvicalcar</i> Thomson, 1878	
				<i>procioni</i> Erdős, 1944	
				<i>pseudonigritarsis</i> Erdős, 1944	
				<i>punctiscapus</i> Thomson, 1878	
				<i>spareta</i> Walker, 1839	
				<i>squamosus</i> Thomson, 1878	
				<i>subfumatus</i> Erdős, 1944	
				<i>ulmi</i> Erdős, 1944	
				<i>urticarii</i> Erdős, 1951	
				<b>Pediobius</b> Walker, 1846	
				<i>acantha</i> (Walker, 1839)	
				<i>alcaeus</i> (Walker, 1839)	
				<i>brachycerus</i> (Thomson, 1878)	
				<i>bruchicida</i> (Rondani, 1872)	
				<i>cassidae</i> Erdős, 1958	
				<i>chilaspidis</i> Bouček, 1965	
				<i>claviger</i> (Thomson, 1878)	
				<i>clita</i> (Walker, 1839)	
				<i>coxalis</i> Bouček, 1965	
				<i>crassicornis</i> (Thomson, 1878)	
				<i>deplanatus</i> Bouček, 1965	
				<i>epeus</i> (Walker, 1839)	
				<i>epigonus</i> (Walker, 1839)	
				<i>eubius</i> (Walker, 1839)	
				<i>facialis</i> (Giraud, 1863)	
				<i>foliorum</i> (Geoffroy in Fourcroy, 1875)	

<i>grunini</i> (Nikolskaja, 1954)	B							
<i>lysis</i> (Walker, 1839)		B M S						
<i>nigritarsis</i> (Thomson, 1878)	B							
<i>oviventris</i> Bouček, 1965	S							
<i>phragmitis</i> Bouček, 1965	S							
<i>phyllotretae</i> (Riley, 1885)	B M							
<i>polanensis</i> Bouček, 1965	S							
<i>pyrgo</i> (Walker, 1839)	B M S							
<i>saulinus</i> (Walker, 1839)	B M S							
<i>termerus</i> (Walker, 1839)	B S							
<i>tetratomus</i> (Thomson, 1878)	B							
<i>ulmi</i> (Erdös, 1954)	B							
<b>Mestocharis</b> Foerster, 1878								
<i>bimacularis</i> (Dalman, 1820)	B M							
<i>maculata</i> (Foerster, 1841)	M S							
<b>Kratoysma</b> Bouček, 1965								
<i>usticrus</i> (Erdös, 1954)	S							
<b>Derostenus</b> Westwood, 1833								
<i>gemmeus</i> Westwood, 1833	B	S						
<i>punctiscuta</i> Thomson, 1878	B	S						
<b>Chrysocharis</b> Foerster, 1856								
<i>acoris</i> (Walker, 1839)	B	S						
<i>albicoxis</i> Erdös, 1958	B	S						
<i>amanus</i> (Walker, 1839)	B	S						
<i>amasis</i> (Walker, 1839)	B							
<i>amyite</i> (Walker, 1839)	B	S						
<i>argyropezae</i> Graham, 1963	B							
<i>assis</i> (Walker, 1839)	B							
<i>avia</i> Hansson, 1985	B							
<i>budensis</i> Erdös, 1954	B							
<i>cerris</i> Erdös, 1961		S						
<i>chilo</i> (Walker, 1839)	B	S						
<i>chlora</i> Graham, 1963	B							
<i>clarkae</i> Yoshimoto, 1973	B							
<i>collaris</i> Graham, 1963	B							
<i>crassiscapus</i> (Thomson, 1878)	B	S						
<i>entedenoides</i> (Walker, 1872)	B							
<i>equiseti</i> Hansson, 1985	M							
<i>eurynota</i> Graham, 1968		S						
<i>idya</i> (Walker, 1839)	B							
<i>illustris</i> Graham, 1963	B							
<i>laomedon</i> (Walker, 1839)	B M S							
<i>laricinellae</i> (Ratzeburg, 1848)			B					
<i>larina</i> (Walker, 1839)			S					
<i>liriomyzae</i> Delucchi, 1954	B	S						
<i>loranthellae</i> Erdös, 1954			S					
<i>mediana</i> Foerster, 1861	B	S						
<i>moravica</i> (Maláč, 1943)	B M S							
<i>nautius</i> (Walker, 1846)	B	S						
<i>nephereus</i> (Walker, 1839)	B M S							
<i>nigricrus</i> (Thomson, 1878)	B							
<i>nitetis</i> (Walker, 1839)	B							
<i>nitidifrons</i> Graham, 1963	B							
<i>novellus</i> (Walker, 1839)	B	S						
<i>orbicularis</i> (Nees, 1834)	B							
<i>orchestis</i> (Ratzeburg, 1844)			B					
<i>paradoxa</i> Hansson, 1985	B							
<i>pentheus</i> (Walker, 1839)	B M S							
<i>phryne</i> (Walker, 1839)	B M							
<i>pilosa</i> Delucchi, 1954	B							
<i>pilicoxa</i> (Thomson, 1878)	B	S						
<i>polyzo</i> (Walker, 1839)	B M S							
<i>prodice</i> (Walker, 1839)	B M S							
<i>pubens</i> Delucchi, 1954	B							
<i>pubicornis</i> (Zetterstedt, 1838)	B M S							
<i>purpurea</i> Bukowski, 1938	B							
<i>submutica</i> Graham, 1963	B M S							
<i>truncatula</i> Graham, 1963	B							
<i>viridis</i> (Nees, 1834)	B M S							
<b>Grahamia</b> Erdös, 1966								
<i>clinius</i> (Walker, 1839)	B							
<i>tatrica</i> Erdös, 1966	S							
<b>Achrysocharoides</b> Girault, 1913								
<i>albiscapus</i> (Delucchi, 1957)	B M S							
<i>atys</i> (Walker, 1839)	B M S							
<i>butus</i> (Walker, 1839)	B M							
<i>cilla</i> (Walker, 1839)	B M S							
<i>latreillei</i> (Curtis, 1826)	B M S							
<i>nigricoxae</i> (Delucchi, 1954)	B M							
<b>Chrysotomomyia</b> Ashmead, 1904								
<i>albipes</i> (Kurdjumov, 1912)	M							
<i>albiscapus</i> (Erdös, 1954)	B M S							

<i>aratus</i> (Walker, 1838)	B	M	<i>salicis</i> Haliday, 1833	B	M	S
<i>arvensis</i> (Graham, 1963)	B		<i>versicolor</i> (Nees, 1834)	B		
<i>chlorogaster</i> (Erdös, 1966)	B		<b>Ionympha</b> Graham, 1959			
<i>clinias</i> (Walker, 1838)	B	M	<i>carne</i> (Walker, 1839)	B	S	
<i>cuprifrons</i> (Erdös, 1954)	B	M	<i>ochus</i> (Walker, 1839)	B	M	S
<i>formosa</i> (Westwood, 1833)	B	M	<b>Eugerium</b> Graham, 1959			
<i>lanassa</i> (Walker, 1839)	B		<i>isander</i> (Walker, 1839)	B	M	S
<i>nunbergi</i> (Szcepanski, 1960)	B		<b>Ceranisus</b> Walker, 1841			
<i>ovulorum</i> (Ratzeburg, 1848)	M		<i>lepidotus</i> Graham, 1963	B		
<i>ruforum</i> (Krausse, 1917)	B	S	<i>menes</i> (Walker, 1839)	B	S	
<i>smaragdula</i> (Graham, 1963)	B		<i>planitianus</i> Erdös, 1966	B	S	
<b>Closterocerus</b> Westwood, 1833			<i>pacuvius</i> (Walker, 1838)	B		
<i>trifasciatus</i> Westwood, 1833	B	M	<b>Thriopctenoides</b> Erdös, 1954			
<b>Teleopterus</b> Silvestri, 1914			<i>carbonarius</i> Erdös, 1954			S
<i>erxias</i> (Walker, 1848)	B		<b>Euderomphale</b> Girault, 1916			
<b>Desmatocharis</b> Graham, 1959			<i>callunae</i> Erdös, 1966	B		
<i>turcica</i> (Nees, 1834)	B		TRICHOGRAMMATIDAE			
<b>Holcopelte</b> Foerster, 1856			<b>Aphelinoidea</b> Girault, 1911			
<i>obscura</i> (Foerster, 1841)	B		<i>melanosoma</i> Novicky, 1940	B		
<i>sulciscuta</i> (Thomson, 1878)	B	M	<i>subexserta</i> Novicky, 1940			S
<b>Asecodes</b> Foerster, 1856			<b>Brachista</b> Haliday in Walker 1851			
<i>coronis</i> (Walker, 1838)	B	M	<i>pungens</i> (Mayr, 1904)	B	M	
<i>hyperion</i> Graham, 1963	B		<b>Centrobia</b> Foerster, 1856			
<i>mento</i> (Walker, 1839)	B	M	<i>doricha</i> (Walker, 1839)	B		
<b>Omphale</b> Haliday, 1833			<i>fumipennis</i> Blood, 1923	B		
<i>amirabilis</i> (Westwood, 1833)	B		<b>Chaetostricha</b> Haliday in Walker, 1851			
<i>aetius</i> (Walker, 1839)	B		<i>dimidiata</i> Walker, 1851	B		
<i>brevis</i> Graham, 1963	B		<b>Lathromeris</b> Foerster, 1856			
<i>chryseis</i> Graham, 1963	S		<i>scutellaris</i> Foerster, 1856	B	M	
<i>clymene</i> (Walker, 1839)	B		<b>Mirufens</b> Girault, 1915			
<i>clypealis</i> (Thomson, 1878)	B	S	<i>longicauda</i> (Blood, 1923)			S
<i>coilus</i> (Walker, 1839)	B	M	<b>Monorthrochaeta</b> Blood, 1923			
<i>epaphus</i> (Walker, 1839)	S		<i>nigra</i> Blood, 1923	B		
<i>lugens</i> (Nees, 1834)	B	M	<b>Oligosita</b> Haliday, 1851			
<i>matrana</i> Erdös, 1954	B	S	<i>collina</i> Walker, 1851	B		
<i>radialis</i> (Thomson, 1878)	B	S	<i>krygeri</i> Girault, 1929	B		
<i>rubigus</i> (Walker, 1839)	B		<i>lutulenta</i> Novicky, 1940	M		
			<i>pallida</i> Kryger, 1918	B		

<i>subfasciata</i> Westwood, 1879	M	<i>ater</i> Foerster, 1841	B
<b>Ophioneurus</b> Ratzeburg, 1852		<i>litoralis</i> (Haliday, 1833)	B M S
<i>signatus</i> Ratzeburg, 1852	B	<i>longicornis</i> Nees, 1834	B S
<b>Paracentrobia</b> Howard, 1897		<i>pictus</i> (Haliday, 1833)	B S
<i>monotricha</i> (Novicky, 1940)	B	<b>Ooctonus</b> Haliday, 1833	
<b>Poropoea</b> Foerster, 1851		<i>askhamensis</i> Hincks, 1952	B
<i>bohemica</i> (Novicky, 1936)	B	<i>hemipterus</i> Haliday, 1833	B S
<i>stollwercki</i> Foerster, 1851	M	<i>heterotomus</i> Foerster, 1847	B
<b>Prestwichia</b> Lubbock, 1863		<i>insignis</i> Haliday, 1833	B S
<i>aquatica</i> Lubbock, 1863	B	<i>soykai</i> Hincks, 1952	B S
<b>Trichogramma</b> Westwood, 1833		<i>vulgatus</i> Haliday, 1833	B
<i>cephalciae</i> Hochmut et Martinek, 1963	B M	<b>Erythmelus</b> Enock, 1909	
<i>embryophagum</i> (Hartig, 1838)	B M S	<i>dichromocnemus</i> Novicki, 1953	B
<i>evanescens</i> Westwood, 1833	B M S	<b>Anaphes</b> Haliday, 1833	
<b>Tumidiclava</b> Girault, 1911		<i>latissimus</i> (Soyka, 1956)	B
<i>bimaculata</i> (Blood, 1923)	M	<i>pallidus</i> (Soyka, 1956)	B
<i>minuscula</i> Novicky, 1940	S	<i>rufoniger</i> (Soyka, 1956)	B
<b>Ufens</b> Girault, 1911		<i>stammeri</i> (Soyka, 1946)	B
<i>foersteri</i> (Kryger, 1918)	B	<b>Eustochus</b> Haliday, 1833	
MYMARIDAE		<i>atripennis</i> (Curtis, 1832)	B
ALAPTINAE		<b>Mymar</b> Curtis, 1832	
<b>Litus</b> Haliday, 1833		<i>pulchellum</i> Curtis, 1832	B
<i>cynipseus</i> Haliday, 1833	B	<b>Stephanodes</b> Enock, 1909	
MYMARINAE		<i>similis</i> (Foerster, 1847)	B
<b>Arescon</b> Walker, 1846		<b>Polynema</b> Haliday, 1833	
<i>rufulum</i> (Foerster, 1847)	B	<i>gracile</i> (Nees, 1834)	B
<b>Gonatocerus</b> Nees, 1834		<i>flavipes</i> Walker, 1846	B
<i>acuminatus</i> (Walker, 1846)	B S	<i>fuscipes</i> Haliday, 1833	S
		<i>pusillum</i> Haliday, 1833	B
		<i>waterhousei</i> Hincks, 1950	B
		MYMAROMMATIDAE	
		<b>Palaeomyrmecomyia</b> Meunier, 1901	
		<i>anomalum</i> (Blood et Kryger, 1922)	B



## CYNIPOIDEA

PAVEL MIKULA

In Czechoslovakia, the superfamily Cynipoidea is among the less known groups of hymenopterous insects. More data are available on gall-producing than on parasitic species. Data on the occurrence of species of the family Cynipidae are available especially in the literature on forestry entomology and in the papers by Baudyš (1912, 1962) and Bayer (1910, 1946). Both authors mainly described the plant galls produced by Cynipoidea, not the insects that produce them. Their extensive collections of galls are deposited in the Moravian Museum in Brno. After 1950, the Cynipoidea was temporarily studied by P. Masner, who described several new species. A large part of his collection is deposited in the National Museum in Prague.

The present check-list is only a tentative one. Systematic faunal investigations will supplement it by additional genera and species, particularly of the parasitic families of Cynipoidea. New discoveries can be expected to be made in the whole territory of Czechoslovakia.

The Cynipoidea includes parasitic species whose larvae attack larvae of Coleoptera, puparia of Diptera (Eucoilidae, Figitidae), or Neuroptera (Anacharitidae). The species of Ibalidae parasitize larvae of Siricidae, those of the subfamily Charipinae parasitize aphids. Phytophagous species of the family Cynipidae produce galls on numerous wild-growing and cultivated plants.

In compiling the list, the classification of taxa according to Kierych (1966, 1979) and Ionescu (1973) has been used.

References: Baudyš, E., 1912: Pro Čechy nové hálky. *Acta Soc. Ent. Boh.* 9: 118—120. — Baudyš, E. 1962: Zoocecidie Osoblažska. *Sbor. vlast. mus.* Olomouc 23: 73—83. — Bayer, E. 1910: Les Zoocécides de la Bohême. *Marcellia Anellino* 9: 63—158. — Bayer, E., 1946: Hálky Olomoucká ze sběru Jos. Otruby 1944. *Acta Soc. Zool. Csl.* 10: 81—89. — Ionescu, M. A., 1973: Biologia galeror — Monografie cecidologica. Editura Acad. RSR, Bucureşti, 178 pp. — Kierych, E., 1966: Materialen zur Kenntnis der phytophagischen Gallwespen (Hym. Cynip.). *Pol. Fragm. faun.*, 13: 159—170. — Kierych, E., 1979: Galasówkowate Cynipoidea. Catalogus Faunae Poloniae, PAN. Inst. Zool. Warszawa, 26, Vol. 6: 103 pp.

CYNIPODEA			
IBALIIDAE			
<b>Ibalia</b> Latreille, 1802			
<i>drewseni</i> Borries, 1891	B M S		
<i>leucospoides</i> (Hochenwarth, 1785)	B M S		
ANACHARITIDAE			
<b>Anacharis</b> Dalman, 1823			
<i>ensifer</i> Walker, 1835	B M		
<i>eucharoides</i> Dalman, 1823	B M S		
<i>immunis</i> Walker, 1835	B M		
<i>rotiventris</i> (Hartig, 1840)	B M		
<i>tincta</i> Walker, 1835	B M		
<i>typica</i> Walker, 1835	B M S		
ASPICERIDAE			
<b>Callaspidia</b> Dahlbom, 1842			
<i>dufouri</i> Giraud, 1860	B M		
FIGITIDAE			
<b>Sarothrus</b> Hartig, 1840			
<i>areolatus</i> Hartig, 1840	B M S		
<i>haemiscutellaris</i> P. Masner, 1964	B S		
<i>longitarsus</i> (Reinhard, 1860)	B M		
<i>tibialis</i> (Zetterstedt, 1838)	B M		
<b>Melanips</b> Giraud, 1860			
<i>granulatus</i> (Hartig, 1841)	B M		
<i>opacus</i> (Hartig, 1840)	B		
<i>parvus</i> (Hartig, 1840)	B S		
<b>Lonchidia</b> Thomson, 1862			
<i>clavicornis</i> Thomson, 1862	B S		
<i>lisonota</i> Thomson, 1862	B		
<b>Figites</b> Latreille, 1802			
<i>anthomyiarum</i> Bouché, 1834	B M		
<i>erythropus</i> (Walker, 1835)	B		
<i>scutellaris</i> (Rossi, 1794)	B M S		
<i>susterai</i> P. Masner, 1965	M S		
EUCOILIDAE			
<b>Kleidotoma</b> Westwood, 1833			
		brevicornis	Thomson, 1862
			B
		caledonica	(Cameron, 1888)
			B S
		erythropus	Thomson, 1877
			B
		formicaria	(Kieffer, 1902)
			B
		striaticollis	(Cameron, 1888)
			B
		Trybliographa	Foerster, 1869
		rapae	(Westwood, 1835)
			B M S
		simulatrix	(Ruthe, 1859)
			B M
		Rhopromeris	Foerster, 1869
		graciliclava	(Kieffer, 1902)
			B
		Pseudeucoila	Ashmead, 1903
		ciliaris	(Zetterstedt, 1838)
			B M S
		coronata	(Hartig, 1841)
			B
		insignis	(Giraud, 1860)
			B M
		Cothonaspis	Hartig, 1840
		albipennis	(Thomson, 1862)
			B
		giraudi	(Dalla Torre et Kieffer, 1910)
			B M S
		nigricornis	Kieffer, 1902
			B M
		Microstilba	Foerster, 1869
		tibialis	Kieffer, 1901
			B
		Disorygma	Foerster, 1869
		divulgata	Foerster, 1869
			B M
		Diglyphosema	Foerster, 1869
		punctata	Kieffer, 1901
			B
		Eucoila	Westwood, 1833
		maculata	Hartig, 1841
			B M S
		ALLOXYSTIDAE	
		Alloxysta	Foerster, 1869
		circumscripta	(Hartig, 1841)
			B
		crassa	(Cameron, 1889)
			B M
		curvata	(Kieffer, 1902)
			B

<i>flavicornis</i> (Hartig, 1841)	B M S	<b>Trigonaspis</b> Hartig, 1840
<i>fracticornis</i> (Thomson, 1862)	B M	<i>megaptera</i> (Panzer, 1801) B M S
<i>macrophadna</i> (Hartig, 1841)	B M S	<i>synaspis</i> (Hartig, 1841) M
<i>minuta</i> (Hartig, 1840)	B M	<b>Biorhiza</b> Westwood, 1840
<i>victrix</i> (Westwood, 1833)	B	<i>pallida</i> (Olivier, 1791) B M S
<b>Pezophycta</b> Foerster, 1869		<b>Aphelonyx</b> Mayr, 1881
<i>brachyptera</i> (Hartig, 1840)	B M	<i>cerricola</i> (Giraud, 1859) ?
<i>cursor</i> (Hartig, 1840)	B	
CYNIPIDAE		
<b>Diplolepis</b> Geoffroy, 1762		<b>Andricus</b> Hartig, 1840
<i>centifoliae</i> (Hartig, 1840)	B M S	<i>aestivalis</i> Giraud, 1859 M S
<i>eglantariae</i> (Hartig, 1840)	B M S	<i>albopunctatus</i> Schlechtendal, 1870 M
<i>mayri</i> (Schlechtendal, 1876)	B M S	<i>paradoxus</i> (Radoszkowski, 1866) M
<i>rosae</i> (Linnaeus, 1758)	B M S	<i>amblycera</i> (Giraud, 1859) M
<i>rosarum</i> (Giraud, 1859)	B M	<i>aries</i> (Giraud, 1859) M S
<i>spinossimae</i> (Giraud, 1859)	B M S	<i>caliciformis</i> (Giraud, 1859) B M S
<b>Neuroterus</b> Hartig, 1840		<i>callidoma</i> (Hartig, 1841) M S
<i>aprilinus</i> (Giraud, 1859)	B M	<i>caputmedusae</i> (Hartig, 1843) M S
<i>fumipennis</i> Hartig, 1841	B M S	<i>conglomerata</i> (Giraud, 1859) M S
<i>glandiformis</i> (Giraud, 1859)	S	<i>conifica</i> (Hartig, 1843) M
<i>laevisculus</i> Schenck, 1863	B M S	<i>coriaria</i> (Hartig, 1843) M S
<i>lanuginosus</i> Giraud, 1859	M	<i>coronata</i> (Giraud, 1859) M
<i>macropterus</i> (Hartig, 1843)	M S	<i>corruptrix</i> (Schlechtendal, 1870) B M
<i>numismalis</i> (Fourcroy, 1885)	B M S	<i>curvator</i> Hartig, 1840 B M S
<i>quercusbaccarum</i> (Linnaeus, 1758)	B M S	<i>cydoniae</i> (Giraud, 1859) M S
<b>Cynips</b> Linnaeus, 1758		<i>fecundatrix</i> (Hartig, 1840) B M S
<i>agama</i> Hartig, 1841	M S	<i>gallaetinctoriae</i> (Olivier, 1791) M S
<i>cornifex</i> Hartig, 1840	M S	<i>giraudianus</i> (Dalla Torre et Kieffer, 1910) M
<i>disticha</i> Hartig, 1841	M S	<i>glandulae</i> (Hartig, 1840) M
<i>divisa</i> Hartig, 1840	B M S	<i>hartigi</i> (Hartig, 1843) B M S
<i>longiventris</i> Hartig, 1840	B M S	<i>hungarica</i> (Hartig, 1843) M S
<i>quercusfolii</i> Linnaeus, 1758	B M S	<i>inectorius</i> (Hartig, 1843) M
		<i>inflator</i> Hartig, 1840 B M S
		<i>kollari</i> (Hartig, 1843) B M S
		<i>lignicola</i> (Hartig, 1840) B M S
		<i>lucidus</i> (Hartig, 1843) M
		<i>marginalis</i> (Schlechtendal, 1876) S
		<i>mayri</i> (Wachtl, 1881) B M
		<i>multiplicatus</i> Giraud, 1859 M S
		<i>ostrea</i> (Hartig, 1840) B M S

<i>polycera</i> (Giraud, 1859)	M S	<i>jaceae</i> (Schenck, 1863)	B M
<i>quadrilineatus</i> Hartig, 1840	B M	<i>kerneri</i> (Wachtl, 1891)	M
<i>quercuscalicis</i> (Burgsdorf, 1783)	B M S	<i>minor</i> Hartig, 1840	B M S
<i>quercuscorticis</i> (Linnaeus, 1761)	M S	<i>papaveris</i> (Perris, 1840)	B M S
<i>quercusradicis</i> (Fabricius, 1798)	B M S	<i>salviae</i> (Giraud, 1859)	B M S
<i>quercusramuli</i> (Linnaeus, 1761)	M S	<b>Liposthenes</b> Foerster, 1869	
<i>rhyzomae</i> (Hartig, 1843)	M	<i>glechomae</i> (Linnaeus, 1758)	B M S
<i>seminationis</i> (Giraud, 1859)	M S	<b>Periclistus</b> Foerster, 1869	
<i>solitarius</i> (Fonscolombe, 1832)	B M S	<i>brandti</i> (Ratzeburg, 1831)	M S
<i>testaceipes</i> (Hartig, 1841)	B M S	<i>caninae</i> (Hartig, 1840)	B M
<b>Diastrophus</b> Hartig, 1840		<b>Ceroptres</b> Hartig, 1840	
<i>mayri</i> Reinhard, 1876	B M S	<i>arator</i> Hartig, 1841	B M S
<i>rubi</i> (Bouché, 1834)	B M S	<b>Saphonecrus</b> Dalla Torre et Kieffer, 1910	
<b>Xestophanes</b> Foerster, 1869		<i>connatus</i> (Hartig, 1840)	B M
<i>brevitarsis</i> (Thomson, 1877)	B M S	<i>undulatus</i> (Mayr, 1876)	B M S
<i>potentillae</i> (Retzius, 1783)	B M S	<b>Synergus</b> Hartig, 1840	
<i>szepligetii</i> Ballás, 1948	M S	<i>albipes</i> Hartig, 1841	B M S
<b>Isocolus</b> Foerster, 1869		<i>apicalis</i> Hartig, 1841	B M
<i>rogenhoferi</i> Wachtl, 1881	B M S	<i>clandestinus</i> Eady, 1952	B
<i>scabiosae</i> (Giraud, 1859)	B M S	<i>evanescens</i> Mayr, 1873	B M S
<b>Aulacidea</b> Ashmead, 1897		<i>gallaepommiformis</i> (Fonscolombe, 1832)	B M S
<i>hieracii</i> (Linnaeus, 1758)	B M S	<i>hayneanus</i> (Ratzeburg, 1831)	B M
<i>macula</i> Forsius, 1921	M S	<i>incrassatus</i> Hartig, 1840	B
<i>pigeoti</i> (Kieffer, 1902)	B	<i>nervosus</i> Hartig, 1840	B M S
<i>pilosellae</i> (Kieffer, 1901)	M S	<i>pallicornis</i> Hartig, 1841	B M S
<i>tragopogonis</i> (Thomson, 1877)	B M S	<i>pallidipennis</i> Mayr, 1873	B M
<b>Phanacis</b> Foerster, 1860		<i>radiatus</i> Mayr, 1873	B M
<i>centaureae</i> Foerster, 1860	B M S	<i>reinhardi</i> Mayr, 1873	B M S
<i>hypocoeridis</i> (Kieffer, 1887)	M S	<i>ruficornis</i> Hartig, 1840	B M
<i>lamsanae</i> (Perris, 1873)	S	<i>thaumacerus</i> (Dalman, 1823)	B M S
<b>Timaspis</b> Mayr, 1881		<i>tscheki</i> Mayr, 1873	B M
<i>cichorii</i> Ballás, 1948	B M S	<i>umbraculus</i> (Olivier, 1791)	B M S
<i>papaveris</i> Kieffer, 1905	B M S	<i>variabilis</i> Mayr, 1873	M S
<b>Aylax</b> Hartig, 1840		<i>vulgaris</i> Hartig, 1840	B M
<i>hypocheridis</i> Kieffer, 1902	M	<i>xanthocerus</i> Hartig, 1841	M
		<b>Pediaspis</b> Tischbein, 1852	
		<i>aceris</i> (Gmelin in Linnaeus, 1790)	M

## EVANIOIDEA

JOSEF ŠEDIVÝ

Faunal studies of the superfamily Evanioidea have no tradition in Czechoslovakia. The hitherto findings come from accidental collections only. So far, 24 species have been recorded in Czechoslovakia. The genera known to occur in the territory of Czechoslovakia were elaborated by Bouček (1957).

Of the family Evanidae, only two genera containing two species are known to occur in Czechoslovakia so far. The species develop in the egg cases of cockroaches. Discoveries of additional species may be expected in the warm southern parts of Czechoslovakia.

Until recently, data on the distribution of species of the family Aulacidae in Czechoslovakia have been sporadic. The larvae are thought to be parasitoids of xylophagous Coleoptera or Hymenoptera.

The author evaluated the hitherto findings of species of the family Gasteruptiidae (Šedivý, 1958). The larvae are predators of sphecids and solitary bees.

This check-list was prepared on the basis of the collections of the National Museum in Prague, Moravian Museum in Brno and the private collection of the present author.

New discoveries of species of the families Aulacidae and Gasteruptiidae may be expected in the whole territory of Czechoslovakia.

The classification in this list is based mainly on works Oehlke (1983, 1984).

References: Bouček, Z. (in Kratochvíl, J. et al.), 1957: Sprušky — Evanioidea. Klíč zvířeny ČSR, II, Praha: 190—191. — Oehlke, J., 1983: Revision der europäischen Aulacidae (Hymenoptera — Evanioidea). *Beitr. Ent.* 33: 439—447. — Oehlke, J., 1984: Beiträge zur Insektenfauna der DDR: Hymenoptera — Evanioidea, Stephanoidea, Trigonalyoidea. *Faun. Abhandl.* 11: 161—190. — Šedivý, J., 1958: Tschechoslowakische Arten der Gasteruptioniden, Hym. Čas. Čsl. spol. ent. 55, 1: 34—43.

EVANIOIDEA	Brachygaster Leach, 1819	
EVANIIDAE	<i>minuta</i> (Olivier, 1791) B M S	
<b>Prosevania</b> Kieffer, 1911	AULACIDAE	
<i>fuscipes</i> (Illiger, 1807)	<b>Aulacus</b> Jurine, 1807	
<i>punctata</i> (Brulle, 1833)	<i>striatus</i> Jurine, 1807	B M S

<b>Pristaulacus</b> Kieffer,			
1900			
<i>Aulacostethus</i> Philippi, 1873			
<i>bimaculatus</i> Kieffer, 1900	*B	S	
<i>chlapowskii</i> Kieffer, 1900		*B*M	
<i>gloriator</i> (Fabricius, 1804)		B M S	
<i>obscuripennis</i> (Westwood, 1841)		B M S	
<b>GASTERUPTIIDAE</b>			
<b>Gasteruption</b> Latreille, 1796			
<i>assector</i> (Linnaeus, 1758)	B M S		
<i>bidentulum</i> (Thomson, 1883)			
<i>diversipes</i> (Abeille, 1879)	M S		
<i>erythrostomum</i> (Dahlbom, 1831)	B M S		
<i>freyi</i> (Tournier, 1877)	B	M S	
<i>goberti</i> (Tournier, 1877)	M		
<i>hastator</i> (Fabricius, 1804)	B	M S	
<i>hungaricus</i> Szépligeti, 1895		M S	
<i>jaculator</i> (Linnaeus, 1758)		B M S	
<i>granulithorax</i> (Tournier, 1877)			
<i>laticeps</i> (Tournier, 1877)	M	S	
<i>minutum</i> (Tournier, 1877)	B	M	
<i>nigrescens</i> Schletterer, 1889		S	
<i>opacum</i> (Tournier, 1877)	B	M S	
<i>paternum</i> Schletterer, 1889	M		
<i>pedemontanum</i> (Tournier, 1877)	B	M S	
<i>pyrenaicum</i> (Guérin, 1844)	M	S	
<i>tibiale</i> Schletterer, 1885	M	S	
<i>tournieri</i> Schletterer, 1885	B	M S	

## PROCTOTRUPOIDEA, CERAPHRONOIDEA

JAN MACEK

The two superfamilies are heterogeneous and taxonomically poorly studied groups, little being known of the occurrence of their species in Czechoslovakia. Hence, the lists of species given below are incomplete, and discoveries of new genera and species may be expected in the whole of Czechoslovakia. The first reports on the occurrence of Proctotrupoidea and Ceraphronoidea in the territory of Bohemia are contained in the papers compiled by Kirchner (e.g. 1855) but they are inaccurate.

In Czechoslovakia, occasional faunal investigations of the two groups began in the nineteen-twenties by the papers of A. A. Ogloblin, F. Gregor sen. and G. Szelényi. These entomologists described certain new genera and species and published faunal data on their distribution. The fundaments of regular faunal investigations were not laid until after 1944 when Tomšík (1944, 1946, 1947) worked out taxonomical and faunal data on certain genera of Diapriidae and Proctotrupidae. His collection is deposited in the Moravian Museum in Brno. Later, the taxonomy and faunistics of the two superfamilies were studied by L. Masner (1957) who contributed a great deal to the knowledge of their distribution in Czechoslovakia. His collection until 1968 is deposited in the National Museum in Prague. The author of this chapter has been studying the fauna of Proctotrupoidea in central and southern Bohemia, southern Moravia, and in the mountains of central Slovakia. His collection is deposited in the National Museum in Prague.

The species of Proctotrupoidea are parasitoids of larvae of Coleoptera (Carabidae, Staphylinidae, Elateridae, etc.), Diptera (especially Cecidomyiidae), Aleyrodidae or, rarely, Chilopoda. Part of the species parasitize eggs of insects and spiders. The species of Ceraphronoidea are hyperparasitoids of larvae of Aphidiidae, Braconidae, Ichneumonidae, Bethylidae, Dryinidae and Tachinidae. They also parasitize species of Aleyrodidae and certain Neuroptera, gall-producing Cynipoidea, and myrmecophilous Diptera.

The classification of higher taxa in this list is based on the papers by Kozlov (1978), Masner (1976), Nixon (1957) and Pschorn-Walcher (1958).

References: Kirchner, L., 1855: Die Gallenauswüchse des budweiser Kreises nebst nomineller Angabe der Gallen Erzeuger und deren Schmarotzer. *Lotos* 5, Praha: 13—33. Kozlov, M. A., 1978: Proctotrupoidea. In. Opred. nas. evr. SSSR 3 (2): 538—664. Masner, L., 1956: The first preliminary report on occurrence of genera of the group Proctotru-

poidea in Czechoslovakia. Scelionidae. *Acta faun. ent. Mus. nat. Pragae*, 1: 99–126. Masner, L., 1957: The first preliminary report on the occurrence of genera of the group Proctotrupoidea in Czechoslovakia. *Acta faun. ent. Mus. nat. Pragae* 2: 83–106. Masner, L., 1976: A revision of the Ismarinae of the New World. Can. Ent. 108: 1243–1266. Nixon, G. E. J., 1957: Diapriidae (Belytiniae). Handb. Ident. Brit. Insect. VIII/3: 107 pp. Pschorn-Walcher, H., 1958: Vorläufige Gliederung der palaearktischen Proctotrupidae. Mitt. Schweiz. Ent. Ges., 31: 57–64. Tomšík, B., 1944: Rod Phaenoserphus (Hym. Serph.) v našich zemích. Fol. ent., 5: 73–76. Tomšík, B., 1946: Rody Galesus a Schizogalesus. Fol. ent. 9: 119–132. Tomšík, B., 1947: Rod Spilomicrus Westw. (Hym., Proct.). Fol. Ent., 10: 33–43.

PROCTOTRUPOIDEA			
HELORIDAE			
<b>Helorus</b> Latreille, 1802			
<i>anomalipes</i> (Panzer, 1798)	B M S		
<i>nigripes</i> Foerster, 1856	B M		
<i>rugosus</i> Thomson, 1856			
<i>ruficornis</i> Foerster, 1856	B M		
<i>striolatus</i> Cameron, 1906	B M		
PROCTOTRUPIDAE			
<b>Disogmus</b> Foerster, 1856			
<i>basalis</i> (Thomson, 1857)	M S		
<i>carinatus</i> Kieffer, 1907	M		
<i>nigripennis</i> (Thomson, 1857)	B		
<b>Cryptoserpheus</b> Kieffer, 1907			
<i>aculeator</i> (Haliday, 1839)	B		
<i>cumaeus</i> Nixon, 1938	B		
<i>laricis</i> (Haliday, 1839)	M		
<i>longicalcar</i> Kieffer, 1908	B M		
<b>Brachyserphus</b> Hellén, 1941			
<i>parvulus</i> (Nees, 1834)	B		
<b>Proctotrupes</b> Latreille, 1796			
<i>gladiator</i> (Haliday, 1839)	B M S		
<i>hofferi</i> Tomšík, 1944			
<i>gravidator</i> (Linnaeus, 1758)	B M S		
<b>Phaenoserphus</b> Kieffer, 1908			
<i>calcator</i> (Haliday, 1839)	B M S		
<i>curtipennis</i> (Haliday, 1839)	B M		
<i>dubiosus</i> Nixon, 1938	B		
<i>fuscipes</i> (Haliday, 1839)	B		
<i>gregori</i> Tomšík, 1942	M		
<i>micrurus</i> Kieffer, 1908	B M		
<i>pallipes</i> (Latreille, 1809)	B M S		
<b>vexator</b> Nixon, 1938	B		
<b>viator</b> (Haliday, 1839)	B M S		
<b>Codrus</b> Panzer, 1801			
<i>ater</i> (Nees, 1834)	B M		
<i>fumipennis</i> (Kieffer, 1908)	M S		
<i>ligatus</i> (Nees, 1834)	B M S		
<i>brevicornis</i> (Haliday, 1839)	B M S		
<i>longicornis</i> (Nees, 1834)	M		
<i>microcerus</i> (Kieffer, 1908)	B M		
<i>niger</i> Panzer, 1801	B M S		
<b>Paracodrus</b> Kieffer, 1907			
<i>apterogynus</i> (Haliday, 1839)	M		
<b>Parthenocordus</b>			
Pschorn-Walcher, 1958			
<i>elongatus</i> (Haliday, 1839)	B		
DIAAPRIIDAE			
<b>Ismarus</b> Haliday, 1835			
<i>flavicornis</i> (Thomson, 1858)	B		
<i>halidayi</i> Foerster, 1850	B		
<i>moravicus</i> Ogloblin, 1925	B M		
<i>rugulosus</i> Foerster, 1850	B M S		
<b>Anommatium</b> Foerster, 1856			
<i>ashmeadi</i> Mayr, 1904	B M S		
<b>Atelopsilus</b> Kieffer, 1908			
<i>ciliatus</i> (Thomson, 1858)	B		
<b>Synacra</b> Foerster, 1856			
<i>brachialis</i> (Nees, 1834)	B M		
<i>sociabilis</i> (Kieffer, 1904)	B M		
<b>Sundholmiella</b> Hedquist, 1975			
<i>giraudi</i> (Kieffer, 1910)	B		
<b>Psilomma</b> Foerster, 1856			

<i>dubia</i> Kieffer, 1908	B M	<i>bicolor</i> Jurine, 1807	B M
<i>fuscicornis</i> Kieffer, 1908	B M	<i>brachyptera</i> Thomson, 1858	B M S
<i>fusciscapus</i> Foerster, 1861	B M S	<i>carinifrons</i> (Kieffer, 1909)	B S
<b>Cardiopsilus</b> Kieffer, 1908		<i>depressa</i> Thomson, 1858	B M S
<i>heterocerus</i> (Haliday, 1857)	B S	<i>elegans</i> Kieffer, 1909	B M S
<i>nervosus</i> (Thomson, 1858)	B M S	<i>lativentris</i> Cameron, 1887	B M S
<i>productus</i> Kieffer, 1908	B M	<i>lubrica</i> Kieffer, 1909	B M S
<b>Pantolyta</b> Foerster, 1856		<i>pelias</i> Nixon, 1957	B M S
<i>atrata</i> Foerster, 1856		<i>rugosicollis</i> Kieffer, 1909	B M
<i>pallida</i> Kieffer, 1908	B M S	<i>seron</i> Nixon, 1957	B M S
<i>stylata</i> Kieffer, 1908	B M	<b>Diphora</b> Foerster, 1856	
<i>vernalis</i> Nixon, 1957	B	<i>westvoodi</i> Foerster, 1856	B M S
<b>Acropista</b> Foerster, 1856		<b>Pantoclis</b> Foerster, 1856	
<i>flavipes</i> Kieffer, 1909	B	<i>brevicornis</i> Kieffer, 1909	B M
<i>flaviventris</i> (Thomson, 1858)	B M S	<i>eulimine</i> Nixon, 1957	B M S
<i>macrocera</i> (Thomson, 1858)	B M S	<i>leviventris</i> (Kieffer, 1907)	B
<i>marginalis</i> Kieffer, 1909	B M	<i>longipennis</i> (Thomson, 1858)	B M S
<i>nigrocincta</i> Kieffer, 1909	B M	<i>macrotoma</i> Kieffer, 1909	B M
<i>radialis</i> Hellén, 1964	B S	<i>merope</i> Nixon, 1957	B
<i>rufiventris</i> (Kieffer, 1907)	B S	<i>obscuripes</i> Kieffer, 1907	B M S
<i>sciariadora</i> (Kieffer, 1907)	B M S	<i>scotica</i> (Kieffer, 1909)	B M S
<i>fuscicornis</i> (Kieffer, 1909)	B M S	<i>similis</i> (Thomson, 1858)	B M S
<b>Opazon</b> Haliday, 1857		<i>subatricornis</i> Kieffer, 1909	B M
<i>Rhynchosilus</i> Kieffer, 1908		<i>sulcata</i> (Thomson, 1858)	B M S
<i>parvulum</i> (Haliday, 1857)	B M	<i>trisulcata</i> Kieffer, 1907	B M
<i>incrassatum</i> (Thomson, 1858)	M	<b>Zygota</b> Foerster, 1856	
<i>apertum</i> (Kieffer, 1910)	B M	<i>brevipennis</i> (Kieffer, 1908)	B M
<b>Oxylabis</b> Foerster, 1856		<i>brevinervis</i> (Kieffer, 1909)	B
<i>bisulca</i> (Nees, 1834)	B M S	<i>breviscula</i> (Thomson, 1858)	B M S
<i>cameroni</i> (Kieffer, 1909)	B	<i>claviscapa</i> (Thomson, 1858)	B S
<i>thomsoni</i> Kieffer, 1916	B M S	<i>croton</i> Nixon, 1957	B S
<b>Aprestes</b> Nixon, 1957		<i>cursor</i> (Kieffer, 1908)	B M S
<i>aberrans</i> Nixon, 1957	B M	<i>excisipes</i> (Kieffer, 1908)	B M S
<b>Synbelyta</b> Hellén, 1964		<i>excisor</i> (Zetterstedt, 1838)	B S
<i>fuscipennis</i> (Thomson, 1858)	B	<i>fossulata</i> (Thomson, 1858)	B S
<b>Belyta</b> Jurine, 1807		<i>fuscata</i> (Thomson, 1858)	B
<i>abrupta</i> Thomson, 1858	B M S	<i>macroneura</i> (Kieffer, 1909)	B S

<i>microtoma</i> (Kieffer, 1909)	B	M	S
<i>nigra</i> (Thomson, 1858)	B		S
<i>praetor</i> Nixon, 1957	B		S
<i>ruficornis</i> (Curtis, 1831)	B	M	S
<i>soluta</i> (Kieffer, 1907)	B	M	
<i>spinosa</i> (Kieffer, 1908)	B		S
<i>subaptera</i> (Thomson, 1858)	B	M	
<b>Paroxylabis</b> Kieffer, 1907			
<i>semirufa</i> Kieffer, 1907	B	M	S
<b>Acanthopsilus</b> Kieffer, 1908			
<i>marshalli</i> (Kieffer, 1907)	B		S
<b>Aclista</b> Foerster, 1856			
<i>analis</i> (Kieffer, 1910)	B	M	
<i>angusta</i> (Kieffer, 1909)	B	M	
<i>bitensis</i> (Kieffer, 1909)	B		
<i>cantianus</i> (Curtis, 1831)	B	M	
<i>clavicornis</i> (Kieffer, 1909)	B	M	
<i>elevata</i> (Thomson, 1858)	B	M	
<i>folia</i> Nixon, 1957	B	M	
<i>fractinervis</i> (Kieffer, 1910)			S
<i>insolita</i> Nixon, 1957	B		
<i>janssoni</i> Nixon, 1957	B	M	
<i>mycale</i> Nixon, 1957	B	M	
<i>neglecta</i> (Kieffer, 1909)	B	M	
<i>prolongata</i> (Kieffer, 1907)	B	M	
<i>rufopetiolata</i> (Nees, 1834)	B	M	S
<i>scutellaris</i> (Thomson, 1858)	B	M	
<i>soror</i> (Kieffer, 1909)	B		
<i>striolata</i> (Thomson, 1858)	B	M	
<i>subfusciformis</i> (Kieffer, 1910)	B	M	
<b>Cinetus</b> Jurine, 1807			
<i>atriceps</i> Kieffer, 1910	B		S
<i>cameroni</i> Kieffer, 1910	B	M	
<i>iridipennis</i> Lepeletier, 1825	B	M	
<i>fuliginosus</i> Curtis, 1831	B	M	
<b>Leptonetus</b> Masner, 1964			
<i>compressus</i> (Thomson, 1858)	B	M	
<i>monilicornis</i> (Kieffer, 1910)	B	M	S
<i>politus</i> (Thomson, 1858)	B	M	
<b>Scorpioteleia</b> Ashmead, 1897			
<i>cebes</i> (Nixon, 1957)			B
<i>compressa</i> (Kieffer, 1910)	B	M	S
<i>longipetiolata</i> (Thomson, 1858)			B M
<i>longiventris</i> (Kieffer, 1910)			B M
<b>Paramesius</b> Westwood, 1832			
<i>brevipennis</i> Kieffer, 1911	B		
<i>dolichocerus</i> Kieffer, 1911	B	M	
<i>dolosus</i> Kieffer, 1911			B M
<i>inermis</i> Kieffer, 1910	B		
<i>rufipes</i> Westwood, 1832	B	M	
<i>unifoveatus</i> Kieffer, 1911	B	M	
<b>Spilomicrus</b> Westwood, 1832			
<i>carinatus</i> Kieffer, 1911	B	M	
<i>cursor</i> Kieffer, 1911			M
<i>flavipes</i> Thomson, 1858	B		
<i>inaequalis</i> Tomšík, 1947	B		
<i>integer</i> Thomson, 1858			B
<i>integer</i> fossus Tomšík, 1947			B
<i>modestus</i> Tomšík, 1947	B	M	
<i>nigriclavus</i> Marshall, 1868	B		
<i>nigripes</i> Thomson, 1858	B	M	
<i>simplex</i> Tomšík, 1947	B		
<i>thomsoni</i> Kieffer, 1911	B	M	
<b>Idiotypa</b> Foerster, 1856			
<i>marii</i> Gregor, 1939			M
<i>rufiventris</i> (Thomson, 1858)			B
<b>Entomacis</b> Foerster, 1856			
<i>biguttata</i> (Kieffer, 1911)	B		
<i>excisa</i> (Kieffer, 1911)	B		
<i>subtruncata</i> Kieffer, 1909			B
<b>Monelata</b> Foerster, 1856			
<i>atricornis</i> (Kieffer, 1911)	B		
<i>cincta</i> Haliday, 1858			M
<i>parvula</i> (Nees, 1834)			S
<b>Basalys</b> Westwood, 1833			
<i>amphoralis</i> (Tomšík, 1948)			M
<i>bispinosa</i> (Tomšík, 1948)			B M

<i>collaris</i> Kieffer, 1911	B	<i>gregori</i> Tomšík, 1946	M
<i>crassiclava</i> (Kieffer, 1908)	B M	<i>lapponicus</i> (Thomson, 1858)	B
<i>exigua</i> (Marshall, 1868)	B M	<i>ornatus</i> (Tomšík, 1946)	M S
<i>fulvicornis</i> (Kieffer, 1911)	B M	<i>parvulus</i> (Kieffer, 1911)	B M
<i>inaequalis</i> (Kieffer, 1911)	B M	<i>rufipes</i> (Thomson, 1858)	M
<i>pedestris</i> (Kieffer, 1907)	B M	<i>rufitarsis</i> (Kieffer, 1911)	B M
<i>plannifrons</i> (Tomšík, 1948)	M	<i>tenuisulcus</i> (Kieffer, 1911)	B M
<i>rufocincta</i> (Kieffer, 1911)	B M	<i>transsylvanicus</i> Szabó, 1977	B
<i>suesica</i> (Kieffer, 1911)	B M	<b>Platymischus</b> Westwood, 1832	
<i>tritoma</i> (Thomson, 1858)	B M	<i>dilatatus</i> Westwood, 1832	B
<b>Diapria</b> Latreille, 1796		<b>Solenopsis</b> Wasmann, 1899	
<i>conica</i> (Fabricius, 1775)	B M S	<i>imitatrix</i> Wasmann, 1899	B S
<b>Trichopria</b> Ashmead, 1893		<b>Acidopria</b> Kieffer, 1913	
<i>atricornis</i> Kieffer, 1911	B	<i>amphoralis</i> Tomšík, 1949	M
<i>ciliaris</i> Kieffer, 1911	B	<i>bispinosa</i> Tomšík, 1949	B M
<i>longiclava</i> Masi, 1931	B		
<i>oogaster</i> (Thomson, 1858)	M	<b>SCELIONIDAE</b>	
<i>rotundata</i> (Kieffer, 1911)	M		
<i>subimpressa</i> (Kieffer, 1911)	M	<b>Tiphodytes</b> Bradley, 1902	
<i>tetratoma</i> Kieffer, 1911	S	<i>gerriphagus</i> (Marshall, 1900)	B
<b>Viennopria</b> Jansson, 1953		<b>Trissolcus</b> Ashmead, 1893	
<i>priesneri</i> Jansson, 1953	B	<i>Microtelenomus</i> auct. nec. Dodd, 1913	
<b>Tetramopria</b> Wasmann, 1899		<i>artus</i> (Kozlov, 1963)	S
<i>aurocincta</i> Wasmann, 1899	B M S	<i>latus</i> (Kozlov, 1963)	M
<i>donisthorpei</i> Kieffer, 1911	B	<i>rufiventris</i> (Mayr, 1908)	B M S
<b>Oxypria</b> Kieffer, 1908		<i>semistriatus</i> (Nees, 1834)	M S
<i>bouceki</i> Masner, 1959	S	<b>Opisthacantha</b> Asmead, 1893	
<b>Aneuropria</b> Kieffer, 1911		<i>Protrimorus</i> Kieffer, 1908	
<i>foersteri</i> (Kieffer, 1910)	B M S	<i>aethra</i> (Walker, 1836)	?
<b>Labolips</b> Haliday, 1857		<b>Telenomus</b> Haliday, 1833	
<i>innupta</i> Haliday, 1857]	B M S	<i>Dissolcus</i> Ashmead, 1893	
<b>Psilus</b> Panzer, 1801		<i>brachialis</i> Haliday, 1833	S
<i>acutangulus</i> (Jansson, 1942)	M	<i>fodori</i> Szelényi, 1941	S
<i>alticeps</i> (Kieffer, 1911)	B	<b>Platytelesnomus</b> Dodd, 1914	
<i>bispinosus</i> (Kieffer, 1911)	B M	<i>danubialis</i> Szelényi, 1939	B S
<i>clypeatus</i> (Thomson, 1858	B M	<b>Trimorus</b> Foerster, 1856	
<i>cornutus</i> Panzer, 1801	B M	<i>Paragrypon</i> Kieffer, 1908	
		<i>algicola</i> (Kieffer, 1910)	B S
		<i>bohemicus</i> Masner, 1962	B
		<i>flavipes</i> (Haliday, 1830)	S



<i>walkeri</i> Kieffer, 1835	B M	<b>Xestonotidea</b> Gahan, 1919
<b>Platystasius</b> Nixon, 1937		<i>pini</i> (Debauche, 1947) B M
<i>transversus</i> (Thomson, 1859)	B	<b>Fahringeria</b> Kieffer, 1921
<i>strangaliophagus</i> Nixon, 1937		<i>synergorum</i> Kieffer, 1921 S
<b>Isostasius</b> Foerster, 1856		<b>Misocyclops</b> Kieffer, 1914
<i>punctiger</i> (Nees, 1834)	B M S	<i>oeclus</i> (Walker, 1835) B
<b>Pseudaphanomerus</b>		<i>ornatus</i> Kieffer, 1914 S
Szelényi, 1941		<b>Urocyclops</b> Maneval, 1936
<i>hyalinatus</i> Szelényi, 1941	S	<i>bettiae</i> Maneval, 1936 B
<b>Allotropa</b> Foerster, 1856		<b>Prosactogaster</b> Kieffer, 1914
<i>jacobsoni</i> Ogloblin, 1939	B	<i>phragmitis</i> (Schrank, 1781) B
<i>mecrida</i> (Walker, 1835)	M	<b>CERAPHRONOIDEA</b>
<b>Fidiobia</b> Ashmead, 1895		<b>CERAPHRONIDAE</b>
<i>hofferi</i> Masner, 1957	M	<b>Neoceraphron</b> Ashmead, 1893
<i>rugosifrons</i> Crawford, 1916	B	<i>clavatellus</i> Szelényi, 1937 B S
<i>tatrae</i> Szelényi, 1941		<i>neglectus</i> Szelényi, 1937 B
<b>Amitus</b> Haldeman, 1850		<b>Ceraphron</b> Jurine, 1807
<i>minervae</i> Silvestri, 1911	B	<i>bispinosa</i> (Nees, 1834) B S
<b>Iphitracelus</b> Haliday, 1835		<i>brevipennis</i> Kieffer, 1907 M
<i>gracilis</i> Masner, 1957	B S	<i>brachypterus</i> Kieffer, 1907
<i>lar</i> Walker, 1835	B M	<i>clavatus</i> (Kieffer, 1906) M
<b>Ectadius</b> Foerster, 1856		<b>Synarsis</b> Foerster, 1878
<i>mamertes</i> (Walker, 1835)	B S	<i>planifrons</i> Kieffer, 1907 M
<b>Leptacis</b> Foerster, 1856		<i>roubali</i> Gregor, 1939 B
<i>fimbriata</i> Kieffer, 1917	B	<b>Elysoceraphron</b> Szelényi, 1936
<i>foersteri</i> (Kieffer, 1914)	B M	<i>hungaricus</i> Szelényi, 1936 S
<b>Sactogaster</b> Foerster, 1856		<b>Aphanogmus</b> Thomson, 1858
<i>tarsa</i> (Walker, 1835)	B	<i>abdominalis</i> (Thomson, 1858) B
<b>Anopedias</b> Foerster, 1856		<i>angustipennis</i> Szelényi, 1940 S
<i>lacustris</i> Kieffer, 1926	S	<i>gracilicornis</i> Foerster, 1861 S
<b>Trichacis</b> Foerster, 1856		<i>nanus</i> (Nees, 1834) B S
<i>pisis</i> (Walker, 1835)	B	<i>polymorphus</i> Hellén et Dessart, 1965 S
<b>Piestopleura</b> Foerster, 1856		MEGASPILIDAE
<i>catillus</i> (Walker, 1835)	B	
<b>Platygaster</b> Latreille, 1809		<b>Trichosteresis</b> Foerster, 1856
<i>manto</i> Walker, 1835	B	<i>glabra</i> (Boheman, 1831) M S
<b>Amblyaspis</b> Foerster, 1856		<i>foersteri</i> Kieffer, 1907
<i>nodicornis</i> (Ratzeburg, 1844)	B	

<b>Dendrocerus</b> Ratzeburg, 1852		<b>Lagynodes</b> Foerster, 1841	
<i>rufipes</i> (Thomson, 1858)	M	<i>acuticornis</i> (Kieffer, 1906)	S
<b>Conostigmus</b> Dahlbom, 1858		<i>crassicornis</i> Kieffer, 1906	S
<i>lentus</i> Kieffer, 1907	M	<i>pallidus</i> (Bohemian, 1831)	M S
		<i>thoracicicus</i> Kieffer, 1906	M S

#### TRIGONALOIDEA

JOSEF ŠEDIVÝ

A single species is known to occur in the whole of Czechoslovakia. The female lays eggs on leaves. If the eggs are ingested by a lepidopterous or a tenthredinoid larva, they produce larvae of the parasitoid, which penetrate into the body cavity of the host where they attack incidental larvae of parasitoids of the orders Hymenoptera or Diptera.

#### TRIGONALOIDEA TRIGONALIDAE

<b>Trigonalis</b> Spinola, 1840	
<i>Pseudogonalos</i> Schulz, 1906	
<i>hahnii</i> Spinola 1840	B M S

ACULEATA

CHRYSIDOIDEA

BETHYLOIDEA

JAROMÍR STREJČEK

In the territory of the ČSSR, Chrysidoidea are represented by 5 families comprising 213 species. In an earlier concept, the superfamily included the families Dryinidae, Embolemidae, Bethylidae and Cleptidae. In the present one, the superfamily Chrysidoidea also contains the family Chrysididae. The first four families have so far been poorly studied in the territory of Czechoslovakia. In greater detail, they were studied by A. A. Oglöblin (1924, 1925) who described several species new for science. His study was continued by A. Hoffer (1935a, 1935b, 1936a, 1936b) who collected an extensive, mostly indetermined collection and described several species. Numerous material was also collected by M. Kocourek. Most of these collections are deposited in the National Museum in Prague. The collections of the entomologists mentioned above and the authors' own collections have been utilised in compiling the list. The collections have been only partly determined and thus the present list will have to be supplemented by a number of new genera and species known to occur in the neighbouring countries or new for science as soon as the collections are determined and thorough faunal investigations are carried out.

The life habits of Bethyloidea are those of parasitoids toids. The species of Dryinidae parasitize nymphae or, rarely, adults of Homoptera. The larvae produce a sac-like case and pupate on plants. The larvae of the subfamily Aphelopinae are endoparasitoids of adult Jassidae. The species of Bethylidae parasitize larvae of certain Coleoptera (Cerambycidae, Cisidae, Anobiidae) and small species of Tortricidae. The species of Cleptidae parasitize fully grown larvae of sawflies ready to pupate. The life habits of the species of Embolemidae are unknown.

In compiling the list, the author adhered to the nomenclature used by Richards (1939).

References: Hoffer, A., 1935a: Nový druh čeledi Bethylidae (Hymenoptera, Vespoidea) z Moravy (Pseudisobrachium Fialai n. sp.) [Eine neue Art der Fam. Bethylidae (Hym. Vespoidea) aus Mähren (Pseudisobrachium Fialai n. sp.)] *Věst. čsl. zool. spol. v Praze*, III: 1—5. — Hoffer, A., 1935b: Nový druh čeledi Bethylidae (Hym. Vespoidea) Epyris Bayeri n. sp. [Une espèce nouvelle de la famille Bethylidae (Hym. — Vespoidea) Epyris Bayeri, n. sp. Brno (Vlastní náklad: 1—4.)] Hoffer, A., 1936a: Nový druh čeledi Dryinidae (Trib. Lestodryini) z Československa. [Eine neue Art der Familie Dryinidae (Trib. Lestodryini) aus der Tschechoslowakei.] *Sbor ent. odd. Nář. Mus. v Praze*, XIV: 164—166. Hoffer, A., 1936b: Rod Mesitius Thoms. v Československé republice. [Genus Mesitius Thoms. in der Tschechoslowakei], *Cas. čsl. spol. ent.* XXXIII: 119—124.

Ogloblin A. A., 1924: Nový druh rodu Prenanteon (Anteoninae, Proctotrupidae, Hym.) z Vysokých Tater. *Jubilejní sbor. čsl. spol. ent.*: 102—106. Ogloblin, A. A., 1925: Popisy nových druhů a poznámky o některých Anteoninech (Hym. Proctotrupidae) Československa. [Description of new species and notes on some Anteoninae (Proctotrupidae) of Czechoslovakia.] *Sbor. ent. odd. Národní Mus. Praha*, II: 117—131. Richards O. W., 1939: The British Bethylidae (s. l.) (Hymenoptera). *Trans. R. Ent. Soc. Lond.* 89, P. 8: 185—344.

BETHYLOIDEA

DRYINIDAE

DRYININAE

GONOTOPODINI

**Gonatopus** Ljungh, 1810

*Digonatopus* Kieffer, 1913

*barbatellus* Richards, 1939 \*S  
*formicarius* Ljungh, 1810 B\*M\*S

*pastoralis* Ogloblin, 1925

*sepsoides* Westwood, 1833 B\*M\*S

*sociabilis* var. *ignotus*  
Ogloblin, 1925

**Dicondylus** Haliday  
in Curtis, 1829—30

*bicolor* (Haliday  
in Curtis, 1828) B M S

*decretorius* Haupt, 1916

*lindbergi* Heikenheimo,  
1957 \*B

*multidens* Ogloblin, 1925 B

**Neogonatopus** Perkins, 1905

*lunatus* (Klug, 1810) B M S

**Pseudogonatopus** Perkins,  
1905

*distinctus* (Kieffer, 1906) B M

*terrestris* Ogloblin, 1925

*separatus* Richards, 1939 \*B\*M\*S

**Platygonatopus** Kieffer, 1906

*pedestris* (Dahman, 1818) \*B

*planiceps* (Kieffer, 1905) \*S

**Tetradontochelys** Richards,  
1939

*ljunghii* (Westwood, 1833) ?

**Agonatopoides** Perkins, 1907

*Plectrogonatopus* Richards,  
1939

*striatus* (Kieffer, 1905) B M S

**Allogonatopus** Haupt, 1938

*albidus* (Kieffer, 1906) \*B\*M\*S  
*distinguendus* (Kieffer,  
1905) B \*S

*luteicornis* (Kieffer, 1905) \*M  
*procerus* Haupt, 1938 \*S

BOCCHINI

**Bocchus** Ashmead, 1893

*slovacus* Strejček, 1964 S

DRYININI

**Mesodryinus** Kieffer, 1906

*niger* (Kieffer, 1904) \*B M

**Dryinus** Latreille, 1804

*Plastodryinus* Kieffer, 1906

*Lestodryinus* Kieffer, 1911

*formicarius* Latreille,  
1804 \*B

*tarracensis* Marshall,  
1818 B M S

*gregori* Hoffer, 1936

**Mercetia** Kieffer, 1906

*baetica* Ceballos, 1927 \*S

ANTEONINI

**Mystrophorus** Foerster, 1856

*formicaeformis* Ruthe,  
1856 B M\*S

**Prenanteon** Kieffer, 1913

*Psilanteon* Kieffer, 1913

*daos* (Walker, 1837) B

*filicornis* Ogloblin, 1925 B

*frontalis* (Dalman, 1818) B\*M\*S

*palustris* Ogloblin, 1924 S

*ruficornis* (Dalman, 1818) B\*M\*S

*pallidinervis* Kieffer, 1905

*dolichocerus* Kieffer, 1905

**Chelogynus** Haliday, 1838

*albidocolle* (Kieffer, 1905) B

<i>albidicollis</i> Kieffer, var.		<i>Polyplanus</i> Nees, 1834
<i>abdominalis</i> Ogloblin, 1925		<i>ruddii</i> Westwood, 1833 B*M S
<i>albidicollis</i> Kieffer, var. <i>incertus</i> Ogloblin, 1925		
<i>angusticollis</i> Ogloblin, 1925	B	
<i>cameroni</i> (Kieffer, 1905)	*B*M*S	
<i>fulviventre</i> (Haliday, in Curtis, 1828)	*B*M*S	
<i>gaullei</i> (Kieffer, 1905)	B*M*S	
<i>infectum</i> (Haliday in Walker, 1837)	*B*M*S	
<i>kiefferi</i> (Chitty, 1908)	*B*M*S	
<i>angusticollis</i> Berland, 1928, non Ogloblin, 1925		
<i>lucidus</i> (Haliday in Curtis, 1828)	*B	
<i>rufoniger</i> Ogloblin, 1925	B	
<i>scapulare</i> (Haliday in Walker, 1837)	B	
<i>longiforceps</i> Kieffer, 1914		
<b>Anteon</b> Jurine, 1807		
<i>Prosanteon</i> Perkins, 1905		
<i>Xenanteon</i> Kieffer, 1913		
<i>brachycerum</i> (Dalman, 1823)	*B	
<i>brevicornis</i> (Dalman, 1818)	B M*S	
<i>flavicone</i> (Dalman, 1823)	*B	
<i>vicus</i> Kieffer, 1905		
<i>jurineanum</i> Latreille, 1809	*M	
<i>sekerae</i> Ogloblin, 1925	*B	
<b>APHELOPINAE</b>		
<b>Aphelopus</b> Dalman, 1823		
<i>heidelbergensis</i> Richards, 1939	*B *S	
<i>holomelas</i> Richards, 1939	*B*M*S	
<i>melaleucus</i> (Dalman, 1818)	B*M*S	
<i>nigriceps</i> Kieffer, 1905	*B*M*S	
<i>serratus</i> Richards, 1939	*B*M	
<b>EMBOLEMIDAE</b>		
<b>Embolemus</b> Westwood, 1833		
<i>Myrmecomorphus</i> Westwood, 1833		
<i>Pedinomma</i> Foerster, 1856		
		<b>BETHYLIDAE</b>
		<b>BETHYLINAE</b>
		<b>BETHYLIINI</b>
		<b>Parasierola</b> Cameron, 1883
		<i>erucarum</i> Szélenyi, 1958 S
		<b>Perisierola</b> Kieffer, 1914
		<i>gallicola</i> (Kieffer, 1905) S
		<b>Goniozus</b> Foerster, 1856
		<i>claripennis</i> (Foerster, 1851) B M S
		<i>mobilis</i> Foerster, 1860 *M*S
		<b>Anoxus</b> Thomson, 1862
		<i>Omalus</i> Kieffer, 1904 p. p.
		<i>coniceps</i> Kieffer, 1904 *B*M*S
		<i>boops</i> Thomson, 1862 *B*M
		<b>Bethylus</b> Latreille, 1802
		<i>Perisemus</i> Foerster, 1856
		<i>Episemus</i> Thomson, 1862
		<i>arcuatus</i> Kieffer, 1905 *B
		<i>cephalotes</i> (Foerster, 1860) B*M*S
		<i>fuscicornis</i> (Jurine, 1807) B*M*S
		<b>Israelius</b> Richards, 1952
		<i>carthami</i> Richards, 1952 *S
		<b>EPYRINAE</b>
		<b>CEPHALONOMINI</b>
		<b>Plastonoxus</b> Kieffer, 1905
		<i>chittendenii</i> (Ashmead, 1893) *B
		<i>incompletus</i> Evans, 1965 *B*M
		<i>westwoodi</i> (Kieffer, 1914) *S
		<b>Cephalonomia</b> Westwood, 1833
		<i>Holopedina</i> Foerster, 1850
		<i>Ateleopterus</i> Ashmead, 1893
		<i>formiciformis</i> Westwood, 1833 *B*M*S
		<i>stephanoderis</i> Betrem, 1961 *B
		<i>tarsalis</i> (Ashmead, 1893) *B
		<i>quadridentata</i> Duchaussay, 1917 B

<i>strandii</i> Hoffer, 1936			
<i>hypobori</i> Kieffer, 1919	*B		
<i>xambeui</i> Giard, 1874	*B		
<i>waterstoni</i> Gaham, 1931	*B		
<b>Acephalonoma</b> Strejček, 1989			
<i>cisidophaga</i> Strejček, 1989	*B	*S	
EPYRINI			
<b>Ceratepyris</b> Kieffer, 1905			
<i>fuscipennis</i> Kieffer, 1905	*B		
<b>Epyris</b> Westwood, 1832			
<i>bayeri</i> Hoffer, 1935	M S		
<i>bilineata</i> Thomson, 1861	*S		
<i>marshalli</i> Kieffer, 1906	*B		
<i>minor</i> Kieffer, 1906	*B*M*S		
<i>niger</i> Westwood, 1832	*B*M*S		
<b>Holepyris</b> Kieffer, 1905			
<i>Isobrachium</i> Cameron, 1910			
<i>africanus</i> Kieffer, 1904	*S		
<i>svylvanidis</i> (Brèthes, 1913)	*B	*S	
<b>Laelius</b> Ashmead, 1856			
<i>anthrenivorus</i> Tranii, 1909	*B	*S	
<i>bipartitus</i> Kieffer, 1906	*B*M		
<i>foersteri</i> Kieffer, 1914	*B		
SCLERODERMINI			
<b>Glenosema</b> Kieffer, 1905			
<i>nigra</i> Kieffer, 1906	*M*S		
<b>Sclerodermus</b> Klug, 1809			
<i>fonscolombei</i> (Westwood, 1881)	*M		
<i>domesticus</i> Klug, 1809	*B*M		
MESITIINAE			
<b>Mesitius</b> Spinola, 1851			
<i>fuscicornis</i> Kieffer, 1906	M*S		
<i>zavadili</i> Hoffer, 1936			S
<b>Heterocoelia</b> Dahlbom, 1854			
<i>Mesitius</i> Spinola, 1851, 1851, p. p.			
<i>cursor</i> (Kieffer, 1906)	M		
<i>halidayella</i> (Westwood, 1874)	B M		
<i>hungarica</i> (Kieffer, 1906)	B*M*S		
PRISTOCERINAE			
<b>Pristocera</b> Klug, 1808			
<i>Sclerochora</i> Foerster, 1850			
<i>depressa</i> (Fabricius, 1805)	M S		
<b>Parascleroderma</b> Kieffer, 1904			
<i>nigriceps</i> Kieffer, 1904	*S		
<i>rugosulum</i> Kieffer, 1906	*B		
<b>Pseudisobrachium</b> Kieffer, 1905			
<i>subcyaneum</i> (Haliday, 1838)	*B M*S		
<i>fialai</i> Hoffer, 1935			
CLEPTIDAE			
<b>Cleptes</b> Latreille, 1802			
<i>aerosus</i> Foerster, 1853	*B M S		
<i>chevrieri</i> Frey-Gessner, 1887	S		
<i>ignitus</i> (Fabricius, 1793)	*S		
<i>nitidulus</i> (Fabricius, 1793)	B M S		
<i>orientalis</i> Dahlbom, 1854	*S		
<i>scutellaris</i> Mocsáry, 1901	S		
<i>semauratus</i> (Linnaeus, 1761)	B M S		
<i>pallipes</i> Lepeletier, 1806			
<i>diana</i> Mocsáry, 1889			
<i>semicyaneus</i> Tournier, 1879	*S		
<i>splendens</i> (Fabricius, 1793)	*S		

## CHYSIDIDAE

ZDENĚK PÁDR

The distribution and occurrence of species of the family Chrysidiidae in Czechoslovakia are relatively well-known. In the past, the family was studied by a number of authors (A. Kubes, E. Hachler, K. Špaček, J. Snoflák, V. Zavadil, A. Hoffer, M. Kocourek, J. Pospíšil, V. Balthasar, J. Niedl, K. Deneš, V. Tyrner and the present author). The results of their faunal investigations are mostly contained in the *Prodromus hymenopterorum Čechoslovakiae* (Balthasar, 1946) and in the monograph by Balthasar (1954). The collections of the authors mentioned above are deposited in the National Museum in Prague and in the Moravian Museum in Brno. Part of the collections is private. In compiling the list the author adhered to the system used by Balthasar (1954).

The females of Chrysidiidae lay eggs in nests of various species of Sphecoidea, Vespoidea and Apoidea. The larvae are parasitic, feeding on food stored for the host larva, regardless of whether it is nectar and pollen or paralysed arthropods. The time and way of killing the host larva are different in different species of the cuckoo wasp.

In recent years, attempts at nomenclatural changes have occurred, stimulated by the paper of W. Lisenmaier (1959, 1968). The re-evaluations of the species contained in the collections will probably result in supplements, especially in the genera *Holopyga*, *Hedychrum*, *Hedychradium* and *Chrysis*. New discoveries are to be expected particularly in the southern parts of the territory of the ČSSR.

References: Balthasar, V., 1946: *Prodromus Chrysidiidarum Rei Publicae Čechoslovakiae. Acta ent. Mus. Nat. Pragae*, XXIV: 221–261. Balthasar, V., 1954: *Zlatňky — Chrysidoidea. Fauna ČSR*, Vol. 3, Praha: 271 pp. Lisenmaier, W., 1959: Revision der Familie Chrysidiidae (Hymenoptera) mit Nachtrag. *Mitt. Schweiz. Ent. Ges.* XXXII, H. 1: 1–240. — Lisenmaier, W., 1968: Revision der Familie Chrysidiidae (Hymenoptera). Zweiter Nachtrag. *Mitt. Schweiz. Ent. Ges.* XLI: 1–144.

### CHYSIDIDAE

<b>Notozus</b> Foerster, 1853		<i>panzeri</i> (Fabricius, 1804)	B	M	S
<i>ambiguus</i> (Dahlbom, 1854)	*M	<i>panzeri coeruleus</i> (Dahlbom, 1854)		B	M
		<i>soror</i> Mocsary, 1889			S
<i>bidens</i> Foerster, 1853	B M*S	<i>pyrosomus</i> Foerster, 1853		M	S
<i>constrictus</i> Foerster, 1853	B M S	<i>sanzii</i> Gogorza, 1887		M	S

<i>spina</i> (Lepeletier, 1806)	*M S	<i>coriaceum</i> (Dahlbom, 1854)	B M S
<b>Omalus</b> Panzer, 1804		<i>elegantulum</i> Buysson, 1887	B M*S
<i>aeneus</i> (Fabricius, 1787)	B M S	<i>femoratum</i> (Dahlbom, 1854)	B S
<i>auratus</i> (Linnaeus, 1761)	B M S	<i>flavipes</i> (Eversmann, 1857)	S
<i>biacinctus</i> Buysson, 1891	B M S	<i>integrum</i> (Dahlbom, 1831)	B S
<i>bidentulus</i> (Lepeletier, 1806)	B M S	<i>jucundum</i> Mocsary, 1889	S
<i>bogdanovi</i> (Radoszkovsky, 1877)	S	<i>krajniki</i> Balthasar, 1946	S
<i>horvathi</i> Mocsary, 1889	S	<i>monochroum</i> Buysson, 1888	M S
<i>politus</i> Buysson, 1887	S	<i>parkanense</i> Balthasar, 1946	M S
<i>pusillus</i> (Fabricius, 1804)	B M S	<i>purpurascens</i> (Dahlbom, 1854)	B
<i>sculpticollis</i> Abeille, de Perrin, 1878	S	<i>roseum</i> (Rossi, 1790)	B M S
<i>triangulifer</i> Abeille de Perrin, 1877	*B	<i>sculpturatum</i> Abeille de Perrin, 1877	B M S
<i>truncatus</i> Dahlbom, 1831	B M S	<i>scutellare</i> Tournier, 1878	M S
<i>violaceus</i> (Scopoli, 1763)	B M S	<i>valesiense</i> Lisenmaier, 1959	B M S
<b>Holopyga</b> Dahlbom, 1854		<i>zelleri</i> (Dahlbom, 1845)	B M S
<i>amoenula</i> (Dahlbom, 1845)	B M S	<b>Stilbum</b> Spinola, 1806	
<i>austrialis</i> Lisenmaier, 1959	B M S	<i>cyanurum</i> (Foerster, 1771)	M S
<i>chrysonota</i> (Foerster, 1853)	B M S	<b>Euchroeus</b> Latreille, 1809	
<i>fervida</i> (Fabricius, 1781)	B M S	<i>purpuratus</i> (Fabricius, 1787)	M S
<i>ignicollis</i> Dahlbom, 1854	S	<b>Spinolia</b> Dahlbom, 1854	
<i>inflammata</i> (Foerster, 1853)	*B*M*S	<i>dallatorreana</i> Mocsary, 1896	S
<i>jurinei</i> Chevrier, 1862	*S	<i>unicolor</i> Dahlbom, 1831	B M
<i>minuma</i> Lisenmaier, 1959	*S	<b>Pseudochrysis</b> Semenov, 1891	
<i>ovata</i> Dahlbom, 1854	B M S	<i>coeruleiventris</i> Abeille de Perrin, 1878)	S
<b>Hedychrum</b> Latreille, 1806		<i>neglecta</i> (Shuckard, 1837)	B M S
<i>aureicolle niemeläi</i> Lisenmaier, 1959	B M S	<i>uniformis</i> (Dahlbom, 1854)	M
<i>chalybaeum</i> Dahlbom, 1854	B M	<b>Chrysidea</b> Bischoff, 1913	
<i>gerstaeckeri</i> Chevrier, 1869	B M S	<i>pumila</i> (Klug, 1845)	M S
<i>intermedium</i> Dahlbom, 1845	B M S	<b>Spintharis</b> Dahlbom, 1854	
<i>rutilans</i> Dahlbom, 1854	B M S	<i>versicolor</i> (Spinola, 1808)	S
<i>nobile</i> Scopoli, 1763	B M S	<b>Parnopes</b> Latreille, 1796	
<b>Hedychridium</b> Abeille de Perrin, 1878		<i>grandior</i> (Pallas, 1771)	B M S
<i>ardens</i> (Coquebert, 1801)	B M S	<b>Chrysis</b> Linnaeus, 1767	
		<i>analis</i> Spinola, 1808	B M S

<i>angustifrons</i>			
Abeille de Perrin, 1878	M S		
<i>angustula</i> Schenck, 1856	*B *S		
<i>austriaca</i> Fabricius, 1804	B M S		
<i>bicolor</i> Lepeletier, 1806	B M S		
<i>brevitarsis</i> Thomson, 1870	*B *S		
<i>cingulicornis</i> Foerster, 1853	M S		
<i>comparata</i> Lepeletier, 1806	M S		
<i>cuprea</i> Rossi, 1790	M S		
<i>cyanea</i> (Linnaeus, 1761)	B M S		
<i>diacantha</i> Mocsary, 1889	S		
<i>dichroa</i> Dahlbom, 1854	B M S		
<i>equestris</i> Dahlbom, 1845	B		
<i>fasciata</i> Olivier, 1790	B M S		
<i>fivaldskyi</i> Mocsary, 1882	M S		
<i>fulgida</i> Linnaeus, 1761	B M S		
<i>germari</i> Wesmael, 1839	S		
<i>gracillima</i> Foerster, 1853	B M S		
<i>grohmanni</i> Dahlbom, 1854	S		
<i>helleni</i> Linsenmaier, 1959	*B *M *S		
<i>hirsuta</i> Gerstaecker, 1869	B S		
<i>hybrida</i> Lepeletier, 1806	S		
<i>ignita</i> (Linnaeus, 1761)	B M S		
<i>inaequalis</i> Dahlbom, 1845	B M S		
<i>indigotea</i> Dufour et Perris, 1840	B M S		
<i>interjecta</i> Buysson, 1891	S		
<i>rosina</i> Balthasar, 1949			
<i>iris</i> Christ, 1791	B M S		
<i>leachi</i> Shuckard, 1837	B M S		
<i>longula</i> Abeille de Perrin, 1879	*B *M *S		
<i>marginata</i> Mocsary, 1889	M S		
<i>mediata</i> Linsenmaier, 1951	*B *M *S		
<i>phryne</i> Abeille de Perrin, 1878	S		
<i>placida</i> Mocsary, 1879	S		
<i>pseudobrevitarsis</i>			
Linsenmaier, 1951	*B		
<i>pulchella</i> Spinola, 1808	S		
<i>pustulosa</i> Abeille de Perrin, 1878	B M S		
<i>pyrrhina</i> Dahlbom, 1845	?		
<i>ragusai</i> De Stefani, 1888	S		
<i>ramburi</i> Dahlbom, 1854	S		
<i>ruddii</i> Shuckard, 1837	B M S		
<i>rufitarsis</i> Brullé, 1832	S		
<i>rutilans</i> Olivier, 1790	B M S		
<i>rutiliventris</i> Abeille de Perrin, 1879	B M S		
<i>scutellaris</i> Fabricius, 1794	B M S		
<i>sexdentata</i> Christ, 1791	B M S		
<i>simplex</i> Dahlbom, 1854	B M S		
<i>splendidula</i> Rossi, 1790	M S		
<i>subsinuata</i> Marquet, 1879	S		
<i>succincta</i> Linnaeus, 1767	B M S		
<i>sybarita</i> Foerster, 1853	B M S		
<i>trimaculata</i> Foerster, 1853	B M S		
<i>viridula</i> Linnaeus, 1761	B M S		

## SCOLIOIDEA

### ZDENĚK PÁDR

In Czechoslovakia, the superfamily Scolioidea is represented by six families. All have been intensively studied in faunal respect since the nineteen thirties. The collections yielded numerous materials which are deposited in the state collections of the National Museum in Prague, the Moravian Museum in Brno and the Slovak National Museum in Bratislava. The said collections contain materials collected by A. Hoffer, O. Šus-

tera, V. Zavadil, J. Šnoflák, V. Balthasar, etc. Additional numerous material is kept in private collections of M. Kocourek and the present author.

Of the family Tiphidae, 9 species of two genera were found occurring in Czechoslovakia. The faunal findings have been evaluated by A. Hoffer (1938). The larvae parasitize beetle larvae of the family Scarabaeidae. Numerous specimens can be found on daucaceous plants in summer.

The family Methochidae is represented in Czechoslovakia by single species the larvae of which parasitize those of tiger beetles.

Of the family Mutillidae, 21 species are known to occur in Czechoslovakia. Only two of these occur in colder regions whereas the rest are thermophilous. The larvae parasitize those of Eumenidae, Sphecidae and Apidae. In Czechoslovakia, the family has been thoroughly studied as to the occurrence of species. The results of faunal investigations have been published by Hoffer (1938). The nomenclature is based on the paper Lelej (1985).

Of the family Scoliidae, five thermophilous species occur in Czechoslovakia, two of which are known to occur only in the warmest parts of the territory. Scoliid larvae parasitize those of scarabaeiid beetles. The results of faunal investigations have been published by Baťa (1938).

Of the family Sapygidae, 5 species are known to occur in Czechoslovakia. Their larvae kill eggs of larvae Hymenoptera, most of all solitary bees, and feed on pollen stored in the hosts' cells. The results of faunal investigations have been published by Baťa (1938).

The nomenclature used in this check-list is based on several papers on the individual families.

References: Baťa, L., 1938: Sapygidae, Scoliidae. in Prodromus hymenopterorum Čechoslovakiæ, Pars 2. *Sbor. entom. odd. Nář. Musea v Praze*: 167—170. — Hoffer, A., 1938: Tiphidae, Methocidae, Myrmosidae, Mutillidae. in Prodromus hymenopterorum Čechoslovakiæ, Pars 2. *Sbor. entom. odd. Nář. Musea v Praze*: 171—195. — Lelej, A. S., 1985: Osy-nemci fauny SSSR i sopredel'nykh stran, Nauka, Leningrad, 268 pp.

SCOLIOIDEA			
TIPHIIDAE			
<b>Tiphia</b> Fabricius, 1775			
<i>distincta</i> Tournier, 1889	?		
<i>femorata</i> Fabricius, 1775	B M S		
<i>fulvipennis</i> Smith, 1879	S		
<i>minuta</i> Linden, 1827	B M S		
<i>morio</i> Fabricius, 1787	B M S		
<i>ruficornis</i> Klug, 1810	B M S		
<i>polita</i> Costa, 1851			
<i>rugosa</i> Tournier, 1889	?		
<b>Meria</b> Illiger, 1807			
<i>dorsalis</i> (Fabricius, 1804)	S		
		polita (Tournier, 1889)	
		<i>tripunctata</i> (Rossi, 1790)	B M S
		METHOCHIDAE	
		<b>Methocha</b> Latreille, 1805	
		<i>articulata articulata</i>	
		(Latreille, 1792)	B M S
		<i>ichneumonides</i> Latreille, 1805	
		<i>articulata nigrescens</i>	
		Hoffer, 1938	M S
		MUTILLIDAE	
		<b>Myrmosa</b> Latreille, 1796	
		<i>atra</i> Panzer, 1801	B M S

<i>melanocephala</i> (Fabricius, 1793)		<b>Smicromyrme</b> Thomson, 1870	
<b>Krombeinella</b> Pate, 1947		<i>pouzdranensis</i> Hoffer, 1936	M
<i>thoracica</i> (Fabricius, 1793)	S	<i>rufipes</i> (Fabricius, 1787)	B M S
<i>ephippium</i> (Fabricius, 1793)		<i>septentrionalis</i> Hoffer, 1936	M S
<b>Paramyrmosa</b> Saussure, 1880		<i>pusilla septentrionalis</i> Hoffer, 1936	
<i>brunnipes</i> (Lepelletier, 1845)	M S	<i>viduata</i> (Pallas, 1773)	M S
<b>Myrmilla</b> Wesmael, 1851		SCOLIIDAE	
<i>calva calva</i> (Villers, 1789)	M S	<b>Campsoscolia</b> Betrem, 1933	
<i>calva distincta</i> (Lepeletier, 1845)	B M S	<i>quinquecineta</i> (Fabricius, 1793)	S
<i>erythrocephala</i> (Latreille, 1792)	M S	<i>sexmaculata</i> (Fabricius, 1781)	M S
<b>Cystomutilla</b> André, 1896		<i>interrupta</i> (Fabricius, 1781)	
<i>ruficeps</i> (Smith, 1855)	S	<b>Scolia</b> Fabricius, 1775	
<b>Dasylabris</b> Radoszkowski, 1855		<i>hirta</i> (Schrank, 1781)	B M S
<i>maura</i> (Linnaeus, 1758)	M S	<i>maculata</i> (Drury, 1773)	S
<i>regalis</i> (Fabricius, 1793)	M S	<i>flavifrons</i> Fabricius, 1775 <i>s. haemorrhoidalis</i> Eversman, 1849	
<i>italica</i> (Fabricius, 1793)		<i>quadripunctata</i> Fabricius, 1775	B M S
<b>Mutilla</b> Linnaeus, 1758		SAPYGIDAE	
<i>europea</i> Linnaeus, 1758	B M S	<b>Sapygina</b> Costa, 1887	
<i>marginata</i> Baer, 1848	B M S	<i>decemguttata</i> (Jurine, 1807)	B M S
<b>Ronisia</b> Costa, 1858		<b>Sapyga</b> Latreille, 1796	
<i>brutia</i> (Petagna, 1787)	M S	<i>clavicornis</i> (Linnaeus, 1758)	B M S
<b>Psysetopoda</b> Schuster, 1949		<i>quinquepunctata</i> (Fabricius, 1781)	B M S
<i>halensis</i> (Fabricius, 1787)	B M S	<i>similis</i> (Fabricius, 1793)	B M S
<i>montana</i> (Panzer, 1806)		<b>Polochrum</b> Spinola, 1806	
<i>punctata pseudomontana</i> (Hoffer, 1936)		<i>repandum</i> Spinola, 1806	S
<i>moravica</i> (Hoffer, 1936)	M S		
<i>variabilis</i> (Hoffer, 1938)			
<i>punctata</i> (Latreille, 1792)	S		
<i>pusilla</i> (Klug, 1835)	M S		
<i>subcomata</i> (Wesmael, 1851)	M		



## FORMICOIDEA

PETR WERNER

The first sporadic notes on the superfamily Formicoidea from the territory of Czechoslovakia can be found in the classical work of Mayr (1855). Lokay E. sen. (1860) published the first list of the ants of Bohemia. He started myrmecological studies as a part of his research on myrmecophilous beetles. Several other coleopterists contributed to our knowledge of Czechoslovakian ant fauna thanks to their interest in myrmecophilous beetles (E. Lokay, jun., T. Krása, J. Roubal, F. Rambousek). The first list of Moravian Formicoidea was published by Zdobnitzky (1910). The first faunal records from Slovakia were published in the work Mocsáry (1918).

More extensive faunal research of the Czechoslovakian ants began in the nineteen twenties and thirties. The results of this intensive faunal research was summarized, up to October, 1939, in the work Záleský (1939). Kratochvíl (1940) published a supplement to this work. With the exception of the above-mentioned authors, the following entomologists participated significantly in the research of the Czechoslovakian ant fauna: Š. Soudek, V. Šilhavý, J. Sadil, V. J. A. Novák, K. Samšiňák, A. Beláková, M. Labuda and P. Bezdečka.

This check-list was prepared on the basis of the collections of the National Museum in Prague, Moravian Museum in Brno, Slovakian Nat. Museum in Bratislava, other Czechoslovak museums and certain private collections.

The ant fauna of Moravia, central Bohemia and southwestern Slovakia are relatively best known. The ant fauna of central and eastern Slovakia and of all mountain regions have been poorly investigated.

References: Kratochvíl, J., 1940: Doplněk nalezišť k Záleského Prodromu mravenců. *Sbor. ent. odděl. Nář. Mus. v Praze* 18: 241—249. — Lokay, E., 1860: Popsání hlavních druhů mravenců v Čechách žijících, s ohledem na hosti u nás dosud v mraveništích nalezené. *Živa* 8: 238—253. — Mayr, G., 1855: Formicina austriaca. Verh. Zool.-bot. Ver., Wien 5: 273—478. — Mocsáry, A., 1918: Ordo Hymenoptera. Fauna Regni Hungariae III., Budapest — Záleský, M., 1939: Formicoidea. Prodromus našeho hmyzu blanokřídlého III. *Sbor. ent. odděl. Nář. Mus. v Praze* 17: 192—240. — Zdobnitzky, W., 1910: Beitrag zur Ameisenfauna Mährens. Zeitschr. mährischen Landesmuseums 10: 1—13.

FORMICOIDEA			
FORMICIDAE			
PONERINAE			
<b>Proceratium</b> Roger, 1863			
<i>Sysphincta</i> Roger, 1863			
<i>melatum</i> (Roger, 1860)	M		
<i>fialai</i> (Kratochvíl, 1944)			
<b>Hypoponera</b> Santschi, 1938			
<i>punctatissima</i> (Roger, 1859)	B M		
<b>Ponera</b> Latreille, 1802			
<i>coarctata</i> (Latreille, 1802)	B M S		
MYRMICINAE			
<b>Manica</b> Jurine, 1807			
<i>rubida</i> (Latreille, 1802)	B M S		
<b>Myrmica</b> Latreille, 1804			
<i>deplanata</i> Ruzsky, 1905	B M S		
<i>gallieni</i> Bondroit, 1919	B*M*S		
<i>bergi</i> (sensu Sadil, 1952)			
<i>hirsuta</i> Elmes, 1978	*B *S		
<i>lobicornis</i> Nylander, 1846	B M S		
<i>rubra</i> (Linnaeus, 1758)	B M S		
<i>laevinodis</i> Nylander, 1846			
<i>ruginodis</i> Nylander, 1846	B M S		
<i>rugulosa</i> Nylander, 1848	B M S		
<i>sabuleti</i> Meinert, 1860	B M S		
<i>salina</i> Ruzsky, 1905	B*M*S		
<i>slovaca</i> Sadil, 1952			
<i>scabrinodis</i> Nylander, 1846	B M S		
<i>schencki</i> Emery, 1894	B M S		
<i>specioides</i> Bondroit, 1918	B M S		
<i>balcanica</i> Sadil, 1952			
<i>sulcinodis</i> Nylander, 1846	B M S		
<i>vandeli</i> Bondroit, 1920	*B *S		
<b>Sifolinia</b> Emery, 1907			
<i>karavajevi</i> (Arnoldi, 1930)			
1930)	B M		
<b>Aphaenogaster</b> , Mayr, 1853			
<b>Attomyrma</b> Emery, 1915			
<b>subterranea</b> (Latreille, 1798)	B M S		
<b>Messor</b> Forel, 1890			
<b>meridionalis</b> (André, 1882)	?		
<b>muticus</b> (Nylander, 1849)	M S		
<b>Monomorium</b> Mayr, 1855			
<i>pharaonis</i> (Linnaeus, 1758) introd.	B M S		
<b>Diplorhoptrum</b> Mayr, 1855			
<i>fugax</i> (Latreille, 1798)	B M S		
<b>Anergates</b> Forel, 1874			
<i>atratulus</i> (Schenck, 1852)	B M S		
<b>Myrmecina</b> Curtis, 1829			
<i>graminicola</i> (Latreille, 1802)	B M S		
<b>Leptothorax</b> Mayr, 1855			
<b>Leptothorax</b> s. str.			
<i>acervorum</i> (Fabricius, 1793)	B M S		
<i>gredleri</i> Mayr, 1855	B M S		
<i>muscorum</i> (Nylander, 1846)	B M S		
<b>Myrafant</b> Smith, 1950			
<i>affinis</i> Mayr, 1855	B M S		
<i>clypeatus</i> (Mayr, 1853)	M*S		
<i>corticalis</i> (Schenck, 1852)	B M S		
<i>interruptus</i> (Schenck, 1852)	B M*S		
<i>lichtensteinii</i> Bondroit, 1918	?		
<i>luteus</i> Forel, 1874	?		
<i>nigriceps</i> Mayr, 1855	B M*S		
<i>nylanderi</i> (Foerster, 1850)	B M S		
<i>parvulus</i> (Schenck, 1852)	B M S		
<i>sordidulus</i> Müller, 1923	B M*S		
<i>tuberum</i> (Fabricius, 1775)	B M S		
<i>unifasciatus</i> (Latreille, 1798)	B M S		
<b>Epimyrma</b> Emery, 1915			
<i>ravouxi</i> (André, 1896)	*B S		
<i>zaleskyi</i> Sadil, 1954			
<b>Formicoxenus</b> Mayr, 1855			
<i>nitidulus</i> (Nylander, 1846)	B M S		
<b>Harpagoxenus</b> Forel, 1893			
<i>sublaevis</i> (Nylander, 1848)	B M*S		

<b>Stenamma</b> Westwood, 1840		<b>Aporomyrmex</b> Faber, 1969	
<i>westwoodi</i> Westwood, 1840	B M S	<i>ampeloni</i> Faber, 1969	*S
<b>Tetramorium</b> Mayr, 1855		<b>Camponotus</b> Mayr, 1861	
<i>caespitum</i> (Linnaeus, 1756)	B M S	<b>Camponotus</b> s. str.	
<i>ferox</i> Ruzsky, 1903	M S	<i>herculeanus</i> (Linnaeus, 1758)	B M S
<i>impurum</i> (Foerster, 1850)	B M S	<i>ligniperda</i> (Latreille, 1802)	B M S
<i>moravicum</i> Kratochvíl, 1944	B M S	<i>vagus</i> (Scopoli, 1763)	M S
<i>semilaeve</i> André, 1883	*S	<b>Colobopsis</b> Mayr, 1861	
<b>Strongylognathus</b> Mayr, 1853		<i>truncatus</i> (Spinola, 1808)	B M S
<i>bulgaricus</i> Viehmeyer, 1922	M	<b>Myrmentoma</b> Forel, 1912	
<i>kratochvili</i> Šilhavý, 1937		<i>dalmaticus</i> (Nylander, 1848)	S
<i>testaceus</i> (Schenck, 1852)	B M S	<i>fallax</i> (Nylander, 1856)	B M S
DOLICHODERINAE		<i>lateralis</i> (Olivier, 1791)	?M?S
<b>Dolichoderus</b> Lund, 1831		<i>piceus</i> (Leach, 1825)	M S
<b>Hypoclinea</b> Mayr, 1855		<b>Tanaemyrmex</b> Ashmead, 1906	
<i>quadripunctatus</i> (Linnaeus, 1771)	B M S	<i>aethiops</i> (Latreille, 1798)	B M S
<b>Liometopum</b> Mayr, 1861		<i>sylvaticus</i> (Olivier, 1791)	?
<i>microcephalum</i> (Panzer, 1798)	M S	<b>Cataglyphis</b> Foerster, 1850	
<b>Tapinoma</b> Foerster, 1850		<b>Monocombus</b> Mayr, 1855	
<i>ambiguum</i> Emery, 1925	B M*S	<i>aenescens</i> (Nylander, 1849)	S
<i>erraticum</i> (Latreille, 1798)	B M S	<i>nodus</i> (Brullé, 1832)	S
<b>Bothriomyrmex</b> Emery, 1869		<b>Lasius</b> Fabricius, 1804	
<i>corsicus mohelensis</i>		<b>Austrolasius</b> Faber, 1967	
Novák, 1944	M	<i>reginae</i> Faber, 1967	B S
<i>gibbus</i> Soudek, 1924	M S	<b>Cautolasius</b> Wilson, 1955	
FORMICINAE		<i>flavus</i> (Fabricius, 1781)	B M S
<b>Plagiolepis</b> Mayr, 1861		<i>myops</i> Forel, 1894	B M S
<b>Plagiolepis</b> s. str.		<b>Chthonolasius</b> Ruzsky, 1913	
<i>pygmaea</i> (Latreille, 1798)	B M S	<i>affinis</i> (Schenck, 1952)	B M S
<i>vindobonensis</i> Lomnicki, 1925	M S	<i>bicornis</i> (Foerster, 1850)	*B M S
<b>Paraplagiolepis</b> Faber, 1969		<i>distinguendus</i> (Emery, 1916)	*M*S
<i>xene</i> Stärcke, 1936	*M*S	<i>jensi</i> Seifert, 1982	*B*M*S
<b>Prenolepis</b> Mayr, 1861		<i>meridionalis</i> (Bondroit, 1919)	*B
<i>nitens</i> Mayr, 1852)	S	<i>mixtus</i> (Nylander, 1846)	B M S
		<i>rabaudi</i> (Bondroit, 1917)	*M*S
		<i>umbratus</i> (Nylander, 1846)	B M S
		<b>Dendrolasius</b> Ruzsky, 1913	
		<i>fuliginosus</i> (Latreille, 1798)	B M S
		<b>Lasius</b> s. str.	
		<i>alienus</i> (Foerster, 1850)	B M S

<i>brunneus</i> (Latreille, 1798)	B M S	<i>truncorum</i> Fabricius, 1804	B M S
<i>emarginatus</i> (Olivier, 1791)	B M S	<b>Raptiformica</b> Forel, 1913	
<i>niger</i> (Linnaeus, 1758)	B M S	<i>sanguinea</i> Latreille, 1798	B M S
<b>Formica</b> Linnaeus, 1758		<b>Serviformica</b> Forel, 1913	
<b>Coptoformica</b> Müller, 1923		<i>cinerea</i> Mayr, 1853	B M S
<i>exsecta</i> Nylander, 1846	B M S	<i>cunicularia</i> Latreille, 1798	B M S
<i>pressilabris</i> Nylander, 1846	B M*S	<i>fusca</i> Linnaeus, 1758	B M S
<b>Formica</b> s. str.		<i>gagates</i> Latreille, 1798	M S
<i>aquilonia</i> Yarrow, 1955	*B	<i>lemani</i> Bondroit, 1917	B M S
<i>lugubris</i> Zetterstedt, 1840	B M	<i>picea</i> Nylander, 1846	B M S
<i>nigricans</i> Emery, 1909	B M*S	<i>rufibarbis</i> Fabricius, 1793	B M S
<i>polyctena</i> Foerster, 1850	B M S	<b>Polyergus</b> Latreille, 1805	
<i>pratensis</i> Retzius, 1783	B M S	<i>rufescens</i> (Latreille, 1798)	B M S
<i>rufa</i> Linnaeus, 1758	B M S		

## VESPOIDEA

ZDENĚK PÁDR

The distribution of species of the superfamily Vespoidea in Czechoslovakia is fairly well-known. The hitherto findings have been made all over the territory. Specimens evidencing them are deposited in the National Museum in Prague, the Moravian Museum in Brno, the Slovak National Museum in Bratislava, in several regional and district museums as well as in numerous private collections. O. Šustera, J. Šnoflák, V. Zavadil, A. Hoffer, V. Balthasar, Z. Bouček, J. Niedl, K. Deneš, M. Kocourek and the present author deserved well of the faunal investigations of Vespoidea.

In Czechoslovakia, the Vespoidea are represented by three families among which the rare family Masaridae is the oldest evolutionarily. The infrequent species of this family are solitary. Their larvae feed on pollen and nectar, or on paralysed arthropods which the female deposits in earthen cells built on plants. The species of the most numerous family, Eumenidae are solitary. The cells in which food for their larvae is stored are built in the soil or in cavities in dead wood. The life habits of the larvae are similar to those of the foregoing family. The Vespidae comprises social wasps living in colonies and building their nests above or below the soil surface. All species are predominantly carnivorous. The family also comprises social parasitoids.

New findings may only be expected in some parts or localities in Czechoslovakia.

The genera and species are arranged according to Blüthgen (1961) and Gusenleitner (1981).

References: Blüthgen, P., 1961: Die Faltenwespen Mitteleuropas (Hymenoptera, Diptoptera). *Abhandlungen der Deutsch. Akad. d. Wissenschaft.*, Nr. 2: 248 p. — Gusenleitner, J., 1981: Catalogus Faunae Austriae, Teil XVI. K. Ü. — Fam. Vespoidea. Verlag d. Österreich. Akad. d. Wissenschaften, Wien, 13 p.

### VESPOIDEA MASARIDAE

**Celonites** Latreille, 1802  
*abbreviatus abbreviatus*  
Villers, 1789

### EUMENIDAE

### ZETHINAE

**Discoelius** Latreille, 1809

*dufourii* Lepeletier, 1831 B M S  
*priesneri* Mader, 1936  
**S** *zonalis* (Panzer, 1801) B M S

### EUMENINAE

<b>Paragymnomerus</b> Blüthgen, 1938		<i>nugdunensis</i> (Saussure, 1856)	B M S
<i>spicornis</i> (Spinola, 1808)	M S	<i>timidus</i> (Saussure, 1856)	B M S
<b>Tropidodynerus</b> Blüthgen, 1939		<b>Leptochilus</b> Saussure, 1852	
<i>interruptus interruptus</i> (Brullé, 1832)	S	<i>Lionotulus</i> Blüthgen, 1938	
<b>Odynerus</b> Latreille, 1802		<i>alpestris</i> (Saussure, 1856)	B M S
<i>Oplomerus</i> Westwood, 1840		<i>tarsatus</i> (Saussure, 1856)	S
<b>Odynerus</b> s. str.		<i>Neoleptochilus</i> Blüthgen, 1961	
<i>melanocephalus</i>		<i>regulus</i> (Saussure, 1856)	M S
<i>melanocephalus</i> (Gmelin in Linnaeus, 1790)	B M S	<b>Jucancistrocerus</b> Blüthgen, 1938	
<i>poecilus</i> Saussure, 1856	M S	<i>jucundus</i> (Mocsáry, 1883)	S
<i>spinipes spinipes</i> (Linnaeus 1758)	B M S	<b>Stenodynerus</b> Saussure, 1863	
<i>Spinicoxa</i> Blüthgen, 1938		<i>Nannodynerus</i> Blüthgen, 1938	
<i>reniformis</i> (Gmelin in Linnaeus, 1790)	B M S	<i>blüthgeni</i> van der Wecht, 1971	B M S
<b>Gymnomerus</b> Blüthgen, 1938		<i>dentisquama</i> (Thomson, 1870)	
<i>laevipes laevipes</i> (Shuckard, 1837)	B M S	<b>chevrieranus</b> (Saussure, 1856)	B M S
<b>Hemipterochilus</b> Ferton, 1909		<i>laticinctus</i> (Schulthess, 1897)	?S
<i>bembeciformis</i>		<i>orenburgensis</i> (André, 1884)	B M S
<i>terricola</i> (Mocsáry, 1883)	S	<i>orbitalis</i> (Thomson, 1874)	
<b>Pterocheilus</b> Klug, 1805		<i>punctifrons</i> (Thomson, 1874)	?M
<i>phaleratus phaleratus</i> (Panzer, 1797)	B M S	<i>steckianus</i> (Schulthess, 1897)	B M S
<b>Alastor</b> Lepeletier, 1841		<i>teutonicus</i> Blüthgen, 1937	
<i>biegelebeni pannonicus</i> Blüthgen, 1956	M S	<i>xanthomelas</i> (Herrich- -Schaeffer, 1939)	B M S
<b>Alastorynerus</b> Blüthgen, 1938		<b>Parodontodynerus</b>	
<i>alastoroides ludendorffi</i> (Dusmet, 1917)	S	Blüthgen, 1938	
<b>Pseudomicrodynerus</b>		<i>ephippium</i> (Klug, 1817)	S
Blüthgen, 1938		<b>Antepipona</b> Saussure, 1855	
<i>parvulus</i> (Herrich- -Schaeffer, 1838)	B M S	<i>Odontodynerus</i> Blüthgen, 1938	
<b>Microdynerus</b> Thomson, 1874		<i>deflenda</i> (Saunders, 1853)	B M S
<i>exilis</i> (Herrich- -Schaeffer, 1839)	B M S	<i>orbitalis</i> (Herrich- -Schaeffer, 1839)	B M S
<i>longicollis</i> Morawitz, 1895	S	<b>Allodynerus</b> Blüthgen, 1938	
		<i>delphinalis</i> (Giraud, 1866)	B M S
		<i>floricola</i> (Saussure, 1853)	S
		<i>rossii</i> (Lepeletier, 1841)	M S
		<b>Pseudepipona</b> Dalla Torre, 1894	

<i>augusta</i> (Morawitz, 1867)	S	<i>angustatus</i> (Zetterstedt, 1838)	B	S
<i>herrichii herrichii</i> (Saussure, 1856)	M S	<i>alternans</i> (Zetterstedt, 1838)		
<i>variegata</i> (Herrick-Schaeffer, 1839)		<i>bifasciatus</i> (Linnaeus, 1761)	B	M S
<i>lativentris</i> (Saussure, 1856)	M	<i>connexus</i> (Curtis, 1826)	B	M S
<b>Euodynerus</b> Dalla Torre, 1904		<i>crassicornis crassicornis</i> (Panzer, 1798)	B	M S
<b>Euodynerus</b> s. str.		<i>debilitatus</i> (Saussure, 1856)	B	M S
<i>dantici dantici</i> (Rossi, 1790)	B M S	<i>gracilis gracilis</i> (Brullé, 1832)	B	M S
<i>egregius</i> (Herrick-Schaeffer, 1939)	S	<i>elegans</i> (Wesmael, 1833)		
<b>Paronodynerus</b> Blüthgen, 1938		<i>murarius</i> (Linnaeus, 1758)	B	M S
<i>notatus notatus</i> (Jurine, 1807)	B M S	<i>mutinensis</i> (Baldini, 1894)	B	M S
<i>quadrifasciatus</i>		<i>sinuatus</i> (Fabricius, 1793)		
<i>quadrifasciatus</i> (Fabricius, 1793)	B M S	<b>Eumenes</b> Latreille, 1802		
<b>Ancistrocerus</b> Wesmael, 1836		<i>coarctatus coarctatus</i> (Linnaeus, 1758)	B	M S
<i>antilope</i> (Panzer, 1789)	B M S	<i>coronatus coronatus</i> (Panzer, 1799)	B	
<i>auctus auctus</i> (Fabricius, 1793)	M S	<i>lunulatus lunulatus</i> (Fabricius, 1804)	M	
<i>renimacula</i> (Lepeletier, 1841)		<i>mediterraneus</i>		
<b>claripennis claripennis</b> Thomson, 1874	B M S	<i>mediteraneus</i> Kriechbaumer, 1879	M	S
<i>quadratus</i> (Panzer, 1799)		<i>papillarius papillarius</i> (Christ, 1791)	M	S
<i>dusmetiulus</i> (Strand, 1914)	B M S	<i>pedunculatus pedunculatus</i> (Panzer, 1799)	M	S
<i>gazella</i> (Panzer, 1789)	B M S	<i>pomiformis pomiformis</i> (Fabricius, 1781)	B	M S
<i>ichneumonideus</i> (Ratzeburg, 1844)	B M S	<i>sareptanus insolitus</i> Müller, 1923	M	S
<i>nigricornis</i> (Curtis, 1826)	B M S	<i>subpomiformis</i> Blüthgen, 1938	B	M S
<i>callosus</i> (Thomson, 1870)		<b>Katamenes</b> Meado Waldo, 1910		
<i>oviventris oviventris</i> Westmael, 1836	B M S	<i>arbustorum arbustorum</i> (Panzer, 1799)	B	M S
<i>parietinus</i> (Linnaeus, 1761)	B M S	VESPIDAE POLISTINAE		
<i>parietum</i> (Linnaeus, 1758)	B M S	<b>Polistes</b> Latreille, 1802		
<i>scoticus scoticus</i> (Curtis, 1826)	B M	<i>biglumis bimaculatus</i> (Geoffroy in Fourcroy, 1758)	B	M S
<i>trifasciatus trifasciatus</i> (Müller, 1776)	B M S			
<b>Symmorphus</b> Wesmael, 1836				
<b>Symmorphus</b> s. str.				

<i>foederatus</i> Kohl, 1898	S	<i>paravespula</i> Blüthgen, 1938
<i>gallicus gallicus</i> (Linnaeus, 1767)	B M S	<i>allovespula</i> Blüthgen, 1943
<i>nimpha</i> (Christ, 1791)	B M S	<i>austriaca</i> (Panzer, 1799) B M S
<b>Sulcopolistes</b> Blüthgen, 1938		<i>germanica germanica</i>
<i>atrimandibularis</i> (Zimmermann, 1930)	M S	(Fabricius, 1793) B M S
<i>sulcifer</i> (Zimmermann, 1930)	S	<i>rufa rufa</i> (Linnaeus, 1758) B M S
		<i>vulgaris</i> (Linnaeus, 1758) B M S
		<b>Dolichovespula</b> Rohwer, 1916
		<i>pseudovespula</i> Bischoff, 1931
		<i>boreovespula</i> Blüthgen, 1943
		<i>metavespula</i> Blüthgen, 1943
VESPINAE		
<b>Vespa</b> Linnaeus, 1758		<i>adulterina</i> (Buysson, 1905) B M
<i>crabro crabro</i> Linnaeus, 1758	B M S	<i>ingrica</i> (Birula, 1931) B
<i>crabro germana</i> Christ, 1791	B M S	<i>omissa</i> Bischoff, 1931
<b>Vespa</b> Thomson, 1869		<i>media</i> (Retzius, 1783) B M S
<b>Vespula</b> s. str. <i>pseudovespula</i> Smiedeknecht, 1881		<i>norwegica norwegica</i> (Fabricius, 1781) B M S
		<i>saxonica</i> (Fabricius, 1793) B M S
		<i>sylvestris sylvestris</i> (Scopoli, 1763) B M S

## POMPILOIDEA

JOSEF ŠEDIVÝ

At the end of the nineteenth and at the beginning of the twentieth centuries, J. Slavíček and P. A. Kubes were engaged in faunal studies of the superfamily Pompiloidea in the territory of present Czechoslovakia. Later, the taxonomy and faunistics of this group were studied by O. Šusterá (1938) who published the first synoptic list of species ascertained in Czechoslovakia. Numerous material from the whole territory was collected by V. Zavadil, A. Hoffer, M. Kocourek, J. Šnoflák, Z. Pádr and other entomologists.

Wolf, H. (1971) published the second synoptic list of Pompiloidea in which he evaluated the hitherto findings in the territory of the ČSSR. The list comprises species contained in the collections of the four most important museums in Prague, Brno, Opava and Bratislava. Besides, he utilised materials collected by 80 entomologists in the territory of the ČSSR. Part of these materials is kept in private collections.

The Pompiloidea is a predatory group of insects. Their larvae develop as euparasites of other arthropods, paralysed by the females sting. The paralysed prey is then provided with an egg and buried in loose soil by the female.

So far, Czechoslovakian Pompiloidea have been insufficiently studied in the northern parts of Bohemia, Moravia and Slovakia as well as in the Bohemian-Moravian Highland.

The present list has been compiled exclusively from the data presented in Wolf (1971).

References: Šusterá, O., 1938: Čeled Psammocharidae (olim Pompilidae — Hrabalky) — *Sbor. Ent. odd. Nář. mus. v Praze*, 16: 196—223. In *Prodromus Hymenopterorum Čecho-Slovakiæ*, 2. Wolf, H. (1971): *Prodromus der Hymenopteren der Tschechoslowakei*. Pars 10. Pompiloidea. — *Acta faun. ent. Mus. nat. Pragae*, 14: 76 p.

POMPILOIDEA		
POMPILIDAE		
PEPSINAE		
PEPSINI		
<b>Cryptocheilus</b> Panzer, 1806		
<i>decemguttatus</i>	<i>egregius egregius</i> (Lepeletier, 1845)	S
<i>decemguttatus</i>	<i>fabricii fabricii</i> (Van der Linden, 1827)	B M S
(Jurine, 1807)	<i>freygessneri</i> (Kohl, 1883)	M S
	<i>notatus notatus</i> (Rossi, 1792)	S

<i>notatus affinis</i> (Van der Linden, 1827)	B M S	
<i>richardsi</i> Móczár, 1953	S	
<i>versicolor versicolor</i> (Scopoli, 1763)	B M S	
<b>Priocnemis</b> Schioedte, 1837		
<i>Anapriocnemis</i> Haupt, 1959		
<i>cordivalvata</i> Haupt, 1926	B M S	
<i>coriacea coriacea</i> Dahlbom, 1843	B M S	
<i>enslini enslini</i> Haupt, 1926	M S	
<i>exaltata exaltata</i> (Fabricius, 1776)	B M S	
<i>fastigiata fastigiata</i> Haupt, 1934	M S	
<i>femoralis femoralis</i> (Dahlbom, 1829)	B M	
<i>jennica</i> Haupt, 1926	B M S	
<i>gracilis gracilis</i> Haupt, 1926	B M S	
<i>gracilis huesingi</i> Wolf, 1960	B M S	
<i>hankoi</i> Mocsár, 1944	B M S	
<i>melanosoma</i> Kohl, 1880	M S	
<i>mesobrometi</i> Wolf, 1961	B M S	
<i>mimula mimula</i> Wesmael, 1851	B M S	
<i>minor</i> (Zetterstedt, 1838)	B M S	
<i>minuta minuta</i> (Van der Linden, 1827)	S	
<i>minuta italica</i> Haupt, 1934	S	
<i>obtusiventris</i> Schioedte, 1837	B M S	
<i>parvula parvula</i> Dahlbom, 1845	B M S	
<i>perturbator perturbator</i> (Harris, 1780)	B M S	
<i>pilichi</i> Priesner, 1960	M	
<i>pogoniooides</i> Costa, 1838	B	
<i>propinqua</i> Lepeletier, 1847	B	
<i>pusilla pusilla</i> Schioedte, 1837	B M S	
<i>schioedtei schioedtei</i> Haupt, 1926	B M S	
<i>sulci sulci</i> Balthasar, 1943	M S	
<i>susterai</i> Haupt, 1926	B M	
<b>Calicurgus</b> Lepeletier, 1845		
<i>hyalinatus hyalinatus</i> (Fabricius, 1793)	B M S	
<i>hyalinatus gyllenhali</i> (Dahlbom, 1843)	M S	
<b>Dipogon</b> Fox, 1897		
<i>Deuteragenia Šustera</i> , 1913		
<i>bifasciatum</i> (Geoffroy, 1785)	M S	
<i>hircanum</i> (Fabricius, 1798)	B M S	
<i>nitidum</i> (Haupt, 1926)	B M S	
<i>variegatum variegatum</i> (Linnaeus, 1758)	M S	
<b>Poecilagenia</b> Haupt, 1926		
<i>rubricans</i> (Lepeletier, 1845)	B M	
AUPLOPODINI		
<b>Auplopus</b> Spinola, 1841		
<i>albifrons albifrons</i> (Dalman, 1823)	B M	
<i>carbonarius</i> (Scopoli, 1763)	B M S	
<i>rectus rectus</i> (Haupt, 1926)	B M S	
POMPILINAE		
APORINI		
<b>Aporus</b> Spinola, 1808		
<i>femoralis</i> Van der Linden, 1827	B M S	
<i>fulviventris polux</i> (Kohl, 1888)	M	
<i>tibialis</i> (Tournier, 1890)	B M	
PSAMMODERINI		
<b>Eoferreola</b> Arnold, 1935		
<i>manticata</i> (Pallas, 1771)	S	
<i>rhombica</i> (Christ, 1791)	B M S	
<b>Ferreola</b> Lepeletier, 1844		
<i>diffinis</i> (Lepeletier, 1845)	S	
HOMONOTINI		
<b>Homonotus</b> Dahlbom, 1845		

<i>sanquinolentus</i>		<b>Nanoclavellia</b> Haupt, 1962
<i>sanquinolentus</i>		
(Fabricius, 1793)	B M S	
<i>sanquinolentus</i>		
<i>balcanicus</i> Haupt, 1927	M S	
POMPILINI		
<b>Pompilus</b> Fabricius, 1798		
<i>plumbeus</i> <i>plumbeus</i>		
(Fabricius, 1787)	B M S	
<b>Arachnospila</b> Kincaid, 1900		
<i>abnormis</i> (Dahlbom, 1842)	B M	
<i>anceps</i> <i>anceps</i>		
(Wesmael, 1851)	B M S	
<i>ausa</i> (Tournier, 1890)	B M S	
<i>bohemabnormis</i> (Wolf, 1966)	B	
<i>fumipennis</i> <i>fumipennis</i>		
(Zetterstedt, 1838)	B M S	
<i>fuscomarginata</i> (Thomson, 1870)	B M S	
<i>luctuosa gibbomima</i>		
(Haupt, 1929)	B S	
<i>minutula minutula</i>		
(Dahlbom, 1842)	B M S	
<i>minutula simplicicrus</i>		
(Priesner, 1960)	M	
<i>opinata</i> (Tournier, 1890)	B M S	
<i>pseudabnormis</i> (Wolf, 1965)	B M	
<i>rufa rufa</i> (Haupt, 1927)	B M	
<i>sogdiana sogdiana</i>		
(Haupt, 1927)	B M	
<i>spissa spissa</i> (Schioedte, 1837)	B M S	
<i>trivialis trivialis</i>		
(Dahlbom, 1843)	B M S	
<i>usurata usurata</i>		
(Blüthgen, 1957)	B M	
<i>wesmaeli</i> (Thomson, 1870)	B M S	
<b>Agenioideus</b> Ashmead, 1902		
<i>apicalis apicalis</i>		
(Van der Linden, 1827)	M	
<i>ciliatus</i> (Lepeletier, 1845)	M	
<i>cinctellus</i> (Spinola, 1808)	B M S	
<i>nubecula</i> (Costa, 1874)	B M S	
<i>sericeus</i> (Van der Linden, 1827)	B M S	
<i>usurarius</i> (Tournier, 1889)	B M S	
<b>leucopterus</b> (Dahlbom, 1845)		<b>Nanoclavellia</b> Haupt, 1962
<b>Arachnotheutes</b> Haupt, 1927		
<i>rufithorax</i> (Costa, 1887)	S	
<b>Tachyagetes</b> Haupt, 1930		
<i>dudichi</i> Mócsar, 1944	M S	
<i>filicornis filicornis</i>		
(Tournier, 1890)	B M S	
<b>Aporinellus</b> Banks, 1911		
<i>obtusus</i> (Gussakovskij, 1935)	M S	
<i>sexmaculatus sexmaculatus</i>		
(Spinola, 1805)	B M S	
<b>Dicyrtomellus</b>		
Gussakovskij, 1935		
<i>luctuosus luctuosus</i>		
(Mocsary, 1879)	S	
<b>Anospilus</b> Haupt, 1929		
<i>orbitalis orbitalis</i>		
(Costa, 1863)	S	
<b>Evagetes</b> Lepeletier, 1845		
<i>alamannicus</i> (Blüthgen, 1944)	B S	
<i>contemptus villicus</i>		
(Tournier, 1890)	M S	
<i>crasicornis crassicornis</i>		
(Shuckard, 1837)	B M S	
<i>dubius</i> (Van der Linden, 1827)	B M S	
<i>littoralis</i> (Wesmael, 1851)	B M S	
<i>magrettii</i> (Kohl, 1886)	M S	
<i>pectinipes pectinipes</i>		
(Linnaeus, 1758)	B M S	
<i>pilosellus</i> (Wesmael, 1851)	B M S	
<i>pontomoravicus</i>		
(Šustera, 1938)	B M S	
<i>proximus</i> (Dahlbom, 1843)	B M S	
<i>subglaber</i> (Haupt, 1941)	B S	
<i>subnudus</i> (Haupt, 1942)	B M S	
<i>tumidosus sinuatus</i>		
(Haupt, 1941)	B M S	
<b>Anoplus</b> Dufour, 1834		
<i>alpinobalticus</i> Wolf, 1965	S	

<i>caviventris</i> Aurivillius 1907	S	<i>ordinarius ordinarius</i> Priesner, 1966	B M S
<i>infuscatus infuscatus</i> (Van der Linden, 1827)	B M S	<i>rufipes rufipes</i> (Linnaeus, 1758)	B M S
<i>nigerrimus</i> (Scopoli, 1763)	B M S	<b>Telostegus</b> Costa, 1887	
<i>pannonicus</i> Wolf, 1965	S	<i>inermis inermis</i> (Brullé, 1832)	S
<i>tenuicornis</i> (Tournier, 1889)	B	CEROPALINAE	
<i>viaticus viaticus</i> (Linnaeus, 1758)	B M S	CEROPALINI	
<i>viaticus paganus</i> (Dahlbom, 1843)	B M S	<b>Ceropales</b> Latreille, 1796	
<b>Batozonellus</b> Arnold, 1937		<i>albicinctus albicinctus</i> (Rossi, 1790)	M S
<i>lacerticida lacerticida</i> (Pallas, 1771)	B M S	<i>cribratus cribratus</i> Costa, 1881	S
<b>Episyron</b> Schioedte, 1837		<i>helveticus</i> Tournier, 1889	S
<i>arrogans</i> (Smith, 1873)	B M S	<i>maculatus maculatus</i> (Fabricius, 1775)	B M S
<i>funereipes</i> (Costa, 1881)		<i>pygmaeus</i> Kohl, 1879	M S
		<i>variegatus</i> (Fabricius, 1798)	B M S

## SPHECOIDEA

ZDENĚK PÁDR

So far, 287 species of the superfamily Sphecoidea have been recorded in the territory of Czechoslovakia. A number of entomologists have been engaged in the faunal investigations of this group already since the end of the nineteenth century. The greatest credit for the knowledge of the occurrence of Sphecoidea in our territory is due to P. A. Kubes, J. Slavíček, O. Šustera, V. Zavadil, J. Šnoflák, A. Hoffer, M. Kocourek, V. Balthasar, M. Hrubant, K. Deneš, J. Niedl, P. Tyrner, the present author and others. Most of the collections of the entomologists mentioned are deposited in National Museum in Prague, and in the Moravian Museum in Brno. Part of the collections is private.

The species recorded in the territory of Czechoslovakia before 1937 have been published by V. Zavadil et al. (1937). Later V. Zavadil et J. Šnoflák (1948) elaborated a very good key to the determination of Sphecidae of Czechoslovakia. In detail, the superfamily has been treated in the monograph by V. Balthasar (1972).

Sphecoidea is a group of predatory insects whose larvae feed on other arthropods which the female paralyses with her sting and deposits in a tunnel excavated in the soil. The female then lays on the paralysed prey an egg from which a euparasitic larva hatches.

Most Sphecoidea are thermophilous insects which frequent light sandy soils. The most thoroughly investigated regions include the Labe (Elbe) basin, southern Moravia and southern Slovakia. The occurrence of Sphecoidea in higher mountainous parts of the territory has been less thoroughly investigated. These regions may be penetrated by northern species hitherto unknown in Czechoslovakia.

The check-list of the superfamily Sphecoidea has been arranged according to the classification used by J. de Beaumont (1964). Opinions on the synonymy of individual species have been adopted from more recent papers of W. Pulawski (1974) and his critique of the above-mentioned monograph by V. Balthasar.

References: Balthasar, V., 1972: Grabwespen — Sphecoidea. Fauna ČSSR, Vol. 20, Academia, Praha, 471 pp. — de Beaumont, J.: 1964: Hymenoptera: Sphecidae. Insecta helvetica 3: 169 pp. — Pulawski, W., 1974: Book review of the monograph by Balthasar 1972. *Acta ent. bohemosl.* 71: 349—350. — Zavadil, V., Šustera O., Baťa, L., 1937: Sphecoidea. in *Prodromus Hymenopterorum Čechoslovakiæ*, Pars. 1. *Sbor. ent. odděl. Nář. Mus.* XV: 145—222. — Zavadil, V., Šnoflák, J., 1948: *Kutilky Československé republiky*. Ent. Handbook of Ent. Listy 13: 179 pp.

SPHECOIDEA			
SPHECIDAE			
<b>Ammophila</b> Kirby, 1798			
<i>affinis</i> (Kirby, 1798)	B M S		
<i>apicalis</i> Brullé, 1838	B M S		
<i>campestris</i> Latreille, 1809	B M S		
<i>fallax</i> Kohl, 1883	M S		
<i>heydeni</i> Dahlbom, 1845	B M S		
<i>hirsuta</i> (Scopoli, 1763)	B M S		
<i>luffi</i> Saunders, 1903	B M S		
<i>pubescens</i> Curtis, 1829	B M S		
<i>susterai</i> Šnoflák, 1943			
<i>sabulosa</i> (Linnaeus, 1758)	B M S		
<i>tydei</i> (Guillon, 1841)	S		
<b>Sphex</b> Linnaeus, 1758			
<i>albisectus</i> Lepeletier et Serville, 1825	S		
<i>atrophirtus</i> Kohl, 1885	S		
<i>maxillosus</i> Fabricius, 1793	B M S		
<i>occitanicus</i> misident.			
<i>subfuscatus</i> Dahlbom, 1845	M S		
<b>Sceliphron</b> Klug, 1801			
<i>destillatorium</i> (Illiger, 1807)	M S		
<i>femoratum</i> (Fabricius, 1781)	S		
AMPULICIDAE			
<b>Dolichurus</b> Latreille, 1809			
<i>corniculus</i> (Spinola, 1808)	B M S		
<b>Ampulex</b> Jurine, 1807			
<i>fasciata</i> Jurine, 1807	B M S		
PHILANTHIDAE			
<b>Philanthus</b> Fabricius, 1790			
<i>coronatus</i> Fabricius, 1790	M S		
<i>triangulum</i> (Fabricius, 1775)	B M S		
<i>venustus</i> (Rossi, 1790)	M S		
<b>Cerceris</b> Latreille, 1802			
albofasciata (Rossi, 1790)		S	
<i>luctuosa</i> Costa, 1869			
arenaria (Linnaeus, 1758)	B M S		
<i>bicincta</i> Klug, 1835		S	
<i>variolosa</i> Costa, 1869			
<i>bracteata</i> Eversmann, 1849		S	
<i>circularis dacica</i>			
Schletterer, 1887		S	
<i>eryngii</i> Marquet, 1875		S	
<i>fimbriata</i> (Rossi, 1790)		S	
<i>funerea</i> Schletterer, 1887			
<i>flavilabris</i> (Fabricius, 1793)		B M S	
<i>jerreri</i> Van der Linden, 1829			
<i>hortivaga</i> Kohl, 1880	B M S		
<i>impercepta</i> Beaumont, 1950		S	
<i>somotorensis</i> Balthasar, 1955			
<i>interrupta</i> (Panzer, 1799)	B M S		
<i>quadricincta</i> (Villers, 1799)	B M S		
<i>quadrifasciata</i> (Panzer, 1799)	B M S		
<i>quinquefasciata</i>			
(Rossi, 1792)	B M S		
<i>rubida</i> (Jurine, 1807)		S	
<i>ruficornis</i> (Fabricius, 1793)		B M S	
<i>cunicularia</i> (Schrank, 1802)			
<i>rufipes</i> (Fabricius, 1787)		S	
<i>tuberculata</i> (Villers, 1789)			
<i>rybiensis</i> (Linnaeus, 1771)	B M S		
<i>sabulosa</i> (Panzer, 1799)	B M S		
<i>stratiotes</i> Schletterer, 1887		S	
<i>tenuivittata</i> Dufour, 1849		S	
NYSSONIDAE			
<b>Argogorytes</b> Ashmead, 1899			
<i>fargei</i> (Shuckard, 1837)	B M S		
<i>mystaceus</i> (Linnaeus, 1761)	B M S		
<b>Gorytes</b> Latreille, 1805			
<i>affinis</i> Spinola, 1808	B M S		

<i>albidulus</i> (Lepeletier, 1832)	B M S	<i>olivacea</i> Cyrillo, 1787	S
<i>dissectus</i> (Panzer, 1801)		<i>mediterranea</i> Handlirsch, 1893	
<i>bicinctus</i> (Rossi, 1792)	B M S	<i>rostrata</i> (Linnaeus, 1758)	B M S
<i>bilunulatus</i> (Costa, 1869)	M S		
<i>craverrii</i> (Costa, 1869)	M S		
<i>ottomanus</i> Mocsáry, 1879			
<i>elegans</i> (Lepeletier, 1832)	B M S	<b>Nysson</b> Latreille, 1796	
<i>fallax</i> Handlirsch, 1888	B M S	<i>bohemicus</i> Zavadil, 1950	B
<i>formosus</i> Jurine, 1807	B	<i>dimidiatus</i> Jurine, 1807	B M S
<i>foveolatus</i> Handlirsch, 1888	S	<i>fulvipes</i> Costa, 1859	S
<i>laevis</i> (Latreille, 1792)	B M S	<i>hrubanti</i> Balthasar, 1972	M S
<i>laticinctus</i> (Lepeletier, 1832)	B M S	<i>interruptus</i> (Fabricius, 1798)	B M S
<i>latifrons</i> Spinola, 1808	M S	<i>maculatus</i> (Fabricius, 1787)	B M S
<i>lunatus</i> (Dahlbom, 1832)	B M S	<i>niger</i> Chevrier, 1868	B M
<i>moravicus</i> Šnoflák, 1943	M S	<i>quadriguttatus</i>	
<i>nigrifacies</i> Mocsáry, 1879	S	Gerstaecker, 1866	B S
<i>pleuripunctatus</i> (Costa, 1869)	M S	<i>roubali</i> Zavadil, 1937	S
<i>procrustes</i> Handlirsch, 1888	M S	<i>scalaris</i> Illiger, 1807	B M S
<i>punctuosus</i> Handlirsch, 1888	B M S	<i>spinosus</i> (Foerster, 1771)	B M S
<i>punctatus</i> (Kirschbaum, 1853)		<i>susterai</i> Zavadil, 1950	M
<i>quadrifasciatus</i> (Fabricius, 1804)	B M S	<i>tridens</i> Gerstaecker, 1866	B M S
<i>quinquecinctus</i> (Fabricius, 1793)	B M S	<i>trimaculatus</i> (Rossi, 1790)	B M S
<i>quinqefasciatus</i> (Panzer, 1798)	B M S	<i>variabilis</i> Chevrier, 1867	S
<i>sareptanus</i> Handlirsch, 1888	M S	<b>Allyon</b> Panzer, 1806	
<i>suclifrons</i> (Costa, 1869)	S	<i>fuscatus</i> (Panzer, 1798)	B M S
<i>tauricus</i> Radoszkowski, 1884	S	<i>pertheesi</i> Gorski, 1852	S
<i>tumidus</i> (Panzer, 1801)	B M S	<i>ratzeburgi</i> Dahlbom, 1843	B M
<b>Stizus</b> Latreille, 1802		<i>tricolor</i> Lepeletier et Serville, 1825	M S
<i>perrisi</i> Dufour, 1838	B M S	<b>Didineis</b> Wesmael, 1852	
<b>Bembecinus</b> Costa, 1859		<i>crassicornis</i> Handlirsch, 1887	S
<i>hungaricus</i> (Frivaldzsky, 1876)	M S	<i>unicornis</i> (Fabricius, 1798)	B M S
<i>tridens</i> (Fabricius, 1781)	B M S	<i>wüstneii</i> Handlirsch, 1887	*M
<b>Bembix</b> Fabricius, 1775		<b>Mellinus</b> Fabricius, 1790	
<i>bidentata</i> Van der Linden, 1829	S	<i>arvensis</i> (Linnaeus, 1758)	B M S
<i>integra</i> Panzer, 1805	B M S	<i>sabulosus</i> (Fabricius, 1787)	B M S
<i>megerlei</i> Dahlbom, 1845	S	LARRIDAE	
		<b>Larra</b> Fabricius, 1793	
		<i>anathema</i> (Rossi, 1790)	M S
		<b>Liris</b> Fabricius, 1804	
		<i>nigra</i> (Van der Linden, 1829)	S

<b>Tachysphex</b> Kohl, 1883							
<i>consocius</i> Kohl, 1892	S						
<i>coriaceus</i> (Costa, 1867)	S						
<i>fulvitarsis</i> (Costa, 1867)	B M S						
<i>grandii</i> Beaumont, 1969	S						
<i>helveticus</i> Kohl, 1884	B M S						
<i>mocsaryi</i> Kohl, 1883	S						
<i>nigripennis</i> (Spinola, 1808)	B M S						
<i>nitidus</i> (Spinola, 1805)	B M S						
<i>obscuripennis</i> (Schenck, 1857)	B M S						
<i>lativalvis</i> (Thomson, 1870)							
<i>panzeri</i> (Van der Linden, 1829)	B M S						
<i>pectinipes</i> (Linnaeus, 1758)	B M S						
<i>psammobius</i> Kohl, 1883	B M S						
<i>pygidialis</i> Kohl, 1883	S						
<i>spoliatus</i> (Giraud, 1863)	B M S						
<b>Tachytes</b> Panzer, 1806							
<i>etruscus</i> (Rossi, 1790)	S						
<i>europeus</i> Kohl, 1883	B M S						
<i>obsoletus</i> (Rossi, 1792)	S						
<b>Solierella</b> Spinola, 1851							
<i>compendita</i> (Piccioli, 1869)	B M S						
<b>Misophus</b> Jurine, 1807							
<i>ater</i> Lepeletier, 1845	B						
<i>bicolor</i> Jurine, 1807	B M S						
<i>concolor</i> Dahlbom, 1845	B M S						
<i>moravicus</i> Balthasar, 1957							
<i>spurius</i> (Dahlbom, 1832)	B M S						
<b>Nitela</b> Latreille, 1809							
<i>fallax</i> Kohl, 1883	B M						
<i>spinolai</i> Latreille, 1809	B M S						
<b>Dinetus</b> Jurine, 1807							
<i>pictus</i> (Fabricius, 1793)	B M S						
<b>Palarus</b> Latreille, 1802							
<i>flavipes</i> (Fabricius, 1781)	M S						
ASTATIDAE							
<b>Astata</b> Latreille, 1796							
<i>apostata</i> Mercet, 1910	S						
<i>boops</i> (Schrank, 1781)	B M S						
<i>costai</i> Piccioli, 1869	S						
<b>femoralis</b> Mocsáry, 1877	B M S						
<i>gallica</i> Beaumont, 1942	S						
<i>lineata</i> Mocsáry, 1879	S						
<i>minor</i> Kohl, 1884	B M S						
<i>rufipes</i> Mocsáry, 1883	S						
<i>stecki</i> Beaumont, 1942	B M S						
<i>stigma</i> (Panzer, 1809)	B M S						
<i>tricolor</i> (Van der Linden, 1829)	M S						
PEMPHREDONIDAE							
<b>Psen</b> Latreille, 1796							
<i>ater</i> (Fabricius, 1794)	B M S						
<i>atratinus</i> (Morawitz, 1891)	B S						
<i>bicolor</i> Jurine, 1807	B M S						
<i>bruxellensis</i> (Bondroit, 1933)	B						
<i>caucasicus</i> Maidl, 1914	M S						
<i>crassipes</i> Costa, 1866	M S						
<i>dahlbomi</i> (Wesmael, 1852)	B M S						
<i>equestris</i> (Fabricius, 1804)	B M S						
<i>littoralis</i> Bondroit, 1933	M S						
<i>fulvitarsis</i> Gussakovskij, 1937							
<i>shuckardi</i> (Wesmael, 1852)	B M S						
<i>tenuis</i> Oehlke, 1965	S						
<i>unicolor</i> Van der Linden, 1829	B M S						
<b>Psenulus</b> Kohl, 1896							
<i>concolor</i> (Dahlbom, 1843)	B M						
<i>fuscipennis</i> (Dahlbom, 1843)	B M S						
<i>laevigatus</i> (Schenck, 1857)	B M						
<i>pallipes</i> (Panzer, 1798)	B M S						
<i>atratus</i> (Fabricius, 1804)							
<i>schencki</i> (Tournier, 1889)	B M S						
<b>Pemphredon</b> Latreille, 1796							
<i>austriacus</i> (Kohl, 1888)	B M						
<i>clypealis</i> Thomson, 1970	B M S						
<i>inornatus</i> Say, 1824	B M S						
<i>shuckardi</i> (Morawitz, 1864)							
<i>lethifer</i> (Shuckard, 1837)	B M S						
<i>lugubris</i> Latreille, 1805	B M S						
<i>lugens</i> Dahlbom, 1842	B M S						
<i>montanus</i> Dahlbom, 1845	B M S						

<i>morio</i> Van der Linden, 1829	B M S	<i>fronticorne</i> Gussakovskij, 1936	M S
<i>rugifer</i> Dahlbom, 1843	B M S	<i>kolazyi</i> Kohl, 1893	B M S
<i>unicolor</i> Panzer, 1798		<i>medium</i> Beaumont, 1945	M
<b>Diodontus</b> Curtis, 1834		<i>minus</i> Beaumont, 1945	B
<i>luperus</i> Shuckard, 1837	B M S	<i>scutatum</i> Chevrier, 1867	M S
<i>maior</i> Kohl, 1901	M S		
<i>minutus</i> (Fabricius, 1793)	B M S		
<i>tristis</i> (Van der Linden, 1829)	B M S		
<b>Passaloecus</b> Shuckard, 1837			
<i>borealis</i> Dahlbom, 1845	B	<b>Rhopalum</b> Kirby, 1829	
<i>corniger</i> Shuckard, 1837	B M S	<i>austriacum</i> Kohl, 1899	M S
<i>eremita</i> Kohl, 1893	B M S	<i>clavipes</i> (Linnaeus, 1758)	B M
<i>gracilis</i> (Curtis, 1834)	B M S	<i>coarctatum</i> (Scopoli, 1763)	B M S
<i>insignis</i> (Van der Linden, 1829)	B M S	<i>tibiale</i> (Fabricius, 1798)	
<i>monilicornis</i> Dahlbom, 1842	B M S	<i>nigrinum</i> Kiesenwetter, 1849	B S
<b>Stigmus</b> Panzer et Jurine, 1805			
<i>pendulus</i> Panzer, 1805	B M S	<b>Entomognathus</b> Dahlbom, 1845	
<i>solskyi</i> A. Morawitz, 1864	B M S	<i>brevis</i> (Van der Linden, 1829)	B M S
<b>Spilomena</b> Shuchard, 1837		<i>dentifer</i> Noskiewicz, 1930	B M S
<i>beata</i> Blüthgen, 1953	B	<b>Lindenius</b> Lepeletier et Brullé, 1834	
<i>differens</i> Blüthgen, 1953	B	<i>albibasis</i> (Fabricius, 1793)	B M S
<i>mocsaryi</i> Kohl, 1898	M S	<i>laevis</i> Costa, 1871	B S
<i>zavadili</i> Šnoflák, 1942		<i>rhaibopus</i> Kohl, 1915	
<i>troglodytes</i> (Van der Linden, 1829)	B M S	<i>panzeri</i> (Van der Linden, 1829)	B M S
<i>vagans</i> Blüthgen, 1953	B S	<i>parkanensis</i> Zavadil, 1948	S
<b>Ammoplanus</b> Giraud, 1869		<i>ponticus</i> Beaumont, 1956	
<i>handlirschi</i> Gussakovskij, 1931	B M	<i>pygmaeus armatus</i>	
<i>hofferi</i> Šnoflák, 1943	M S	<i>(Van der Linden, 1829)</i>	B M S
<i>pragensis</i> Šnoflák, 1945	B	<i>subaeneus</i> Lepeletier et Brullé, 1834	S
<i>wesmaeli</i> Giraud, 1869	B M S		
<i>perrisi</i> Giraud, 1869		<b>Crossocerus</b> Lepeletier et Brullé, 1834	
TRYPOXYLONIDAE		<i>acanthophorus</i> Kohl, 1892	S
<b>Pison</b> Spinola, 1808		<i>annulipes</i> Lepeletier	
<i>atrum</i> Spinola, 1808	S	<i>et Brullé, 1834</i>	B M S
<b>Trypoxyylon</b> Latreille, 1796		<i>ambiguus</i> (Dahlbom, 1842)	
<i>attenuatum</i> Smith, 1851	B M S	<i>assimilis</i> (Smith, 1856)	B M S
<i>clavicerum</i> Lepeletier et Serville, 1825	B M S	<i>tiroensis</i> Kohl, 1879	
<i>figulus</i> (Linnaeus, 1758)	B M S	<i>barbipes</i> (Dahlbom, 1845)	B M
		<i>binotatus</i> Lepeletier et Brullé, 1834	B M S
		<i>confusus</i> Schulz, 1906	

<i>capitosus</i> (Shuckard, 1837)	B M	<i>laponicus</i> Zetterstedt, 1838	B M S
<i>cetratus</i> (Shuckard, 1837)	B M S	<i>loewi</i> Dahlbom, 1845	M
<i>cinxius</i> (Dahlbom, 1838)	B M S	<i>peltarius</i> (Schreber, 1784)	B M S
<i>congener</i> (Dahlbom, 1845)	B M	<i>scutellatus</i> (Scheven, 1781)	B M S
<i>curvitarsis</i> (Herrich et Schaeffer, 1840)	M		
<i>denticoxa</i> Bischoff, 1932	M	<b>Lestica</b> Billberg, 1820	
<i>denticrus</i> (Herrich- -Schaeffer, 1841)	B M	<i>alata</i> (Panzer, 1797)	B M S
<i>distinguendus</i> (Morawitz, 1866)	B M S	<i>clypeata</i> (Linnaeus, 1767)	B M S
<i>dimidiatus</i> (Fabricius, 1781)	B M S	<i>crassicornis</i> (Spinola, 1808)	M S
<i>elongatus</i> (Linden, 1829)	B M S	<i>subterranea</i> (Fabricius, 1775)	B M S
<i>exiguus</i> (Van der Linden, 1829)	B M S	<b>Ectemnius</b> Dahlbom, 1845	
<i>heydeni</i> Kohl, 1880	B M S	<i>borealis</i> (Zetterstedt, 1838)	B M S
<i>imitans</i> Kohl, 1915	S	<i>nigrinus</i> (Herrich et Schaeffer, 1841)	
<i>inermis</i> Thomson, 1870	B M	<i>cavifrons</i> (Thomson, 1870)	B M S
<i>leucostomus</i> (Linnaeus, 1758)	B M S	<i>cephalotes</i> (Olivier, 1811)	B M S
<i>megacephalus</i> (Rossi, 1790)	B M S	<i>quadricinctus</i> (Fabricius, 1787)	
<i>leucostomoides</i> Richards, 1935		<i>continuus</i> (Fabricius, 1804)	B M S
<i>ovalis</i> Lepeletier et Brullé, 1834	B M S	<i>vagus</i> (Linnaeus, 1758)	
<i>anxius</i> Wesmael, 1852		<i>dives</i> (Lepeletier et Brullé, 1834)	B M S
<i>palmipes</i> (Linnaeus, 1767)	B M S	<i>fessorius</i> (Linnaeus, 1758)	B S
<i>podagricus</i> (Van der Linden, 1829)	B M S	<i>guttatus</i> (Van der Linden, 1829)	B M S
<i>snojaki</i> Zavadil, 1948		<i>spinicollis</i> (Herrich et Schaeffer, 1841)	
<i>pubescens</i> (Shuckard, 1837)	B M S	<i>impressus</i> Smith, 1856	S
<i>pusillus</i> Lepeletier et Brullé, 1834	B M S	<i>laevigatus</i> (De Stefani, 1884)	B M S
<i>varius</i> Lepeletier et Brullé, 1834		<i>lapidarius</i> (Panzer, 1804)	B M S
<i>quadrimaculatus</i> (Fabricius, 1793)	B M S	<i>chrysostomus</i> (Lepeletier et Brullé, 1834)	
<i>styrius</i> (Kohl, 1892)	M	<i>lituratus</i> (Panzer, 1805)	M S
<i>tarsatus</i> (Shuckard, 1837)	B M	<i>nigritarsus</i> (Herrich et Schaeffer, 1841)	M S
<i>vagabundus</i> (Panzer, 1798)	B M S	<i>rubicola</i> (Dufour et Perris, 1840)	B M S
<i>walkeri</i> (Shuckard, 1837)	B M	<i>ruficornis</i> (Zetterstedt, 1838)	B M S
<i>wesmaeli</i> (Van der Linden, 1829)	B M S	<i>nigrifrons</i> Cresson, 1865	
<b>Crabro</b> Fabricius, 1775		<i>planifrons</i> Thomson, 1870	
<i>cribrarius</i> (Linnaeus, 1758)	B M S	<i>rugifer</i> Dahlbom, 1845	B S

<i>sexcinctus</i> (Fabricius, 1775)	B M S	<i>melancholicus</i> Chevrier, 1868	B M S
<i>zonatus</i> (Panzer, 1797)		<i>mucronatus</i> (Fabricius, 1793)	B M S
<i>schlettereri</i> Kohl, 1888	B*M S	<i>pugnax</i> Olivier, 1811	B M S
<i>spinipes</i> (Morawitz, 1866)	S	<i>quatuordecimnotatus</i> Jurine, 1807	B M S
<b>Oxybelus</b> Latreille, 1796		<i>trispinosus</i> (Fabricius, 1787)	B M S
<i>argentatus</i> Curtis, 1833	M S	<i>nigripes</i> Olivier, 1811	B M S
<i>aurantiacus</i> Mocsáry, 1883	S	<i>uniglumis</i> (Linnaeus, 1758)	B M S
<i>bipunctatus</i> Olivier, 1811	B M S	<i>variegatus</i> Wesmael, 1852	B M S
<i>dissectus elegans</i> Mocsáry, 1879	S	<i>victor</i> Lepeletier, 1845	B M S
<i>latidens</i> Gerstaecker, 1867	B M S	<b>Belomicrus</b> Costa, 1866	
<i>latro</i> Olivier, 1811	M S	<i>antennalis</i> Kohl, 1899	M S
<i>lineatus</i> (Fabricius, 1787)	B S	<i>italicus</i> Costa, 1866	B M S
<i>maculipes</i> Smith, 1856	S	<i>obscurus</i> Kohl, 1892	
<i>mandibularis</i> Dahlbom, 1845	B M S		
<i>sericatus</i> Gerstaecker, 1867			



## APOIDEA

MIROSLAV KOCOUREK

The beginnings of investigations into solitary bees in the territory of Czechoslovakia date back to the end of the nineteenth and early twentieth centuries. In that period, J. Slaviček (1895) and A. Ducke studied solitary bees in Moravia, P. Kubes (1904) in Bohemia.

The late entomologists O. Šustera, V. Zavadil, L. Bařa, J. Šnoflák, F. Gregor, sen., J. May, V. Balthasar, A. Hoffer and others were the pioneers of Czechoslovak apidology. Their perfectly labelled collections, faunal papers, careful files of O. Šustera as well as experience and data submitted by the above entomologists to the present author provided a solid basis for the compilation of this check-list. Their collections are deposited in the National Museum in Prague, the Moravian Museum in Brno and the Regional Museum in České Budějovice. Investigations into Apoidea in Czechoslovakia were substantially augmented by J. Palásek, F. Urbášek, Z. Pádr, B. Tkalců and the present author Kocourek (1963). The collections of these entomologists are predominantly private.

The superfamily Apoidea comprises the most numerous group of pollinators of entomophilous plants. Some of the species are intentionally utilised for this activity (*Megachile*, *Rhophotoides*, *Apis*, etc.). The species of the family Apidae are social (*Bombus*, *Apis*) or parasitic (*Psithyrus*). Some species of the families Anthophoridae and Halictidae are parasocial or eusocial. Most species are solitary in the nesting behavior. The larvae of some (e.g., those of the genera *Nomada*, *Sphecodes*, *Coelioxys*, *Epeorus*) are cleptoparasitic in other species of bees.

Relatively most thorough faunal studies have been made in central Bohemia, the České Středohoří Mts., central and southern Moravia, southern and southeastern Slovakia. Data on the occurrence of solitary bees in the rest of the territory are sporadic so that future investigations may be expected to yield discoveries of additional species.

In addition to smaller papers by Czechoslovak and foreign authors, the works of O. Schmiedeknecht (1930) and Bouček et Šustera (1957) were used as guidelines for nomenclature and classification in compiling this check-list.

References: Bouček, Z. and Šustera, O. (in Kratochvíl J. et al.), 1957: Včely — Apoidea. Klíč zvřízený ČSR, Praha: 377—406. — Kocourek, M., 1963: Beitrag zur Kenntnis der aculeaten Hymenopteren der Tschechoslowakei. *Acta faun. ent. Mus. Nat. Pragae*, 9: 285—298. — Kocourek, M., 1966: Prodromus der Hymenopteren der Tschoslowakei, Pars 9: Apoidea 1. *Acta faun. ent. Mus. Nat. Pragae* 12: 1—121. — Kubes, A., 1904:

Fauna Bohemica. Conspectus Apidarum, quas in Bohemia collegi. *Acta Soc. ent. Boh.* 1: 26—31. — Schmiedeknecht, O., 1930: Die Hymenopteren Nord- und Mitteleuropas, Jena, 1062 p. — Slavíček, J., 1895: Beitrag zur Fauna v. Mähren. Die Bienen der Umgebung von Mikulov. *Verh. naturf. Ver. Brünn*, 33: 98—104.

APODEA COLLETIDAE			
<b>Colletes</b> Latreille, 1802			
<i>cunicularius</i> (Linnaeus, 1758)	B M S	<i>euryscapa</i> (Foerster, 1891)	M S
<i>daviesanus</i> Smith, 1846	B M S	<i>gibba confusa</i> (Nylander, 1852)	B M S
<i>floralis</i> Eversmann, 1852	B	<i>gracilicornis</i> Morawitz, 1867	B M S
<i>fodiens</i> (Geoffroy in Fourcroy, 1785)	B M S	<i>hyalinata</i> Smith, 1843	B M S
<i>graeffei</i> Alfken, 1900	M S	<i>lineolata</i> Schenck, 1859	B M S
<i>hylaeiformis</i> Eversmann, 1852	M S	<i>nigrifacies</i> (Bramson, 1879)	B M S
<i>impunctatus</i> Nylander, 1852	M	<i>nigrita</i> (Fabricius, 1798)	B M S
<i>inexpectatus</i> Noskiewicz, 1936	B M S	<i>pectoralis</i> (Foerster, 1871)	B M
<i>maidli</i> Noskiewicz, 1936	S	<i>pfankuchi</i> Alfken, 1915	M S
<i>marginatus</i> Smith, 1846	B M S	<i>pictipes</i> (Nylander, 1852)	B M S
<i>mlokossewiczi</i> Radoszkowski, 1891	S	<i>punctata</i> Brullé, 1832	B M S
<i>nasutus</i> Smith, 1853	B M S	<i>punctulatissima</i> Smith, 1843	B M S
<i>punctatus</i> Mocsáry, 1877	M S	<i>rinki</i> Gorski, 1852	B M S
<i>similis</i> Schenck, 1853	B M S	<i>signata</i> (Panzer, 1798)	B M S
<i>spectabilis</i> Morawitz, 1868	B M S	<i>pratensis</i> (Geoffroy in Fourcroy, 1785)	
<i>succinctus</i> (Linnaeus, 1758)	B M S	<i>sinuata</i> (Schenck, 1853)	M S
<b>Prosopis</b> Fabricius, 1804		<i>styriaca</i> (Foerster, 1871)	B M S
<i>angustata</i> Schenck, 1859	B M S	<i>variegata</i> (Fabricius, 1798)	B M S
<i>annularis</i> (Kirby, 1802)	B M S		ANDRENIDAE
<i>annulata</i> (Linnaeus, 1758)	B M S	<b>Andrena</b> Fabricius, 1775	
<i>bifasciata</i> Jurine, 1807	?	<i>aciculata</i> Morawitz, 1896	B M S
<i>bissinuata</i> Foerster, 1871	B M S	<i>aeneiventris</i> Morawitz, 1872	M S
<i>brevicornis</i> (Nylander, 1852)	B M S	<i>agilissima</i> (Scopoli, 1770)	B M S
<i>clypearis</i> Schenck, 1853	M S	<i>albofasciata</i> Thomson, 1870	B M S
<i>communis</i> (Nylander, 1852)	B M S	<i>albopunctata</i> (Rossi, 1792)	S
<i>cornuta</i> (Curtis, 1831)	M S	<i>alfkenella</i> Perkins, 1914	B M S
<i>difformis</i> Eversmann, 1852	B M S	<i>alutacea</i> Stöckhert, 1942	M
<i>duckei</i> Alfken, 1904	M S	<i>anthrisci</i> Blüthgen, 1925	?
		<i>apicata</i> Smith, 1847	B M S
		<i>argentata</i> Smith, 1844	B M S
		<i>assimilis</i> Radoszkowski, 1868	B M S
		<i>atrata</i> Friese, 1887	M S

<i>barbareae</i> Panzer, 1805	B M S	<i>fulvicornis</i> Schenck, 1853	B M S
<i>barbilabris</i> (Kirby, 1802)	B M S	<i>fulvida</i> Schenck, 1853	B M S
<i>batava</i> Pérez, 1902	B M	<i>fuscipes</i> (Kirby, 1802)	B M S
<i>bicolor</i> Fabricius, 1775	B M S	<i>fuscosa</i> Erichson, 1835	M S
<i>bimaculata</i> (Kirby, 1802)	B M S	<i>gelriae</i> van der Wecht, 1927	B M S
<i>bisulcata</i> Morawitz, 1876	M S	<i>granulosa</i> Pérez, 1902	B M S
<i>bluethgeni</i> Stöckhert, 1930	B	<i>gravida</i> Imhoff, 1832	B M S
<i>braunsiana</i> Friese, 1887	S	<i>haemorrhoa</i> (Fabricius, 1781)	B M S
<i>bucephala</i> Stephens, 1846	S	<i>hatorfiana</i> (Fabricius, 1775)	B M S
<i>carbonaria</i> (Linnaeus, 1767)	B M S	<i>hedickae</i> Jaeger, 1934	S
<i>chrysopus</i> Pérez, 1902	M S	<i>helvola</i> Linnaeus, 1758	B M S
<i>chrysopyga</i> Schenck, 1832	B M S	<i>hesperia</i> Smith, 1853	S
<i>chrysosceles</i> (Kirby, 1802)	B M S	<i>humilis</i> Imhoff, 1832	B M S
<i>cineraria</i> (Linnaeus, 1758)	B M S	<i>hungarica</i> Friese, 1887	S
<i>cinerascens</i> Eversmann, 1852	B M S	<i>intermedia</i> Thomson, 1872	M
<i>clarkella</i> (Kirby, 1802)	B M S	<i>impunctata</i> Pérez, 1895	B M S
<i>clypella hasitata</i> Warncke, 1973	S	<i>paula</i> Noskiewicz, 1938	
<i>coitana</i> (Kirby, 1802)	B M S	<i>labialis</i> (Kirby, 1802)	B M S
<i>combaella</i> Warncke, 1966	S	<i>labiata</i> (Fabricius, 1781)	B M S
<i>combinata</i> (Christ, 1791)	B M S	<i>lapponica</i> Zetterstedt, 1838	B M S
<i>confinis</i> Stöckhert, 1929	B M S	<i>lathyri</i> Alfken, 1899	B M S
<i>congruens</i> Smiedeknecht, 1883	B M S	<i>lepidia</i> Schenck, 1859	M S
<i>cordialis</i> Morawitz, 1878	S	<i>limata</i> Smith, 1853	M S
<i>curvana</i> Warncke, 1965	M	<i>limbata</i> Eversmann, 1852	S
<i>curvungula</i> Thomson, 1870	B M S	<i>marginata</i> (Fabricius, 1776)	B M S
<i>danuvia</i> Stöckhert, 1950	B M S	<i>mehelyi</i> Alfken, 1936	M S
<i>decipiens</i> Schenck, 1859	B M S	<i>minutula</i> (Kirby, 1802)	B M S
<i>denticulata</i> (Kirby, 1802)	B M S	<i>minutuloides</i> Perkins, 1914	B M S
<i>dorsalis</i> Brullé, 1832	S	<i>mitis</i> Schmiedeknecht, 1883	B M S
<i>dorsata</i> (Kirby, 1802)	B M S	<i>mocsaryi</i> Schmiedeknecht, 1883	M S
<i>enslinella</i> Stöckhert, 1924	M S	<i>morawitzi</i> Thomson, 1872	B M S
<i>falsifica</i> Perkins, 1914	B M S	<i>morio</i> Brullé, 1832	B M S
<i>ferox</i> Smith, 1847	B M S	<i>mucida</i> Kriechbaumer, 1873	M S
<i>figurata</i> Morawitz, 1866	S	<i>nana</i> (Kirby, 1802)	B M S
<i>flavipes</i> Panzer, 1798	B M S	<i>nanaeformis</i> Noskiewitz, 1924	B
<i>floreo</i> Fabricius, 1793	B M S	<i>nanula subnuda</i> Alfken, 1924	B M S
<i>floricola</i> Eversmann, 1852	B M S	<i>nasuta</i> Giraud, 1863	B M S
<i>florivaga</i> Eversmann, 1852	B M S	<i>nigriceps</i> (Kirby, 1802)	B M S
<i>fucata</i> Smith, 1847	B M S	<i>nigroaenea</i> (Kirby, 1802)	B M S
<i>fulva</i> Müller, 1776	B	<i>nigrospina</i> Thomson, 1872	B M S
<i>armata</i> (Gmelin in Linnaeus, 1790)		<i>nitida</i> (Müller, 1776)	B M S
<i>fulvago</i> (Christ, 1791)	B M S		
<i>fulvata</i> Stöckhert, 1930	S		

<i>nitidiuscula</i> Schenck, 1853	B M S	
<i>niveata</i> Friese, 1887	B M S	
<i>nobilis</i> Morawitz, 1873	B M S	
<i>numida hypoplia</i>		
Schmiedeknecht, 1883	M S	
<i>nuptialis</i> Pérez, 1902	B M S	
<i>nycthemera</i> Imhoff, 1866	M S	
<i>obsoleta spongiosa</i>		
Warncke, 1967	B M S	
<i>distinguenda</i> Schenck, 1871		
<i>ocreata</i> (Christ, 1791)	B M S	
<i>oralis</i> Morawitz, 1876	M S	
<i>ovatula</i> (Kirby, 1802)	B M S	
<i>pallitarsis</i> Pérez, 1902	B M S	
<i>pandellei</i> Saunders, 1881	B M S	
<i>parviceps</i> Kriechbaumer, 1873	S	
<i>paucisquama</i> Noskiewicz, 1924	M S	
<i>pauxilla</i> Stöckhert, 1935	B M S	
<i>pillichi</i> Noskiewicz, 1937	B M S	
<i>polita</i> Smith, 1847	B M S	
<i>potentillae</i> Panzer, 1809	B M S	
<i>praecox</i> (Scopoli, 1763)	B M S	
<i>propinqua</i> Schenck, 1853	B M S	
<i>proxima</i> (Kirby, 1802)	B M S	
<i>pusilla</i> Pérez, 1902	B M S	
<i>ratisbonensis</i> Stöckhert, 1924	B M S	
<i>rosae</i> Panzer, 1800	B M S	
<i>roseipes</i> Alfken, 1933	S	
<i>ruficrus</i> (Nylander, 1848)	B M S	
<i>rufizona</i> Imhoff, 1834	B S	
<i>rufula</i> Schmiedeknecht, 1883	S	
<i>rugulosa</i> Stöckhert, 1935	B	
<i>sabulosa</i> (Scopoli, 1763)	B M S	
<i>carantenica</i> Pérez, 1902		
<i>semilaevis</i> Pérez, 1903	B M S	
<i>saundersella</i> Perkins, 1914		
<i>saxonica</i> Stöckhert, 1935	B M S	
<i>schencki</i> Morawitz, 1896	B M S	
<i>schlettereri</i> Friese, 1896	S	
<i>scita</i> Eversmann, 1852	B M S	
<i>seminuda</i> Friese, 1896	S	
<i>sericata</i> Imhoff, 1866	B M S	
<i>simillima</i> Smith, 1851	B M	
<i>spectabilis</i> Smith, 1853	S	
<i>strohmella</i> Stöckhert, 1930	B M S	
<i>subopaca</i> Nylander, 1848	B M S	
<i>suerinensis</i> Friese, 1894	B M S	
<i>susterai</i> Alfken, 1914	B M S	
<i>sympyti</i> Schmiedeknecht, 1883	B M S	
<i>synadelpha</i> Perkins, 1914	B S	
<i>taraxaci</i> Giraud, 1861	M S	
<i>tarsata</i> Nylander, 1848	B	
<i>thoracica</i> (Fabricius, 1775)	B M S	
<i>tibialis tibialis</i> (Kirby, 1802)	B M S	
<i>tibialis vindobonensis</i>		
Stöckhert, 1950	B M S	
<i>transitoria</i> Morawitz, 1872	M S	
<i>trimmerana</i> (Kirby, 1802)	M S	
<i>truncatilabris</i> Morawitz, 1878	M S	
<i>tscheki</i> Morawitz, 1872	B M S	
<i>vaga</i> Panzer, 1799	B M S	
<i>variabilis</i> Smith, 1853	S	
<i>varians</i> (Kirby, 1802)	B M S	
<i>ventralis</i> Imhoff, 1832	B M S	
<i>ventricosa</i> Dours, 1873	S	
<i>viridescens</i> Viereck, 1916	B M S	
<i>wolkella</i> (Kirby, 1802)	B M S	
<b>Melitturga</b> Latreille, 1809		
<i>clavicornis</i> (Latreille, 1806)	B M S	
<i>praestans</i> Giraud, 1861	M S	
<b>Panurgus</b> Panzer, 1806		
<b>Panurgus</b> s. str.		
<i>banksianus</i> (Kirby, 1802)	B M S	
<i>calcaratus</i> (Scopoli, 1763)	B M S	
<i>Panurginus</i> Nylander, 1848		
<i>labiatus</i> (Eversmann, 1852)	B M S	
<b>Camptopoeum</b> Spinola, 1843		
<i>friesei</i> (Mocsáry, 1894)	S	
<i>frontale</i> (Fabricius, 1804)	B M S	
HALICTIDAE		
<b>Halictus</b> Latreille, 1804		
<i>aeneidorsum</i> Alfken, 1920	M S	
<i>aeratus</i> (Kirby, 1802)	B M S	
<i>viridiaeneus</i> Blüthgen, 1918		
<i>albipes</i> (Fabricius, 1781)	B M S	
<i>albomaculatus</i> Lucas, 1846	?	
<i>alpinus</i> Alfken, 1907	?	

<i>bluethgeni</i> Ebmer, 1971	M	<i>marginellus</i> (Schenck, 1853)	B M S
<i>synhirtiventris</i> Blüthgen, 1930		<i>mesosclerus</i> Pérez, 1903	M S
<i>brevicornis</i> Schenck, 1870	B M S	<i>minutissimus</i> (Kirby, 1802)	B M S
<i>breviventris</i> (Schenck, 1853)	B M S	<i>minutulus</i> (Schenck, 1853)	B M S
<i>buccalis</i> Pérez, 1903	B M S	<i>minutus</i> (Kirby, 1802)	B M
<i>calceatus</i> (Scopoli, 1763)	B M S	<i>morbillosus</i> Kriechbaumer, 1873	M S
<i>clypearis</i> (Schenck, 1853)	B M S	<i>morio</i> (Fabricius, 1793)	B M S
<i>convexusculus</i> (Schenck, 1853)	M S	<i>niger</i> Viereck, 1883	B M S
<i>continentalis</i> Blüthgen, 1944	B M S	<i>fratellus</i> , Pérez, 1903	
<i>costulatus</i> Kriechbaumer, 1853	B M S	<i>nigripes</i> Lepeletier, 1841	B M S
<i>damascenus</i> Pérez, 1910	S	<i>nitidiusculus</i> (Kirby, 1802)	B M S
<i>distinctus</i> Schenck, 1868	B M S	<i>obscuratus</i> Morawitz, 1876	S
<i>elegans</i> Lepeletier, 1841	S	<i>pallens</i> Brullé, 1832	M S
<i>euboensis</i> Strand, 1909	M S	<i>patellatus</i> Morawitz, 1873	B S
<i>eurygnathus</i> Blüthgen, 1930	B M S	<i>pauxillus</i> (Schenck, 1853)	B M S
<i>fasciatus</i> Nylander, 1848	B M S	<i>perkinsi</i> Blüthgen, 1926	B M S
<i>fulvicornis</i> (Kirby, 1802)	B M S	<i>pleuralis</i> Moravitz, 1872	?
<i>glabriusculus</i> Morawitz, 1872	B M S	<i>pollinosus</i> Sichel, 1860	B M S
<i>griseolus</i> Morawitz, 1872	M	<i>prasinus</i> Smith, 1848	B M
<i>intermedius</i> Schenck, 1867—68	M	<i>punctatissimus</i> (Schenck, 1859)	B M S
<i>interruptus</i> (Panzer, 1798)	B M S	<i>puncticollis</i> Morawitz, 1872	B M S
<i>kessleri</i> Brams, 1879	M S	<i>pygmaeus</i> (Fabricius, 1804)	B M S
<i>laevigatus</i> (Kirby, 1802)	B M S	<i>quadricinctus</i> (Fabricius, 1774)	B M S
<i>laevis</i> (Kirby, 1802)	B M S	<i>quadrinotatus</i> (Schenck, 1859)	B M S
<i>langobardicus</i> Blüthgen, 1944	B M S	<i>quadrinotatus</i> (Kirby, 1802)	B M S
<i>laticeps</i> Schenck, 1867—68	B M S	<i>quadresignatus</i> (Schenck, 1853)	B M S
<i>lativentris</i> (Schenck, 1853)	B M S	<i>rubicundus</i> Christ, 1791	B M S
<i>leucopus</i> (Kirby, 1802)	B M S	<i>rufitarsis</i> Zetterstedt, 1938	B M
<i>leucozonius</i> (Schrank, 1781)	B M S	<i>sajoi</i> Blüthgen, 1923	M S
<i>limbellus</i> Morawitz, 1876	B M S	<i>scabiosae</i> (Rossi, 1790)	S
<i>linearis</i> Schenck, 1867—68	B M S	<i>seladonius</i> (Fabricius, 1794)	M S
<i>lucidulus</i> (Schenck, 1861)	B M S	<i>geminatus</i> Pérez, 1903	
<i>maculatus</i> Smith, 1848	B M S	<i>semitectus</i> Morawitz, 1873	M S
<i>major</i> Nylander, 1852	B M S	<i>senex</i> (Foerster, 1860)	B M S
<i>malachurus</i> (Kirby, 1802)	B M S	<i>setulosus</i> Strand, 1909	M S
<i>marginatus</i> (Brullé, 1832)	M S	<i>sexcinctus</i> (Fabricius, 1775)	B M S
		<i>sexmaculatus</i> (Schenck, 1853)	M

<i>sexnotatus</i> Nylander,				
1852	?			
<i>sexnotatus</i> (Kirby, 1802)	B M S			
<i>nitidus</i> (Panzer, 1798)				
<i>sexstrigatus</i> Schenck,				
1867—68	B M			
<i>simplex</i> Blüthgen, 1923	B M S			
<i>smeathmanellus nitidulus</i>				
(Fabricius, 1804)	B M S			
<i>subauratus</i> (Rossi, 1792)	B M S			
<i>subfasciatus</i> (Imhoff,				
1832)	B M S			
<i>tarsatus</i> Schenck,				
1867—68	B M			
<i>trichopygus</i> Blüthgen, 1923	M S			
<i>tricinctus</i> Schenck, 1874	B M S			
<i>truncaticollis</i> Morawitz,				
1878	M S			
<i>tumulorum</i> (Linnaeus,				
1758)	B M S			
<i>vestitus</i> Lepeletier, 1841	M S			
<i>villosulus</i> Kirby, 1802	B M S			
<i>xanthopus</i> (Kirby, 1802)	B M S			
<i>zonulus</i> Smith, 1848	B M S			
<b>Sphecodes</b> Latreille, 1805				
<i>albilabris</i> (Kirby, 1802)	B M S			
<i>alternatus</i> Smith, 1853	S			
<i>cephalotes</i> Meyer, 1922	M S			
<i>crassus</i> Thomson, 1870	B M S			
<i>cristatus</i> v. Hagens, 1882	M S			
<i>croaticus</i> Meyer, 1922	B M			
<i>ephippius</i> (Linnaeus, 1767)	B M S			
<i>fasciatus</i> v. Hagens, 1882	B M S			
<i>ferruginatus</i> v. Hagens,				
1882	B M S			
<i>gibbus</i> (Linnaeus, 1758)	B M S			
<i>pinguiculus hungaricus</i>				
Blüthgen, 1923	S			
<i>hyalinatus</i> v. Hagens, 1882	M S			
<i>intermedius</i> Blüthgen, 1923	M S			
<i>longulus</i> v. Hagens, 1882	B M S			
<i>majalis</i> Pérez, 1903	M			
<i>marginatus</i> v. Hagens, 1882	M S			
<i>miniatus</i> v. Hagens, 1882	B M S			
<i>monilicornis</i> (Kirby,				
1802)	B M S			
<i>niger</i> v. Hagens, 1882	B M S			
<i>pellucidus</i> Smith, 1845	B M S			
<i>pseudofasciatus</i> Blüthgen,				
1924	M S			
<i>puncticeps</i> Thomson, 1870	B M			
<i>reticulatus</i> Thomson, 1870	B M S			
<i>ruficrus rubicundus</i>				
v. Hagens, 1882	B M S			
<i>ruficentris</i> (Panzer, 1798)	B M S			
<i>subovalis</i> Schenck, 1866				
<i>scabricollis</i> Wesmael, 1835	B M S			
<i>schencki</i> v. Hagens, 1822	S			
<i>spinulosus</i> v. Hagens, 1875	B M S			
<b>Nomiooides</b> Schenck, 1866				
<i>minutissima</i> (Rossi, 1790)	B M S			
<i>variegata</i> (Olivier, 1789)	M S			
<b>Nomia</b> Latreille, 1804				
<i>diversipes</i> Latreille, 1806	S			
<i>femoralis</i> (Pallas, 1773)	M S			
<b>Dufourea</b> Lepeletier, 1841				
<i>dentiventris</i> (Nylander,				
1848)	B M S			
<i>halictula</i> (Nylander, 1852)	B M			
<i>inermis</i> (Nylander, 1848)	B M S			
<i>vulgaris</i> Schenck, 1895	B M S			
<b>Rhophites</b> Spinola, 1808				
<i>algirus trispinosus</i>				
Pérez, 1902	B M S			
<i>hartmani</i> Friese, 1902	B M S			
<i>quinquespinosus</i> Spinola,				
1808	1808			
<b>Rhophitoides</b> Schenck, 1859				
<i>canus</i> (Eversmann, 1852)	B M S			
<b>Systropha</b> Illiger, 1806				
<i>curvicornis</i> (Scopoli,				
1770)	B M S			
<i>planidens</i> Giraud, 1861	B M S			
<b>MELITTIDAE</b>				
<b>Melitta</b> Kirby, 1802				
<i>dimidiata</i> Morawitz, 1876	M S			
<i>haemorrhoidalis</i> (Fabricius,				
1775)	B M S			
<i>leporina</i> (Panzer, 1799)	B M S			
<i>melanura</i> Nylander, 1852	?			
<i>nigricans</i> Alfken, 1905	B M S			
<i>tricincta</i> Kirby, 1802	B M S			
<i>tomentosa</i> Friese, 1900	S			
<b>Dasypoda</b> Latreille, 1802				
<i>argentata</i> Panzer, 1809	B M S			

<i>hirtipes</i> (Fabricius, 1793)	B M S	
<i>mixta</i> Radoszkowski, 1886	B M	
<b>Macropis</b> Klug, in Panzer, 1809		
<i>friwaldskyi</i> Mocsáry, 1878	S	
<i>fulvipes</i> (Fabricius, 1804)	B M S	
<i>labiata</i> (Fabricius, 1804)	B M S	
<b>MEGACHILIDAE</b>		
<b>Lithurge</b> Latreille, 1825		
<i>chrysurus</i> Fonscolombe, 1834	S	
<i>cornutus</i> (Fabricius, 1787)	M S	
<b>Trachusa</b> Panzer, 1805		
<i>byssina</i> (Panzer, 1798)	B M S	
<b>Icteranthidium</b> Michener, 1948		
<i>laterale</i> (Latreille, 1809)	S	
<b>Paraanthidiellum</b> Michener, 1948		
<i>lituratum</i> (Panzer, 1801)	B M S	
<i>tenellum</i> (Mocsáry, 1879)	S	
<b>Anthidium</b> Fabricius, 1804		
<i>cingulatum</i> Latreille, 1809	B S	
<i>florentinum</i> (Fabricius, 1775)	S	
<i>loti</i> Perris, 1852	S	
<i>variegatum</i> (Fabricius, 1781)		
<i>manicatum</i> (Linnaeus, 1758)	B M S	
<i>montanum</i> Morawitz, 1864	B M S	
<i>oblongatum</i> Latreille, 1809	B M S	
<i>punctatum</i> Latreille, 1809	B M S	
<i>septemspinosum</i> Lepeletier, 1841	S	
<i>tornense</i> (Tkalcú, 1966)	S	
<b>Anthidiellum</b> Cockerell, 1904		
<i>strigatum</i> (Panzer, 1805)	B M S	
<b>Paraanthidium</b> Friese, 1898		
<i>interruptum</i> (Fabricius, 1781)	S	
<b>Rhodanthidium</b> Isensee, 1927		
<i>septemdentatum</i> (Latreille, 1809)	B M S	
<b>Stelis</b> Panzer, 1806		
<i>breviuscula</i> (Klug, 1807)	B M S	
<i>franconica</i> Blüthgen, 1930	7	
<i>minuta</i> Lepeletier et Serville, 1825	B M S	
<i>minima</i> Schenck, 1859		
<i>nasuta</i> (Latreille, 1809)	B M S	
<i>odontopyga</i> Noskiewitz, 1925	B M S	
<i>ornatula</i> (Klug, 1807)	B M S	
<i>phaeoptera</i> (Kirby, 1802)	B M S	
<i>punctulatissima</i> (Kirby, 1802)	B M S	
<i>moravica</i> Tkalcú, 1971		
<i>signata</i> (Latreille, 1809)	B M S	
<b>Dioxys</b> Lepeletier et Serville, 1825		
<b>Dioxoides</b> Popov, 1947		
<i>tridentata</i> (Nylander, 1848)	B M S	
<b>Dioxys</b> Lepeletier et Serville, 1825		
<i>cincta</i> (Jurine, 1807)	M S	
<b>Paradioxys</b> Mocsáry, 1877		
<i>pannonica</i> Mocsáry, 1877	S	
<b>Heriades</b> Spinola, 1808		
<i>crenulatus</i> Nylander, 1856	B M S	
<i>truncorum</i> (Linnaeus, 1758)	B M S	
<b>Chelostoma</b> Latreille, 1809		
<i>campanularum</i> (Kirby, 1802)	B M S	
<i>distinctum</i> Stöckhert, 1929	B M S	
<i>emarginatum</i> Nylander, 1856	M	
<i>florisomme</i> (Linnaeus, 1758)	B M S	
<i>foveolatum</i> (Morawitz, 1868)	S	
<i>fuliginosum</i> (Panzer, 1798)	B M S	
<i>grande</i> Nylander, 1852	?	
<i>handlirschi</i> Schletterer, 1889	S	
<i>ventrale</i> Schletterer, 1889	M S	
<b>Osmia</b> Panzer, 1806		
<i>aurulenta</i> (Panzer, 1799)	B M S	
<i>bicolor</i> (Schrank, 1781)	B M S	

<i>cerinthidis</i> Morawitz, 1876	B	M	S	<i>andrenoides</i> Spinola, 1808	M	S	
<i>coeruleescens</i> (Linnaeus, 1758)	B	M	S	<i>bidentata</i> Morawitz, 1876	B	M	S
<i>cornuta</i> (Latreille, 1805)	B	M	S	<i>ligurica</i> Morawitz, 1868		S	
<i>fulviventris</i> (Panzer, 1798)	B	M	S	<i>mocsaryi</i> Friese, 1895		S	
<i>gallarum</i> Spinola, 1808	B	M	S	<i>papaveris</i> (Latreille, 1799)	B	M	S
<i>inermis</i> (Zetterstedt, 1838)	B	M		<i>scutellaris</i> Morawitz, 1868		S	
<i>labialis tornensis</i> Tkalcù, 1975		S		<i>spinulosa</i> (Kirby, 1802)	B	M	S
<i>leaina</i> Kirby, 1802	B	M	S	<i>tergestensis</i> Ducke, 1897		M	S
<i>maritima</i> Friese, 1885		S		<i>villosa</i> (Schenck, 1853)	B	M	S
<i>melanogaster</i> Spinola, 1808		M	S				
<i>mustelina</i> Gerstaecker, 1869	B	M	S	<b>Metellinella</b> Tkalcù, 1966			
<i>nigriventris</i> Zetterstedt, 1838	B	S		<i>brevicornis</i> (Fabricius, 1798)	B	M	S
<i>parietina</i> Curtis, 1828	B	M	S				
<i>pilicornis</i> Smith, 1846	B	M	S	<b>Chalicodoma</b> Lepeletier, 1841			
<i>rufa</i> (Linnaeus, 1758)	B	M	S	<i>hungarica</i> Mocsáry, 1877		S	
<i>rufohirta</i> Latreille, 1811	B	M	S	<i>parietum</i> (Geoffroy in Fourcroy, 1785)	B	M	S
<i>submicans</i> Morawitz, 1871	M	S					
<i>uncinata</i> Gerstaecker, 1869	B	M		<b>Megachile</b> Latreille, 1802			
<i>ventralis</i> (Panzer, 1798)	B	M	S	<i>albisecta</i> (Klug, 1817)	M	S	
<i>xanthomelaena</i> (Kirby, 1802)	B	M		<i>alpicola</i> Alfken, 1924	B	S	
<b>Hoplitis</b> Klug, 1807				<i>analis</i> Nylander, 1852	B	M	S
<i>acuticornis</i> Dufour				<i>apicalis</i> Spinola, 1808	B	M	S
et Perris, 1840	B	M	S	<i>argentata</i> (Fabricius, 1793)	B	M	S
<i>adunca</i> (Panzer, 1798)	B	M	S	<i>bombycina</i> Radoszkowski, 1876		S	
<i>anthocopoides</i> Schenck, 1853	B	M	S	<i>centuncularis</i> (Linnaeus, 1758)	B	M	S
<i>laevifrons</i> Morawitz, 1872	M	S		<i>circumcincta</i> (Kirby, 1802)	B	M	S
<i>lepeletieri</i> Pérez, 1879	?			<i>deceptoria</i> Pérez, 1890	M	S	
<i>leucomelaena</i> (Kirby, 1802)	B	M	S	<i>ericetorum</i> Lepeletier, 1841	B	M	S
<i>manicata</i> Morice, 1901	M	S		<i>genalis</i> Morawitz, 1880	B	S	
<i>mitis</i> Nylander, 1852	B	M	S	<i>lagopoda</i> (Linnaeus, 1761)	B	M	S
<i>parvula</i> Doufour	B	M	S	<i>lapponica</i> Thomson, 1872	B	M	S
et Perris, 1840	S			<i>leucomalla</i> Gerstseker, 1869	M	S	
<i>praestans</i> Morawitz, 1894	B	M	S	<i>ligniseca</i> (Kirby, 1802)	B	M	S
<i>ravouxi</i> Pérez, 1902	B	M	S	<i>maackii</i> Radoszkowski, 1874	S		
<i>tenuispina</i> Alfken, 1937	S			<i>maritima</i> (Kirby, 1802)	B	M	S
<i>tridentata</i> Doufour	B	M	S	<i>melanopyga</i> Costa, 1863	B	M	S
et Perris, 1840	B	M	S	<i>nigriventris</i> Schenck, 1867—68	B		
<b>Anthocopa</b> Lepeletier et Serville, 1825				<i>octosignata</i> Nylander, 1852	M	S	
				<i>pilicrus</i> Morawitz, 1878	S		
				<i>pilidens</i> Alfken, 1923	M	S	
				<i>pyrenaea</i> Pérez, 1890	B	M	
				<i>rotundata</i> (Fabricius, 1784)	B	M	

<i>rubrimana</i> Morawitz, 1894	S	<i>braunsiana</i> Schmiedeknecht, 1882	B M
<i>versicolor</i> Smith, 1844	B M	<i>bouceki</i> Kocourek, 1985	S
<i>willughbiella</i> (Kirby, 1802)	B M S	<i>calimorpha</i> Schmiedeknecht, 1882	M S
<b>Coelioxys</b> Latreille, 1809		<i>castellana</i> Dusmet, 1913	B M S
<i>africana</i> Lepeletier, 1841	B M S	<i>baeri</i> Stöckhert, 1930	
<i>alata</i> Foerster, 1853	M	<i>chrysopyga</i> Morawitz, 1872	M S
<i>aurolimbata</i> Foerster, 1853	B M S	<i>cinctiventris</i> Friese, 1921	B M S
<i>brevis</i> Eversmann, 1852	B M S	<i>conjugens</i> Herrich et Schaeffer, 1839	B M S
<i>caudata</i> Spinola, 1938	S	<i>connectens</i> Pérez, 1884	S
<i>conoidea</i> (Illiger, 1806)	B M S	<i>corycraea</i> Schmiedeknecht, 1882	S
<i>elongata</i> Lepeletier, 1841	B M S	<i>cruenta</i> Smiedeknecht, 1882	M S
<i>emarginata</i> Foerster, 1853	B M S	<i>discrepans</i> Schmiedeknecht, 1882	S
<i>foersteri</i> Morawitz, 1872	S	<i>distinguenda</i> Morawitz, 1873	B M S
<i>haemorrhoa</i> Foerster, 1853	S	<i>dziesduszyckii</i> Noskiewicz, 1924	M S
<i>inermis</i> (Kirby, 1802)	B M S	<i>emarginata</i> Morawitz, 1878	B M
<i>lanceolata</i> (Nylander, 1852)	M	<i>errans</i> Lepeletier, 1841	B M S
<i>mandibularis</i> Nylander, 1848	B M S	<i>fabriciana</i> (Linnaeus, 1767)	B M S
<i>polycentris</i> Foerster, 1853	S	<i>facilis</i> Schwarz, 1967	B M S
<i>quadridentata</i> (Linnaeus, 1761)	B M S	<i>femoralis</i> Morawitz, 1869	B M S
<i>rufescens</i> Lepeletier, 1825	B M S	<i>ferruginata</i> (Linnaeus, 1767)	B M S
<i>rufocaudata</i> Smith, 1854	B M S	<i>xanthosticta</i> Kirby, 1802	
ANTHOPHORIDAE			
<b>Nomada</b> Scopoli, 1770		<i>flava</i> Panzer, 1798	B M S
<i>alboguttata</i> Herrich et Schaeffer, 1893	B M S	<i>flavoguttata</i> (Kirby, 1802)	B M S
<i>argentata</i> Herrich et Schaeffer, 1839	B M S	<i>flavopicta</i> (Kirby, 1802)	B M S
<i>armata</i> Herrich et Schaeffer, 1839	B M S	<i>fucata</i> Panzer, 1798	B M S
<i>atroscutellaris</i> Strand, 1921	M S	<i>fulvicornis</i> Fabricius, 1793	B M S
<i>baccata baccata</i> Smith, 1844	B M S	<i>fulvipes</i> Brullé, 1832	S
<i>palescens</i> Herrich et Schaeffer, 1839		<i>furva</i> Panzer, 1798	B M S
<i>baccata hrbantii</i> Balthasar, 1958	S	<i>furvoides</i> Stöckhert, 1943	M S
<i>basalis</i> Herrich et Schaeffer, 1839	M S	<i>fuscicornis</i> Nylander, 1848	B M S
<i>bifida</i> Thomson, 1872	B M S	<i>gabella</i> Thomson, 1870	B M
<i>bispinosa</i> Mocsáry, 1883	M S	<i>goodeniana</i> Kirby, 1802	B M S
<i>blepharipes</i> Schmiedeknecht, 1882	S	<i>guttulata</i> Schenck, 1859	B M S
<i>bluethgeni</i> Stöckhert, 1943	B M S	<i>hirtipes</i> Pérez, 1884	S
		<i>immaculata</i> Moravitz, 1873	?
		<i>incisa</i> Schmiedeknecht, 1882	S
		<i>insignipes</i> Schmiedeknecht, 1882	S

<i>integra</i> Brullé, 1832	S	<i>hillana</i> Kirby, 1802
<i>kohli</i> Schmiedeknecht, 1882	B M S	<i>sybarita</i> Schmiedeknecht, 1882
<i>lathburiana</i> (Kirby, 1802)	B M S	<i>syphityi</i> Stöckhert, 1930
<i>lepeletieri</i> Pérez, 1884	B M S	<i>tenella</i> Morcsáry, 1883
<i>leucophthalma</i> (Kirby, 1802)	B M S	<i>thersites</i> Schmiedeknecht, 1882
<i>lineola</i> Panzer, 1798	B M S	<i>tomentillae</i> Alfken, 1901
<i>marshamella</i> Kirby, 1802	B M S	<i>trapeziformis</i>
<i>melonopyga</i> Schmiedeknecht, 1882	S	Schmiedeknecht, 1882
<i>melathoracica</i> Imhoff, 1834	S	<i>trispinosa</i> Schmiedeknecht, 1882
<i>minuscula</i> Noskiewitz, 1930	M S	<i>verna</i> Schmiedeknecht, 1882
<i>mocsaryi</i> Schmiedeknecht, 1882	S	<i>villosa</i> Thomson, 1870
<i>montana</i> Mocsáry, 1894	B S	<i>zonata</i> Panzer, 1798
<i>mutabilis</i> Morawitz, 1871	B M S	
<i>mutica</i> Morawitz, 1872	B M S	
<i>nobilis</i> Herrich et Schaeffer, 1839	B M S	<b>Pasites</b> Jurine, 1807
<i>noskiewiczi</i> Schmiedeknecht, 1882	M S	<b>Parammobatodes</b> Popov, 1932
<i>nuptialis</i> Noskiewicz, 1930	S	<i>minutus</i> (Mocsáry, 1878)
<i>obscura</i> Zetterstedt, 1838	B M S	<b>Pasites</b> Jurine, 1807
<i>obtusifrons</i> Nylander, 1848	B M S	<i>maculatus</i> Jurine, 1807
<i>opaca</i> Alfken, 1913	B M	<b>Ammobates</b> Latreille, 1809
<i>panzeri</i> Lepeletier, 1858	B M S	<i>punctatus</i> (Fabricius, 1804)
<i>pectoralis</i> Morawitz, 1878	S	<i>similis</i> (Mocsáry, 1894)
<i>picciolana jurassica</i> Stöckhert, 1941	B M S	<i>vinctus</i> (Gerstaecker, 1869)
<i>pleurosticta</i> Herrich et Schaeffer, 1839	B	<b>Ammobatoides</b>
<i>pygidialis</i> Schwarz, 1968	S	Radoszkowski, 1868
<i>rhenana</i> Morawitz, 1872	B M S	<i>abdominalis</i> (Eversmann, 1852)
<i>roberjeotiana</i> Panzer, 1799	B M S	<b>Biastes</b> Panzer, 1806
<i>rostrata</i> Herrich et Schaeffer, 1839	M S	<i>brevicornis</i> (Panzer, 1798)
<i>eustalacta</i> Gerstaecker, 1869		<i>emarginatus</i> (Schenck, 1853)
<i>rufipes</i> Fabricius, 1793	B M S	<i>truncatus</i> (Nylander, 1848)
<i>schmiedeknechti</i> Schmiedeknecht, 1882	B M S	<b>Epeorus</b> Latreille, 1802
<i>sexfasciata</i> Panzer, 1799	B M S	<i>cruciger</i> (Panzer, 1799)
<i>shepardana</i> (Kirby, 1802)	B M S	<i>fasciatus</i> Friese, 1895
<i>signata</i> Jurine, 1807	B M S	<i>schummeli</i> Schilling, 1848
<i>similis</i> Morawitz, 1872	B M S	<i>tarsalis</i> Morawitz, 1873
<i>stigma</i> Fabricius, 1804 <i>cinnabarina</i> Morawitz, 1890	B M S	<i>variegatus</i> (Linnaeus, 1758)
<i>stoeckherti</i> Pittioni, 1947	S	<b>Triepeolus</b> Robertson, 1901
<i>striata</i> Fabricius, 1793	B M S	<i>tristis</i> (Smith, 1854)

<b>Epeoloides</b> Giraud, 1863			
<i>coecutiens</i> (Fabricius, 1775)	B M S		
<b>Tetralonia</b> Spinola, 1838			
<i>albicincta</i> (Lepeletier, 1841)	B M S		
<i>acutangula</i> Morawitz, 1878	S		
<i>dentata</i> (Klug, 1835)	B M S		
<i>fulvescens</i> Giraud, 1863	M S		
<i>graia</i> (Eversmann, 1852)	S		
<i>hungarica</i> (Friese, 1895)	B M S		
<i>inulae</i> Tkalcù, 1979	M S		
<i>lyncea</i> (Mocsáry, 1879)	S		
<i>macroglossa</i> Illyger, 1866	B M S		
<i>nana</i> Morawitz, 1873	M S		
<i>pollinosa</i> (Lepeletier, 1841)	S		
<i>ruficollis</i> (Brullé, 1832)	S		
<i>salicariae</i> (Lepeletier, 1841)	M S		
<i>scabiosae</i> Mocsáry, 1879	S		
<i>tricincta</i> (Erichson, 1835)	S		
<b>Eucera</b> Scopoli, 1770			
<i>caspica</i> Morawitz, 1873	M		
<i>chrysopyga</i> Pérez, 1879	S		
<i>cinerea</i> Lepeletier, 1841	M S		
<i>clypeata</i> Erichson, 1835	B M S		
<i>curvitarsis</i> Mocsáry, 1879	S		
<i>eucnemidea</i> Dours, 1873	S		
<i>hispana</i> Lepeletier, 1841	S		
<i>interrupta</i> Baer, 1850	B M S		
<i>longicornis</i> (Linnaeus, 1758)	B M S		
<i>nigrifacies</i> Lepeletier, 1841	S		
<i>nitidiventris</i> Mocsáry, 1879	S		
<i>parvicornis</i> Mocsáry, 1878	S		
<i>platyrrhina</i> Pérez, 1902	S		
<i>pollinosa</i> Smith, 1854	M S		
<i>seminuda</i> Brullé, 1832	M S		
<i>similis</i> Lepeletier, 1841	S		
<i>spectabilis</i> (Mocsáry, 1879)	S		
<i>taurica</i> Morawitz, 1871	S		
<i>tuberculata</i> (Fabricius, 1793)	B M S		
<b>Anthophora</b> Latreille, 1803			
<i>acerorum</i> (Linnaeus, 1758)	B M S		
<i>aestivalis</i> (Panzer, 1801)	B M S		
<i>borealis</i> Morawitz, 1864	B M S		
<i>crassipes</i> Lepeletier, 1841	M S		
<i>crinipes</i> Smith, 1854	B M S		
<i>parietina</i> (Fabricius, 1793)	M S		
<i>plagiata</i> (Illiger, 1806)			
<i>quadrimaculata</i> (Panzer, 1806)	B M S		
<i>retusa</i> (Linnaeus, 1758)	B M		
<b>Amegilla</b> Friese, 1897			
<i>albigena</i> Lepeletier, 1841	M S		
<i>garrula</i> (Rossi, 1790)	S		
<i>magnilabris</i> Fedtschenko, 1875	S		
<i>pipiens</i> Mocsáry, 1879	S		
<i>pubescens</i> (Fabricius, 1781)	B M S		
<i>quadrifasciata</i> (Villers, 1789)	B M S		
<i>salviae</i> Morawitz, 1876	S		
<b>Heliophila</b> Klug, 1807			
<i>bimaculata</i> (Panzer, 1798)	B M S		
<b>Clisodon</b> Patton, 1879			
<i>furcatus</i> (Panzer, 1798)	B M S		
<b>Melecta</b> Latreille, 1802			
<i>aegyptiaca</i> Radoszkowski, 1876	S		
<i>luctuosa</i> (Scopoli, 1770)	B M S		
<i>punctata</i> (Fabricius, 1775)	B M S		
<b>Thyreus</b> Panzer, 1806			
<i>affinis</i> (Morawitz, 1873)	S		
<i>histrionicus</i> (Illiger, 1806)	B M S		
<i>orbatus</i> Lepeletier, 1841	B M S		
<i>ramosus</i> (Lepeletier, 1841)	S		
<i>truncatus</i> (Pérez, 1883)	S		
<b>Ceratina</b> Latreille, 1802			
<i>acuta</i> Friese, 1896	S		
<i>chalybea</i> Chevrier, 1872	M S		
<i>cucurbitina</i> (Rossi, 1792)	M S		
<i>cyanea</i> (Kirby, 1802)	B M S		
<i>dentiventris</i> Gerstaecker, 1869	S		
<i>gravidula</i> Gerstaecker, 1869	S		
<i>laevifrons</i> Morawitz, 1895	S		

<i>nigrolabiata</i> Friese, 1896	S	<i>sylvarum</i> (Linnaeus, 1761)	B M S
<b>Xylocopa</b> Latreille, 1802		<i>soroensis</i> (Fabricius, 1777)	B M S
<i>cyanescens</i> Brullé, 1832	M S	<i>subterraneus</i> (Linnaeus, 1758)	B M S
<i>valga</i> Gerstaecker, 1872	B M S	<i>terrestris</i> (Linnaeus, 1758)	B M S
<i>violacea</i> (Linnaeus, 1758)	B M S	<i>wufleini</i> Radoszkowski, 1859	B M S
APIDAE			
<b>Bombus</b> Latreille, 1802		<b>Psithyrus</b> Lepeletier, 1833	
<i>agrorum</i> (Fabricius, 1787)	B M S	<i>barbutellus</i> (Kirby, 1802)	B M S
<i>argillaceus</i> (Scopoli, 1763)	S	<i>bohemicus</i> (Seidl, 1837)	B M S
<i>armeniacus</i> Boda, 1877	M	<i>campestris</i> (Panzer, 1801)	B M S
<i>confusus</i> Schenck, 1859	B M S	<i>maxillosus</i> (Klug, 1817)	M S
<i>distinguendus</i> Morawitz, 1869	B M S	<i>norvegicus</i> Sparre- -Schneider, 1918	B M S
<i>equestris</i> (Fabricius, 1783)	B M S	<i>quadricolor</i> Lepeletier, 1892	B M S
<i>veteranus</i> (Fabricius, 1793)		<i>rupestris</i> (Fabricius, 1793)	B M S
<i>elegans</i> (Fabricius, 1793)	B M S	<i>sylvestris</i> Lepeletier, 1833	B M S
<i>mesomelas</i> Gerstaecker, 1869		<i>vestalis</i> (Geoffroy in Fourcroy, 1785)	B M S
<i>fragrans</i> (Pallas, 1771)	M S		
<i>hortorum</i> (Linnaeus, 1761)	B M S	<b>Apis</b> Linnaeus, 1758	
<i>humilis</i> Illiger, 1806	B M S	<i>mellifera</i> Linnaeus, 1758	B M S
<i>hypnorum</i> (Linnaeus, 1756)	B M S	<i>mellifica</i> Linnaeus, 1761	
<i>jonellus</i> (Kirby, 1802)	B M S	hybrides of:	
<i>laesus</i> Morawitz, 1875	S	<i>mellifera carnica</i> Pollmann, 1879 introd.	
<i>lapidarius</i> (Linnaeus, 1758)	B M S	<i>mellifera cypria</i> Pollmann, 1879 introd.	
<i>lucorum</i> (Linnaeus, 1761)	B M S	<i>mellifera ligustica</i> Spinola, 1806 introd.	
<i>muscorum</i> (Linnaeus, 1758)	B M S	<i>mellifera mellifera</i> Linnaeus, 1758	
<i>pomorum</i> Panzer, 1805	B M S	<i>mellifera remipes</i> Gerstaecker, 1862 introd.	
<i>pratorum</i> (Linnaeus, 1751)	B M S		
<i>pyrenaeus</i> Pérez, 1879	S		
<i>ruderarius</i> (Müller, 1776)	B M S		
<i>ruderatus</i> (Fabricius, 1775)	M S		

## INDEX

to families and genera. Synonyms and other invalid names are not shown.

JOSEF ŠEDIVÝ

### FAMILIES

- Alloxystidae* — 130
- Ampulicidae* — 166
- Anacharitidae* — 130
- Andrenidae* — 174
- Anthophoridae* — 181
- Aphelinidae* — 118
- Aphidiidae* — 45
- Apidae* — 184
- Argidae* — 15
- Aspiceridae* — 130
- Astatidae* — 168
- Aulacidae* — 133
- Bethylidae* — 145
- Blasticotomidae* — 15
- Braconidae* — 27
- Cephidae* — 25
- Ceraphronidae* — 141
- Chalcididae* — 98
- Chrysidae* — 147
- Cimbicidae* — 15
- Cleptidae* — 146
- Colletidae* — 174
- Crabronidae* — 169
- Cynipidae* — 131
- Diplopidae* — 136
- Dipteronidae* — 16
- Dryinidae* — 144
- Elasmidae* — 119
- Embolemidae* — 145
- Encyrtidae* — 112
- Eucharitidae* — 102
- Eucolidae* — 130
- Eulophidae* — 119
- Eumenidae* — 157
- Eupelmidae* — 111
- Eurytomidae* — 99
- Evaniidae* — 133
- Fligittidae* — 130
- Formicidae* — 154
- Gasteruptiidae* — 134
- Haltictidae* — 176
- Heloridae* — 136
- Ibalidae* — 130
- Ichneumonidae* — 49
- Larridae* — 167
- Leucospidae* — 98
- Masariidae* — 157
- Megachilidae* — 179

- Megalodontidae* — 15
- Megaspilidae* — 141
- Melittidae* — 178
- Methochidae* — 150
- Mymaridae* — 127
- Mymarommatidae* — 127
- Mutillidae* — 150
- Nyssonidae* — 166
- Ormyridae* — 102
- Orussidae* — 25
- Pamphiliidae* — 14
- Pemphredontidae* — 168
- Perilampidae* — 102
- Philantidae* — 166
- Platygasteridae* — 140
- Pompilidae* — 161
- Proctotrupidae* — 136
- Pteromalidae* — 102
- Sapygidae* — 151
- Scelionidae* — 139
- Scoliidae* — 151
- Signiphoridae* — 118
- Siricidae* — 24
- Sphecidae* — 166
- Stephanidae* — 95
- Tenthredinidae* — 16
- Tetracampidae* — 119
- Tiphidae* — 150
- Torymidae* — 100
- Trichogrammatidae* — 126
- Trigonidae* — 142
- Trypoxylonidae* — 169
- Vespidae* — 159
- Xyelidae* — 14
- Xiphydriidae* — 24
- GENERAL**
- A**
- Abia* — 16
- Ablaxia* — 108
- Absyrta* — 70
- Acaenitus* — 84
- Acampsis* — 41
- Acantholyda* — 14
- Acanthopsilus* — 138
- Acephalonomia* — 146
- Aceratoneuromyta* — 124
- Achatus* — 93
- Achalcerinys* — 116

- Achrysocharoides* — 125
- Acidopria* — 139
- Aclista* — 138
- Aclastus* — 58
- Aclitus* — 45
- Aconias* — 63
- Acrisis* — 29
- Acrocormus* — 106
- Acrodactyla* — 51
- Acrolyta* — 58
- Acropiesia* — 137
- Acropimpla* — 51
- Acrotricnus* — 66
- Acrotomus* — 56
- Adelenyrtus* — 115
- Adelius* — 41
- Adelognathus* — 56
- Adelurola* — 34
- Ademon* — 32
- Adialytus* — 45
- Admirenyrtus* — 118
- Aethcerus* — 88
- Agasthenes* — 59
- Agathis* — 38
- Agentoleus* — 163
- Aggelma* — 108
- Aglaostigma* — 23
- Agonatopoides* — 144
- Agriotypus* — 57
- Agrogorytes* — 166
- Agromyzaphagus* — 118
- Agrothereutes* — 64
- Agrypon* — 84
- Alastor* — 158
- Alastorynerus* — 158
- Aleiodes* — 31
- Alexeter* — 71
- Allolus* — 36
- Allants* — 22
- Allocerastichus* — 124
- Allocerchysius* — 118
- Allodynerus* — 158
- Alloea* — 33
- Allogonatopus* — 144
- Allophroides* — 80
- Alloplasta* — 68
- Allotropa* — 141
- Alloxysta* — 130
- Allurus* — 43
- Allyson* — 167

- Alomya* — 87  
*Alysia* — 33  
*Amauronematus* — 20  
*Amblyaspis* — 141  
*Amblyjoppa* — 94  
*Amblyteles* — 93  
*Amegilla* — 183  
*Ametastegia* — 22  
*Amicroterys* — 113  
*Amitus* — 141  
*Ammeia* — 105  
*Ammobatoides* — 182  
*Amnophila* — 166  
*Amnoplatus* — 169  
*Ampulex* — 166  
*Amyras* — 35  
*Anacharis* — 130  
*Anagyrus* — 112  
*Anaphes* — 127  
*Anaprostocetus* — 123  
*Anastatus* — 111  
*Ancistrocerus* — 159  
*Andrena* — 174  
*Andricus* — 131  
*Anergates* — 154  
*Aneuclis* — 80  
*Aneugmenus* — 16  
*Aneuropria* — 139  
*Anisobas* — 90  
*Anisocyrtta* — 33  
*Anisopteromalus* — 106  
*Anisopygus* — 90  
*Anisotacrus* — 73  
*Anogmus* — 108  
*Anomalicornia* — 113  
*Anomalon* — 83  
*Anommatum* — 136  
*Anoncus* — 72  
*Anopedias* — 141  
*Anoplus* — 163  
*Anoplonyx* — 18  
*Anoplius* — 163  
*Anoxus* — 145  
*Anteon* — 145  
*Antepipona* — 158  
*Anteris* — 140  
*Anthemus* — 118  
*Anthidiellum* — 179  
*Anthidium* — 179  
*Anthocopa* — 180  
*Anthophora* — 183  
*Anustia* — 112  
*Aoplus* — 90  
*Apaeleticus* — 89  
*Apanteles* — 39  
*Apechthis* — 52  
*Apegus* — 140  
*Apeloma* — 108  
*Apethymus* — 23  
*Aphaenogaster* — 154  
*Aphaereta* — 33  
*Aphanistes* — 83  
*Aphanogmus* — 141  
*Aphelinidae* — 126  
*Aphelinus* — 118  
*Aphelonyx* — 131  
*Aphelopus* — 145  
*Aphycoidea* — 113  
*Aphyculus* — 114  
*Aphytus* — 114  
*Aphidius* — 45  
*Aphytis* — 119  
*Apis* — 184  
*Apophua* — 66  
*Aporinellus* — 163  
*Aporomyrmex* — 155  
*Aporus* — 162  
*Apotetrastrichus* — 122  
*Aprestes* — 137  
*Apronopa* — 35  
*Aprosthemata* — 15  
*Aprostocetus* — 122  
*Apsilocera* — 109  
*Apsilops* — 63  
*Aptesis* — 63  
*Aquaencyrtus* — 112  
*Arachnospila* — 163  
*Arachnotheutes* — 163  
*Archonomus* — 118  
*Archirileya* — 99  
*Ardis* — 21  
*Areneta* — 68  
*Areopraon* — 46  
*Arescon* — 127  
*Arge* — 15  
*Argogorytes* — 166  
*Aridelus* — 43  
*Aristelix* — 35  
*Aritratis* — 64  
*Arotes* — 84  
*Arotrephes* — 58  
*Arrhenophagus* — 117  
*Arthrolitus* — 110  
*Asaphes* — 103  
*Aschitus* — 113  
*Ascogaster* — 42  
*Asecodes* — 126  
*Asitus* — 112  
*Asobarra* — 34  
*Aspicolpus* — 36  
*Aspilota* — 34  
*Astata* — 168  
*Asthenolabus* — 89  
*Astichus* — 124  
*Astiphromma* — 81  
*Atanycolus* — 30  
*Atelopslus* — 136  
*Athalia* — 21  
*Atractodes* — 62  
*Atrichomalus* — 109  
*Atrometus* — 84  
*Aulacidea* — 132  
*Aulacus* — 133  
*Aulogymnus* — 120  
*Auplopus* — 162  
*Aylax* — 132  
*Azelus* — 71  
*Azotus* — 119  
**B**  
*Baeacis* — 36  
*Baeocharis* — 115  
*Baeognatha* — 39  
*Baeus* — 140  
*Bairamlia* — 103  
*Banchus* — 69  
*Barichneumon* — 91  
*Barycnemis* — 80  
*Baryconus* — 140  
*Barylypa* — 84  
*Baryproctus* — 30  
*Barytarbes* — 71  
*Basalys* — 138  
*Bathyplectes* — 75  
*Bathythrix* — 60  
*Batozonellus* — 164  
*Belaspida* — 98  
*Belomicrus* — 171  
*Belyta* — 137  
*Bembecinus* — 167  
*Bembix* — 167  
*Bethylus* — 145  
*Betuloxys* — 46  
*Blastes* — 182  
*Bioblapsis* — 87  
*Biolytia* — 75  
*Biorhiza* — 131  
*Blosteres* — 33  
*Birka* — 17  
*Blacometeorus* — 38  
*Blacus* — 37  
*Blasticotoma* — 15  
*Blastothrix* — 114  
*Blennocampa* — 21  
*Bocchus* — 144  
*Bombus* — 184  
*Bothriomyrmex* — 155  
*Boucekiella* — 116  
*Brachista* — 126  
*Brachycyrtus* — 56  
*Brachyelatus* — 103  
*Brachyencyrtus* — 113  
*Brachygaster* — 133  
*Brachymeria* — 98  
*Brachyserphus* — 136  
*Brachythops* — 17  
*Bracomorpha* — 30  
*Bracon* — 30  
*Bryopezus* — 120  
*Buathra* — 65  
*Bugacia* — 105  
**C**  
*Caenacis* — 107  
*Coenocrepis* — 106  
*Caenocryptus* — 65

- Caenolyda* — 14  
*Calameuta* — 25  
*Calaphidius* — 46  
*Calicurgus* — 162  
*Caltroa* — 20  
*Callajoppa* — 94  
*Callaspida* — 130  
*Callidora* — 75  
*Callimerismus* — 104  
*Callipteroma* — 112  
*Callitula* — 106  
*Calometropia* — 117  
*Calosota* — 111  
*Campocraspedon* — 86  
*Campodus* — 72  
*Compoletis* — 76  
*Camponotus* — 155  
*Campoplex* — 74  
*Campsoscolia* — 151  
*Capellia* — 107  
*Cardiochiles* — 41  
*Cardiopsilus* — 137  
*Casinaria* — 75  
*Catadelphus* — 94  
*Cataglyphis* — 155  
*Catalytus* — 59  
*Catastenus* — 85  
*Catolaccus* — 107  
*Cea* — 102  
*Ceballosia* — 118  
*Cecidostiba* — 107  
*Cecidotetrastichus* — 122  
*Celonites* — 157  
*Cenocoelius* — 36  
*Centeterus* — 88  
*Centistes* — 43  
*Centrobia* — 126  
*Centrodora* — 119  
*Cephalcia* — 14  
*Cephaloglypta* — 67  
*Cephalonomia* — 145  
*Cephaloplites* — 32  
*Cephus* — 25  
*Ceranisus* — 126  
*Ceraphron* — 141  
*Cerapterocerus* — 115  
*Ceratepyris* — 146  
*Ceratina* — 183  
*Cerceris* — 166  
*Cerchystella* — 116  
*Cerchysius* — 114  
*Cercobelus* — 115  
*Cerocephala* — 103  
*Ceropales* — 103  
*Ceroptres* — 132  
*Chaenon* — 35  
*Chaenusa* — 35  
*Chaetostricha* — 126  
*Chalcis* — 98  
*Challicdoma* — 180  
*Charitopes* — 60  
*Charitopus* — 112  
  
*Charmon* — 38  
*Charops* — 75  
*Chartocerus* — 118  
*Chasmias* — 93  
*Chasmodon* — 33  
*Cheiroleurus* — 116  
*Cheiropachus* — 106  
*Cheilogynus* — 144  
*Chelonus* — 42  
*Chelostoma* — 179  
*Chrotica* — 57  
*Chlorocytus* — 108  
*Choeras* — 40  
*Chorebus* — 35  
*Choreia* — 115  
*Chorinaeus* — 82  
*Chremylus* — 29  
*Chrysidea* — 148  
*Chrysis* — 148  
*Chrysocharis* — 125  
*Chrysolampus* — 103  
*Chrysomala* — 103  
*Chrysonotomyia* — 125  
*Chrysopophthorus* — 44  
*Cidaphus* — 81  
*Cimbex* — 16  
*Cinetus* — 138  
*Cinxaletus* — 88  
*Citrospilus* — 120  
*Cladardis* — 22  
*Cladius* — 17  
*Cleolophus* — 120  
*Cleonymus* — 102  
*Cleptes* — 146  
*Clinocentrus* — 32  
*Clisodon* — 183  
*Clistopyga* — 51  
*Closterocerus* — 126  
*Clytina* — 118  
*Coccobius* — 118  
*Coccophagoidea* — 118  
*Coccophagus* — 118  
*Codrus* — 136  
*Coelichneumon* — 94  
*Coelioxys* — 181  
*Coelinus* — 35  
*Coeloides* — 30  
*Coleopencyrtus* — 117  
*Coelopisthia* — 110  
*Coenocercus* — 116  
*Colastes* — 29  
*Coleocentrus* — 84  
*Colletes* — 174  
*Collyria* — 85  
*Colocnema* — 63  
*Coloneura* — 35  
*Colotrechnus* — 111  
*Colpoclypeus* — 121  
*Colpognathus* — 88  
*Colpotrochia* — 83  
*Comperiella* — 115  
*Conomorium* — 110  
  
*Conostigmus* — 142  
*Copidosoma* — 116  
*Copidosomopsis* — 117  
*Coruna* — 111  
*Corynis* — 16  
*Cosmoconus* — 55  
*Cosmophorus* — 43  
*Cotesia* — 40  
*Cothonaspis* — 130  
*Crabro* — 170  
*Crataeps* — 124  
*Cratichneumon* — 90  
*Cratocryptus* — 62  
*Cratomus* — 111  
*Cratospilla* — 34  
*Cremastus* — 79  
*Cremnodes* — 62  
*Cricellius* — 108  
*Croesus* — 19  
*Crossocerus* — 169  
*Crypteffigies* — 90  
*Cryptocheilus* — 161  
*Cryptopimpla* — 68  
*Cryptopristus* — 100  
*Cryptoprymna* — 104  
*Cryptoserphus* — 136  
*Ctenichneumon* — 96  
*Cteniscus* — 56  
*Ctenochira* — 54  
*Ctenopelma* — 69  
*Cubocephalus* — 62  
*Cuneala* — 24  
*Cyanopterus* — 30  
*Cycasis* — 55  
*Cyclogastrella* — 110  
*Cyclolabus* — 89  
*Cylloceria* — 85  
*Cymodusa* — 77  
*Cynips* — 131  
*Cyrtogaster* — 104  
*Cystomutilla* — 151  
  
 D  
*Dacnusa* — 35  
*Dahlbominus* — 121  
*Dapsilarthra* — 34  
*Dasylabris* — 151  
*Dasypoda* — 178  
*Decanematus* — 20  
*Delomerista* — 53  
*Delopia* — 76  
*Demopheles* — 62  
*Dendrocerus* — 142  
*Dendrosoter* — 28  
*Dentilabus* — 89  
*Dermatopeltis* — 120  
*Derostenus* — 125  
*Desmatochartis* — 126  
*Deuterixys* — 41  
*Deuteroxorides* — 53  
*Diachasma* — 33  
*Diadegma* — 78

- Diadromus* — 88  
*Diaeretiellus* — 46  
*Diaeretiella* — 46  
*Diaeretus* — 46  
*Diaglyptidea* — 58  
*Diaparsis* — 80  
*Diapria* — 139  
*Diaschiaspis* — 132  
*Diastrophus* — 132  
*Diaulinopssiss* — 120  
*Dibrachides* — 110  
*Dibrachys* — 110  
*Dicaelotus* — 87  
*Dicarnosis* — 120  
*Dichrogaster* — 58  
*Dicladocerus* — 120  
*Dicondylus* — 144  
*Dicyrtaspis* — 36  
*Dicyrtomellus* — 163  
*Didactyliocerus* — 101  
*Didineis* — 167  
*Diglochis* — 110  
*Diglyphosema* — 130  
*Diglyphus* — 121  
*Dimachus* — 110  
*Dimeromicrus* — 101  
*Dimmockia* — 120  
*Dimophora* — 79  
*Dinarmoides* — 106  
*Dinarmus* — 106  
*Dinetus* — 168  
*Dineura* — 18  
*Dinocampus* — 44  
*Dinocarnosis* — 112  
*Dinocarsiella* — 112  
*Dinocarsis* — 112  
*Dinotiscus* — 105  
*Dinotoides* — 107  
*Diodontus* — 169  
*Diolcogaster* — 41  
*Diomorus* — 101  
*Diospilus* — 36  
*Dioxys* — 179  
*Dipara* — 103  
*Diphora* — 137  
*Diphyus* — 93  
*Diplazon* — 86  
*Diplolepis* — 131  
*Diplorhoptrum* — 154  
*Dipogon* — 162  
*Dipriocampe* — 119  
*Dipton* — 16  
*Dirhincus* — 110  
*Dtrhinus* — 98  
*Dirophanes* — 88  
*Discodes* — 115  
*Discoelius* — 157  
*Disogmus* — 136  
*Disophrys* — 38  
*Disorygma* — 130  
*Distatrix* — 40  
*Dolerus* — 17  
*Dolichoderus* — 155  
*Dolichogenidea* — 39  
*Dolichomitus* — 51  
*Dolichurus* — 166  
*Doliphoceras* — 112  
*Dolophron* — 77  
*Dolopsidea* — 29  
*Doryctes* — 28  
*Doryctosoma* — 29  
*Dreisbachia* — 51  
*Drepanocontonus* — 83  
*Dryinus* — 144  
*Dujourea* — 178  
*Dusmetia* — 112  
*Dyscritulus* — 46  
*Dyscoletes* — 36  
*Dyspetes* — 55  
  
 E  
*Earinus* — 38  
*Earobia* — 74  
*Eclithroplexiella* — 115  
*Echthrus* — 63  
*Ecytus* — 54  
*Ecphyllus* — 29  
*Ecrizotes* — 105  
*Ectadius* — 141  
*Ectemnius* — 170  
*Ectopius* — 89  
*Ectroma* — 115  
*Elachertus* — 119  
*Elasmosoma* — 43  
*Elasmus* — 119  
*Elinora* — 23  
*Elysoceraphron* — 141  
*Embolemus* — 145  
*Empria* — 22  
*Encarsia* — 118  
*Enclisis* — 64  
*Enrateola* — 57  
*Encyrtoscelio* — 140  
*Encyrtus* — 117  
*Endasys* — 60  
*Endelomyia* — 20  
*Enicospilus* — 81  
*Enizemum* — 86  
*Entedon* — 124  
*Entomacris* — 138  
*Entomognathus* — 169  
*Enytus* — 78  
*Eoferreola* — 162  
*Eopsis* — 22  
*Eparces* — 88  
*Epeoloides* — 183  
*Epeolus* — 182  
*Ephedrus* — 46  
*Ephialtes* — 50  
*Epiclerus* — 119  
*Epmicta* — 35  
*Epimyrma* — 154  
*Epistathmus* — 80  
*Episyron* — 164  
  
*Epitetracnemus* — 115  
*Epitomus* — 88  
*Epyris* — 146  
*Erdoesia* — 104  
*Erdoesina* — 110  
*Eremotylus* — 81  
*Eretmocerus* — 119  
*Ericydinus* — 112  
*Eridolitus* — 56  
*Eridontomerus* — 100  
*Eriocampa* — 22  
*Eriplatys* — 88  
*Eripternus* — 76  
*Erromenus* — 55  
*Erythmelus* — 127  
*Erythromalus* — 107  
*Ethelurgus* — 60  
*Eubazus* — 36  
*Eucera* — 183  
*Euceros* — 56  
*Euchalcis* — 98  
*Eucharis* — 102  
*Euchroeus* — 148  
*Euccidophagus* — 118  
*Eucoila* — 130  
*Eudelus* — 58  
*Euderomphale* — 126  
*Euderus* — 124  
*Eugahania* — 117  
*Eugertium* — 126  
*Eulonchetron* — 108  
*Eulophus* — 121  
*Eumacepolus* — 108  
*Eumenes* — 159  
*Euneura* — 111  
*Eunotus* — 103  
*Euodynerus* — 159  
*Eupalamus* — 90  
*Eupelmus* — 111  
*Euplectrus* — 119  
*Eupoecilopoda* — 117  
*Eurydtnoia* — 107  
*Eurytschia* — 119  
*Eurylabus* — 89  
*Euryproctus* — 73  
*Eurytoma* — 99  
*Eusandalum* — 111  
*Eusemion* — 115  
*Eusterinx* — 85  
*Eustochus* — 127  
*Eutanyacra* — 93  
*Eutetrastrichus* — 121  
*Eutomostethus* — 21  
*Euura* — 20  
*Evagetes* — 163  
*Excavarus* — 55  
*Exenterus* — 56  
*Exeristes* — 50  
*Exephanes* — 93  
*Exetastes* — 68  
*Exochus* — 83  
*Exopristus* — 100

- Exotela* — 35  
*Exyston* — 55  
**F**  
*Fahringeria* — 141  
*Falciconus* — 46  
*Fenella* — 21  
*Fenusia* — 21  
*Ferreola* — 162  
*Fidlobia* — 141  
*Figites* — 130  
*Flavopimpla* — 50  
*Foersterella* — 119  
*Foersteria* — 36  
*Formica* — 156  
*Formicoxenus* — 154  
*Foveobracon* — 31  
**G**  
*Gambrus* — 64  
*Gasteruption* — 134  
*Gastracanthus* — 103  
*Gastracistrus* — 105  
*Gbelcia* — 109  
*Geltis* — 58  
*Gilpinia* — 16  
*Ginsiana* — 118  
*Graudia* — 63  
*Glenosema* — 146  
*Globulencyrtus* — 118  
*Glypticnemis* — 60  
*Glyphognathus* — 104  
*Glyphomerus* — 100  
*Glypta* — 66  
*Glyptapanteles* — 40  
*Glyptorhaestus* — 70  
*Gnaptodon* — 29  
*Gnotus* — 61  
*Goedartia* — 89  
*Gonatocerus* — 127  
*Gonatopus* — 144  
*Gontozus* — 145  
*Gonotyphus* — 76  
*Gorytes* — 166  
*Grahamia* — 125  
*Gravenhorstia* — 84  
*Gregoptimpla* — 51  
*Gryon* — 140  
*Grypocentrus* — 54  
*Gunomeria* — 73  
*Gymnomerus* — 158  
*Gyrinophagus* — 110  
**H**  
*Habritys* — 106  
*Habrobracon* — 31  
*Habroleptis* — 115  
*Habronyx* — 83  
*Hadrodactylus* — 73  
*Halictus* — 176  
*Halidamia* — 22  
*Halomalus* — 110  
*Haltichella* — 98  
*Halticoptera* — 104  
*Halticopterina* — 104  
*Harpagoxenus* — 154  
*Harpiphorus* — 22  
*Hartigia* — 25  
*Hecabolus* — 29  
*Hedychridium* — 148  
*Hedychrum* — 148  
*Helcon* — 36  
*Helconidea* — 36  
*Helcostizus* — 60  
*Helegonatopus* — 114  
*Helictes* — 85  
*Heliophila* — 183  
*Hellwigia* — 74  
*Helorus* — 136  
*Hemicneumon* — 87  
*Hemicroa* — 18  
*Hemiptarsenus* — 120  
*Hemipterochilus* — 158  
*Hemiteles* — 58  
*Hemitrichus* — 110  
*Hepiopelmus* — 93  
*Heptamelus* — 17  
*Hercus* — 54  
*Heriades* — 179  
*Herpestomus* — 88  
*Heterarthrus* — 20  
*Heterischnus* — 87  
*Heterocoelia* — 146  
*Heterocola* — 79  
*Heteropelma* — 83  
*Heteroprymna* — 109  
*Heterospilus* — 28  
*Heydenia* — 102  
*Himerta* — 72  
*Hinatara* — 21  
*Hispaniella* — 118  
*Histeromerus* — 29  
*Hobbya* — 107  
*Hockeria* — 98  
*Holcaeus* — 108  
*Holcopelte* — 126  
*Holcotetrastichus* — 122  
*Holepyris* — 146  
*Holopyga* — 148  
*Holoteleia* — 140  
*Homalotyloidea* — 117  
*Homalotylus* — 117  
*Homaspis* — 69  
*Homolobus* — 38  
*Homonotus* — 162  
*Homoporus* — 106  
*Homotrechus* — 90  
*Homotropus* — 86  
*Hoplismenus* — 94  
*Hoplitis* — 180  
*Hoplocampa* — 18  
*Hoplocampoides* — 21  
*Hoplopsis* — 113  
*Hormius* — 29  
*Hybophanes* — 54  
*Hybothorax* — 99  
*Hybrizon* — 44  
*Hydryta* — 64  
*Hypamblys* — 73  
*Hyperactmus* — 85  
*Hyperbatus* — 72  
*Hyperimerus* — 103  
*Hypomecus* — 89  
*Hypoponera* — 154  
*Hypsotter* — 78  
*Hypsantyx* — 73  
*Hypsicera* — 83  
**I**  
*Ibalia* — 130  
*Ichneumon* — 91  
*Ichneutes* — 43  
*Iconella* — 40  
*Icteranthidium* — 179  
*Idarnotorymus* — 100  
*Idiasta* — 34  
*Idiogramma* — 56  
*Idiolexis* — 34  
*Idiolispa* — 64  
*Idiotypa* — 138  
*Idris* — 140  
*Illidops* — 39  
*Inocerota* — 140  
*Inostemma* — 140  
*Invreia* — 99  
*Ionymptha* — 126  
*Iphiaulax* — 30  
*Iphitachelus* — 141  
*Ipideurytoma* — 100  
*Iseropus* — 51  
*Ischnoceros* — 57  
*Ischnus* — 64  
*Ischyroptyx* — 106  
*Ismarus* — 136  
*Isocolus* — 132  
*Isocyrtus* — 108  
*Isodromus* — 117  
*Isomecus* — 30  
*Isostasius* — 141  
*Israelius* — 145  
*Itamoplex* — 65  
*Itoplectis* — 52  
*Ixodiphagus* — 115  
**J**  
*Jansoniella* — 103  
*Janus* — 25  
*Javra* — 62  
*Jucancistrocerus* — 158  
**K**  
*Kaleva* — 108  
*Karpinskiella* — 108  
*Katamenes* — 159  
*Kleidotoma* — 130  
*Kocourekia* — 124

- Kratoysma* — 125  
*Kriechbaumerella* — 98  
*Kristotomus* — 55  
*Krombeinella* — 151  
*Ksenoplata* — 104  
**L**  
*Labolips* — 139  
*Laelius* — 146  
*Lagarotis* — 71  
*Lagynodes* — 142  
*Lamachus* — 71  
*Lamennaisia* — 118  
*Lampoterma* — 109  
*Lamprotatus* — 105  
*Laotris* — 35  
*Lapton* — 83  
*Lariophagus* — 110  
*Larra* — 167  
*Lasiochalcidita* — 99  
*Lasius* — 155  
*Lathiponus* — 70  
*Lathrolestes* — 71  
*Lathromeris* — 126  
*Lathroplex* — 74  
*Lathrostizus* — 79  
*Latibus* — 66  
*Leiocyrtus* — 116  
*Leiophron* — 44  
*Lemophagus* — 79  
*Leptacis* — 141  
*Leptacoenites* — 84  
*Leptocampoplex* — 76  
*Leptochilus* — 158  
*Leptomastix* — 112  
*Leptomastidea* — 112  
*Leptomerapus* — 109  
*Leptonetus* — 138  
*Leptoteleia* — 140  
*Leptothonax* — 154  
*Lestica* — 170  
*Lestrices* — 36  
*Lethades* — 69  
*Leucospis* — 98  
*Limerodes* — 94  
*Limerodops* — 93  
*Lindentus* — 169  
*Linycus* — 89  
*Liodontomerus* — 101  
*Liometopum* — 155  
*Liotryphon* — 50  
*Lipolexis* — 46  
*Liposthenes* — 132  
*Liris* — 167  
*Lissogaster* — 40  
*Lissonota* — 67  
*Listrocryptus* — 63  
*Listrodromus* — 90  
*Listrognathus* — 65  
*Lithurge* — 179  
*Litus* — 127  
*Loderus* — 17  
*Lonchetron* — 108  
*Lonchidia* — 130  
*Lophyrolectus* — 70  
*Loxocephalus* — 44  
*Lycorina* — 66  
*Lyka* — 112  
*Lymnatrichneumon* — 91  
*Lysaphidus* — 46  
*Lysephedrus* — 46  
*Lysibia* — 58  
*Lysiphlebus* — 46  
*Lysitermus* — 29  
**M**  
*Macrocentrus* — 38  
*Macrodiplion* — 16  
*Macromesus* — 102  
*Macroneura* — 111  
*Macrophya* — 24  
*Macropis* — 179  
*Macroteleia* — 140  
*Macrus* — 77  
*Mahencyrtus* — 116  
*Manica* — 154  
*Marietta* — 119  
*Mastrulus* — 60  
*Mastrus* — 59  
*Mayrencyrtus* — 116  
*Mayridia* — 115  
*Medophron* — 60  
*Megacara* — 61  
*Megachile* — 180  
*Megalodontes* — 15  
*Megaplectes* — 63  
*Megarhyssa* — 53  
*Megastigmus* — 100  
*Megastylus* — 85  
*Melanocistrus* — 105  
*Melanichneumon* — 91  
*Melanips* — 130  
*Melanopus* — 15  
*Melecta* — 183  
*Melitta* — 178  
*Melittobia* — 124  
*Melitturga* — 176  
*Mellinus* — 167  
*Meloboris* — 77  
*Merapori* — 109  
*Mercetia* — 144  
*Meria* — 150  
*Meringopus* — 65  
*Merismus* — 104  
*Merisus* — 106  
*Merostenus* — 112  
*Mesitius* — 146  
*Mesochorus* — 81  
*Mesoclistus* — 84  
*Mesodryinus* — 144  
*Mesoleius* — 72  
*Mesoleptidea* — 73  
*Mesoleptus* — 62  
*Mesoneura* — 18  
*Mesopolobus* — 108  
*Mesostenidea* — 64  
*Mesostenus* — 65  
*Messa* — 21  
*Messor* — 154  
*Mestocharis* — 125  
*Metablastothrix* — 114  
*Metaclysis* — 140  
*Metacolus* — 106  
*Metallus* — 21  
*Metapelma* — 111  
*Metaphaenodiscus* — 112  
*Metaphidius* — 47  
*Metaphycus* — 114  
*Metastenus* — 111  
*Metellinella* — 180  
*Meteorus* — 43  
*Methocha* — 150  
*Metopius* — 82  
*Micradelus* — 105  
*Microchelonus* — 42  
*Microctonus* — 44  
*Microdiprion* — 16  
*Microdontomerus* — 101  
*Microdynerus* — 158  
*Microgaster* — 41  
*Microleptes* — 85  
*Microlycus* — 121  
*Micrope* — 88  
*Microstilba* — 130  
*Microterys* — 113  
*Microtypus* — 113  
*Minotetrastichus* — 122  
*Miotropis* — 120  
*Mira* — 115  
*Mirax* — 41  
*Mirufens* — 126  
*Mischotetrastichus* — 122  
*Miscogaster* — 105  
*Miscophus* — 168  
*Misetus* — 87  
*Misocyclops* — 141  
*Mohegencyrtus* — 113  
*Mohejniella* — 113  
*Mokrzekia* — 107  
*Monardis* — 22  
*Monekaza* — 138  
*Monoblastus* — 55  
*Monocetus* — 16  
*Monoclonia* — 47  
*Monoctonus* — 47  
*Monodiscodes* — 112  
*Monodontomerus* — 100  
*Monolexis* — 29  
*Monomorium* — 154  
*Monophadnoides* — 22  
*Monophadnus* — 21  
*Monorthrochaeta* — 126  
*Monosoma* — 22  
*Monostegia* — 22  
*Moraviella* — 110  
*Muscidiifurax* — 110

- Mutilla* — 151  
*Mymar* — 127  
*Myrmecina* — 154  
*Myrmeleonostenus* — 65  
*Myrmica* — 154  
*Myrmilla* — 154  
*Myrmosa* 150  
*Mystrophorus* — 144  
  
**N**  
*Nanoclavelia* — 163  
*Nasonia* — 110  
*Necremnus* — 121  
*Negenaspidus* — 118  
*Nematinus* — 18  
*Nematopodius* — 66  
*Nematus* — 19  
*Nemeritis* — 75  
*Neoceraphron* — 141  
*Neochalcis* — 98  
*Neodipara* — 103  
*Neodiprion* — 16  
*Neogonatopus* — 144  
*Neohybothorax* — 99  
*Neoneurus* — 43  
*Neorhacodes* — 66  
*Neotrichoporoides* — 123  
*Neotypus* — 90  
*Neoxorides* — 53  
*Nepista* — 76  
*Nesolandria* — 17  
*Netelia* — 53  
*Netomocera* — 103  
*Neurateles* — 86  
*Neurocrassus* — 28  
*Neuroterus* — 131  
*Neurotoma* — 15  
*Nikanoria* — 100  
*Nippocryptus* — 65  
*Nitela* — 168  
*Nodisoplata* — 104  
*Nomada* — 181  
*Nomia* — 178  
*Nomioides* — 178  
*Norbanus* — 106  
*Noserus* — 29  
*Notanisus* — 102  
*Notoglyptus* — 104  
*Notopygus* — 69  
*Notosemus* — 88  
*Notozus* — 147  
*Novitzkianus* — 104  
*Nysson* — 167  
  
**O**  
*Obstphaga* — 58  
*Obtusodonta* — 93  
*Odontocolon* — 57  
*Odynerus* — 158  
*Oetophorus* — 70  
*Oleisticampe* — 78  
*Olethrodottis* — 69  
  
*Oligosita* — 126  
*Omalus* — 147  
*Omphale* — 126  
*Oncophanes* — 29  
*Ontsira* — 28  
*Ooctonus* — 127  
*Oeencyrtus* — 114  
*Oomyzus* — 122  
*Opazón* — 137  
*Opheltes* — 71  
*Opion* — 81  
*Ophioneurus* — 127  
*Opisthacantha* — 139  
*Opius* — 32  
*Oresbius* — 62  
*Orgichneumon* — 91  
*Orgilus* — 38  
*Ormocerus* — 105  
*Ormyrus* — 102  
*Oronotus* — 87  
*Orthizema* — 61  
*Orthocentrus* — 85  
*Orthopelma* — 85  
*Orthostigma* — 34  
*Orussus* — 25  
*Osmia* — 179  
*Otlophorus* — 71  
*Oxybelus* — 171  
*Oxylabis* — 137  
*Oxypria* — 139  
*Oxyrrhexis* — 52  
*Oxysuchus* — 106  
*Oxytorus* — 85  
  
**P**  
*Pachycrepoides* — 111  
*Pachynematus* — 18  
*Pachyneuron* — 111  
*Pachyprotasis* — 24  
*Palaeocimbex* — 16  
*Palaeomyimar* — 127  
*Palarus* — 168  
*Pambolus* — 29  
*Pamphilus* — 14  
*Pandelus* — 106  
*Pannontiella* — 102  
*Panstenon* — 103  
*Pantoclis* — 137  
*Pantolyta* — 137  
*Pantorhaestes* — 73  
*Panurgus* — 176  
*Paraanthidium* — 179  
*Paraanthidium* — 179  
*Parablastothrix* — 117  
*Parablatticida* — 118  
*Paracentrobia* — 127  
*Paracharactus* — 21  
*Paracodus* — 136  
*Paracoelichneumon* — 91  
*Paracopidosoma* — 113  
*Paraenastoidea* — 112  
*Paragymnomerus* — 158  
  
*Paralipsis* — 47  
*Paralitomastix* — 117  
*Paramasia* — 113  
*Paramesius* — 138  
*Paramonocetus* — 47  
*Paramyrmosa* — 151  
*Parania* — 84  
*Paraperithous* — 51  
*Parapraon* — 47  
*Parasauleia* — 118  
*Parascleroderma* — 146  
*Parasecodella* — 124  
*Parasierola* — 145  
*Paratetracnemoidea* — 112  
*Pareophora* — 22  
*Paraeucorystes* — 29  
*Parmortha* — 62  
*Parna* — 21  
*Paropae* — 148  
*Parodontodynerus* — 158  
*Paroxylabis* — 138  
*Parthenocodrus* — 136  
*Pasites* — 182  
*Passaloecus* — 169  
*Patrocloides* — 92  
*Patroclus* — 93  
*Pauesia* — 47  
*Pediaspis* — 132  
*Pediothius* — 124  
*Pegopus* — 109  
*Pempredon* — 168  
*Pentapleura* — 35  
*Periclista* — 22  
*Periclistus* — 132  
*Peridesmia* — 109  
*Perilampus* — 102  
*Perilissus* — 70  
*Perilitus* — 44  
*Perineura* — 23  
*Peristerola* — 145  
*Perispincter* — 84  
*Perispuda* — 71  
*Peristenus* — 44  
*Perithous* — 52  
*Perniphora* — 106  
*Petalodes* — 32  
*Pezophycta* — 131  
*Pezilepsis* — 110  
*Phaenocarpa* — 35  
*Phaenolobus* — 84  
*Phaenoserphus* — 136  
*Phaenogenes* — 88  
*Phanacis* — 132  
*Phanerotoma* — 42  
*Phanerotomella* — 43  
*Philanthus* — 166  
*Phobetes* — 73  
*Phobocampe* — 77  
*Pholetesor* — 39  
*Phradis* — 79  
*Phrudus* — 74  
*Phthorima* 86

- Phygadeuon* — 61  
*Phyllocolpa* — 19  
*Phymatocera* — 21  
*Physetopoda* — 151  
*Phytodietus* — 53  
*Pictoscytoides* — 106  
*Piestopleura* — 141  
*Pigeria* — 31  
*Pikonema* — 18  
*Pimpla* — 52  
*Pion* — 70  
*Pirene* — 105  
*Pison* — 169  
*Plagiolepis* — 155  
*Plastonoxus* — 145  
*Platencyrtus* — 113  
*Platneptis* — 110  
*Platycampus* — 18  
*Platygaster* — 141  
*Platygerrhus* — 103  
*Platigonatopus* — 144  
*Platylabops* — 90  
*Platylabus* — 89  
*Platymischus* — 139  
*Platynocheilus* — 119  
*Platypermalus* — 109  
*Platyrhopus* — 112  
*Platystasius* — 141  
*Platytenomus* — 139  
*Plectiscidea* — 85  
*Plectocryptus* — 63  
*Pleolophus* — 63  
*Plesiobaeus* — 140  
*Plesiomicroterys* — 113  
*Pleroneura* — 14  
*Plutothrix* — 103  
*Pnigalio* — 121  
*Podagrion* — 100  
*Podoschistus* — 53  
*Poecilagenia* — 182  
*Poemenia* — 53  
*Pokomandy* — 33  
*Polemochartus* — 35  
*Polistes* — 159  
*Polochrom* — 151  
*Polyaulon* — 59  
*Polyblastus* — 54  
*Polycystus* — 104  
*Polydegmon* — 36  
*Polyergus* — 156  
*Polynema* — 127  
*Polysphincta* — 52  
*Polystenus* — 29  
*Polytribax* — 63  
*Pompilius* — 163  
*Ponera* — 154  
*Pontania* — 20  
*Pontopristia* — 20  
*Poropoea* — 127  
*Praon* — 47  
*Prenanteon* — 144  
*Prenolepis* — 155  
*Prestwichia* — 127  
*Priocnemis* — 162  
*Prionomastix* — 117  
*Prionomitus* — 115  
*Priophorus* — 17  
*Priopoda* 71  
*Pristaulacus* — 134  
*Pristiceros* — 89  
*Pristiphora* — 18  
*Pristocera* — 146  
*Pristomerus* — 79  
*Probles* — 80  
*Probolus* — 89  
*Proceratum* — 154  
*Prochiloneurus* — 116  
*Procinetus* — 84  
*Proclitus* — 85  
*Proconura* — 99  
*Proctotrupes* — 136  
*Profenus* — 21  
*Promethes* — 87  
*Pronotalia* — 124  
*Prosactogaster* — 141  
*Prosapha* — 35  
*Prosevania* — 133  
*Prosopis* — 174  
*Protaganteles* — 40  
*Proaphidius* — 47  
*Protarchus* — 71  
*Proterops* — 43  
*Protichneumon* — 94  
*Protodacnusa* — 35  
*Psen* — 168  
*Psenulus* — 168  
*Pseudaphanomerus* — 141  
*Pseudaphycus* — 114  
*Pseudectroma* — 114  
*Pseudepipona* — 158  
*Pseuderimerus* — 101  
*Pseudeucoila* — 130  
*Pseudisobrachium* — 146  
*Pseudocatolacus* — 109  
*Pseudochrysis* — 148  
*Pseudoclavellaria* — 16  
*Pseudococcobius* — 114  
*Pseudocymodusa* — 77  
*Pseudodineura* — 20  
*Pseudogonatopus* — 144  
*Pseudomicrodynerus* — 158  
*Pseudoplatylabus* — 90  
*Pseudopraon* — 47  
*Pseudorhopus* — 117  
*Pseudorhyssa* — 53  
*Pseudosystole* — 100  
*Pseudotetramesa* — 99  
*Pseudotorymus* — 101  
*Pseudovipio* — 30  
*Psilanteris* — 140  
*Psilocera* — 107  
*Psilomastax* — 94  
*Psilomma* — 136  
*Psilonotus* — 108  
*Psilophrys* — 117  
*Psilus* — 139  
*Psithyrus* — 184  
*Psychophagus* — 107  
*Psyllaephagus* — 115  
*Psyetopoda* — 151  
*Pterocheilus* — 158  
*Pteromalus* — 107  
*Pteroptrix* — 118  
*Pycnomyces* — 64  
*Pygostolus* — 43  
*Pyracmon* — 76  
*Pyramydophorus* — 89
- Q
- Quadrencyrtus* — 117
- R
- Rakosina* — 110  
*Rastvalva* — 40  
*Ratzeburgiola* — 121  
*Rhaconotus* — 28  
*Rhadinoceraea* — 21  
*Rhaestus* — 70  
*Rhaphitelus* — 106  
*Rhembobius* — 60  
*Rhincocoelia* — 104  
*Rhincopelte* — 120  
*Rhimpoctona* — 75  
*Rhinotorus* — 72  
*Rhodanthidium* — 179  
*Rhogogaster* — 23  
*Rhopalicus* — 106  
*Rhopalum* — 169  
*Rhopites* — 178  
*Rhophitoides* — 178  
*Rhoptronctrus* — 28  
*Rhopromeris* — 130  
*Rhopus* — 112  
*Rhorus* — 70  
*Rhynchobanchus* — 69  
*Rhysipolis* — 29  
*Rhyssa* — 53  
*Rhyssalus* — 29  
*Rhyssella* — 53  
*Rhyssolabus* — 89  
*Rictichneumon* — 91  
*Rogas* — 32  
*Rohatina* — 109  
*Ronisia* — 151  
*Ropalophorus* — 44  
*Roptocerus* — 106  
*Rostrobracon* — 31
- S
- Sactogaster* — 141  
*Saotis* — 72  
*Saphonecrus* — 132  
*Sapysa* — 151  
*Sapygina* — 151  
*Sarothrus* — 130  
*Sathon* — 40

- Sathropterus* — 80  
*Scambus* — 50  
*Scelio* — 140  
*Sceliphron* — 166  
*Sceptrothelys* — 107  
*Schenkia* — 63  
*Schizontonus* — 110  
*Schizoprymnus* — 36  
*Schizopyga* — 51  
*Sciateryx* — 23  
*Sclerodermus* — 146  
*Scoila* — 151  
*Scolioneura* — 21  
*Scolobates* — 71  
*Scopesis* — 72  
*Scorpioteleia* — 138  
*Scutellista* — 103  
*Seciliclava* — 115  
*Selanderma* — 104  
*Selandria* — 17  
*Seleucus* — 73  
*Semen* — 115  
*Semiotelus* — 105  
*Sesioplex* — 74  
*Sharliphora* — 19  
*Shawiana* — 29  
*Stifolinia* — 154  
*Sigalphus* — 42  
*Sigmophora* — 124  
*Sinarachna* — 52  
*Sinophorus* — 74  
*Stobla* — 22  
*Strex* — 24  
*Skeloceras* — 105  
*Smicromyrme* — 151  
*Smicroplectrus* — 56  
*Solenopsis* — 139  
*Sollerella* — 168  
*Spalangia* — 102  
*Spalangiotelta* — 102  
*Spaniopus* — 109  
*Sparsion* — 140  
*Spathius* — 28  
*Spathopus* — 105  
*Sphaenolepis* — 122  
*Sphaeripalpus* — 104  
*Sphecodes* — 176  
*Sphecophaga* — 66  
*Sphiegigaster* — 104  
*Sphe* — 166  
*Sphinctus* — 53  
*Spilichneumon* — 93  
*Spilochalcis* — 98  
*Spilomalus* — 107  
*Spilomena* — 169  
*Spilomicrus* — 138  
*Spinolia* — 148  
*Spintharis* — 148  
*Spintherus* — 107  
*Spudaeus* — 83  
*Stauronematus* — 18  
*Stauropoctonus* — 81  
*Stelis* — 179  
*Stemmatosteres* — 114  
*Stenamma* — 155  
*Stenarella* — 66  
*Stenetra* — 107  
*Stenichneuman* — 92  
*Stenobarichneumon* — 91  
*Stenodontus* — 88  
*Stenodynerus* — 158  
*Stenolabus* — 90  
*Stenomacrus* — 86  
*Stenomalina* — 108  
*Stenomesius* — 119  
*Stenophrus* — 105  
*Stenoselma* — 106  
*Stephanodes* — 127  
*Stephanus* — 95  
*Sterictiphora* — 15  
*Stethomostus* — 21  
*Stibeutes* — 61  
*Stichocrepis* — 110  
*Stictomischus* — 105  
*Stictoplisthus* — 82  
*Stigmus* — 169  
*Stilbops* — 66  
*Stilbula* — 102  
*Stilbum* — 148  
*Stilpnus* — 62  
*Stinoplus* — 109  
*Stizus* — 167  
*Strebocera* — 44  
*Strejcekia* — 110  
*Strombocerina* — 16  
*Strongylogaster* — 16  
*Strongylognathus* — 155  
*Subprionomitus* — 113  
*Sulcarius* — 61  
*Sulcopolistes* — 160  
*Sundholmella* — 136  
*Sussaba* — 87  
*Susterai* — 104  
*Sycophila* — 99  
*Symmorphus* — 159  
*Symphera* — 70  
*Symphya* — 36  
*Sympiesis* — 120  
*Synacra* — 136  
*Synaldis* — 35  
*Synaristis* — 141  
*Synbelyta* — 137  
*Syndipnus* — 73  
*Synechocryptus* — 65  
*Synedrus* — 108  
*Synelix* — 36  
*Syneraus* — 132  
*Synodites* — 72  
*Synoecetes* — 70  
*Synomelix* — 72  
*Syntactus* — 70  
*Syntomopus* — 104  
*Syntretus* — 44  
*Syrphophagus* — 113  
*Syrphophilus* — 86  
*Syspasis* — 91  
*Systasis* — 105  
*Systole* — 100  
*Systropha* — 178  
*Syzeuctus* — 68  
  
*T*  
*Tachyagetes* — 163  
*Tachysphex* — 168  
*Tachytes* — 168  
*Tamarixia* — 122  
*Tanycarpa* — 35  
*Taphaeus* — 36  
*Tapinoma* — 155  
*Taxonus* — 22  
*Teleas* — 140  
*Telenomus* — 139  
*Teleopterus* — 126  
*Teleutea* — 66  
*Telostegus* — 164  
*Temelucha* — 79  
*Tenthredo* — 23  
*Tenthredopsis* — 23  
*Termolampa* — 105  
*Tersilochus* — 80  
*Tetracampe* — 119  
*Tetracnemoidea* — 113  
*Tetradontochelys* — 144  
*Tetracnemus* — 112  
*Tetralonia* — 183  
*Tetramesa* — 99  
*Tetramopria* — 139  
*Tetramorium* — 155  
*Tetrastrichus* — 121  
*Thectogaster* — 104  
*Theocolax* — 103  
*Therlon* — 83  
*Theronia* — 52  
*Theroscopus* — 61  
*Thinodytes* — 104  
*Thomsonisca* — 115  
*Thoron* — 140  
*Tripectenoides* — 126  
*Thrybius* — 64  
*Thymaris* — 54  
*Thyrateles* — 92  
*Thyreus* — 183  
*Thysanus* — 118  
*Timaspis* — 132  
*Tiphia* — 150  
*Tiphodytes* — 139  
*Tomocobia* — 110  
*Tomostethus* — 21  
*Torymus* — 101  
*Townesia* — 51  
*Townesilitus* — 44  
*Toxares* — 47  
*Toxeuma* — 104  
*Trachelus* — 25  
*Trachusa* — 179  
*Trachyarus* — 87

- Trachyusa* — 35  
*Tranosema* — 77  
*Tranosemella* — 78  
*Trechnites* — 114  
*Trematopygus* — 69  
*Tremex* — 24  
*Triaspis* — 37  
*Trieces* — 82  
*Triepolus* — 182  
*Trichacis* — 141  
*Trichiocampus* — 17  
*Trichiosoma* — 16  
*Trichogramma* — 127  
*Tricholabus* — 94  
*Tricholinum* — 61  
*Trichomalopsis* — 109  
*Trichomalus* — 109  
*Trichomasthus* — 112  
*Trichomma* — 83  
*Trichoplectrus* — 119  
*Trichiosoma* — 16  
*Trichopria* — 139  
*Trichosteresis* — 141  
*Tricladia* — 113  
*Triclistus* — 82  
*Tricolas* — 107  
*Tricyclomischus* — 104  
*Trigonalis* — 142  
*Trigonaspis* — 131  
*Trigonoderus* — 103  
*Trimorus* — 139  
*Triptognathops* — 94  
*Triptognathus* — 93  
*Trissacantha* — 140  
*Trissolcus* — 139
- Trisynaldis* — 35  
*Tritneptis* — 110  
*Trioxys* — 47  
*Trjapitzinellus* — 116  
*Trogus* — 94  
*Tromatobia* — 51  
*Tropidodynerus* — 158  
*Tropistes* — 61  
*Trybliographa* — 130  
*Trychnosoma* — 108  
*Trychosis* — 64  
*Tryphon* — 55  
*Trypoxylon* — 169  
*Tumidiclava* — 127  
*Tymmophorus* — 86  
*Tyndarichus* — 116
- U**  
*Ufens* — 127  
*Urocerus* — 24  
*Urocyclops* — 141  
*Urolepis* — 110
- V**  
*Veltrusia* — 110  
*Venanides* — 40  
*Venturia* — 75  
*Vespa* — 160  
*Vespula* — 160  
*Viennopria* — 139  
*Vipio* — 38  
*Virgichneumon* — 91  
*Vrestovia* — 111  
*Vulgichneumon* — 91
- W**  
*Wesmaelia* — 43  
*Woldstedtius* — 86
- X**  
*Xanthellum* — 120  
*Xanthoectroma* — 112  
*Xenaphycus* — 114  
*Xenaproctocetus* — 122  
*Xenarcha* — 29  
*Xenolytus* — 58  
*Xenomerus* — 140  
*Xenoschesis* — 69  
*Xeris* — 24  
*Xestonotidea* — 141  
*Xestophanes* — 132  
*Xiphulcus* — 58  
*Xiphydria* — 24  
*Xiphydriophagus* — 106  
*Xorides* — 57  
*Xyela* — 14  
*Xylocopa* — 184  
*Xylophrurus* — 65
- Z**  
*Zabrychypus* — 51  
*Zaglyptus* — 51  
*Zaomma* — 116  
*Zaplethoconia* — 70  
*Zaraea* — 16  
*Zatypota* — 52  
*Zele* — 43  
*Zoophthorus* — 60  
*Zygota* — 137

**Authors' addresses:**

RNDr. Karel Beneš, Petýrkova 1956, 149 00 Praha 4.

Doc. Ing. Miroslav Čapek, DrSc., Výzkumná stanica VÚLH, Lesnícka ul. 11, 969 23 Banská Štiavnica.

Ing. Vladimír Kalina, Ústav aplikované ekologie a ekotechniky, 281 63 Kostelec nad Černými lesy.

Ing. Miroslav Kocourek, Nejedlého ul. 36, 682 03 Vyškov.

RNDr. Jozef Lukáš, CSc., Zool. ústav PFUK, Mlynská dolina, B2, 842 15 Bratislava.

RNDr. Jan Macek, Národní muzeum, ent. odděl. 148 00 Praha - Kunratice 1.

Pavel Mikula, prom. biol., Federální ministerstvo zemědělství a výživy ČSSR, odb. VTR, Těšnov 65, 117 05 Praha.

RNDr. Zdeněk Pádr, CSc., ul. Lidových milicí 43, 120 00 Praha 2.

RNDr. Jaromír Strejček, Mládežnická ul. 3063, 108 00 Praha 10.

RNDr. Josef Šedivý, DrSc., Výzkumný ústav rostlinné výroby, 161 06 Praha 6 - Ruzyně 507.

RNDr. Petr Werner, Výzkumný ústav potravinářského průmyslu, Na bělidle 21, 150 38 Praha 5.

---

Acta faunistica entomologica Musei Nationalis Pragae 19.  
Redaktor RNDr. Jiří Dlabač, CSc. — Vydává Národní muzeum v Praze.  
Vyšlo 30. XI. 1989 — Náklad 1'000 ks.  
Vytiskly: Tiskařské závody, s. p., provoz 61, Kladno.